

# **FOLLOW-UP SURVEY OF NUTRITIONAL STATUS IN CHILDREN 6-29 MONTHS OF AGE, KYRGYZ REPUBLIC 2013**



---

*Disclaimer:*

*The findings and conclusions of this report do not necessarily represent the official position of the U.S. Centers for Disease Control and Prevention and UNICEF.*

*Use of trade names is for identification only and does not imply endorsement by the U.S. Department of Health and Human Services and UNICEF.*

---

## TABLE OF CONTENTS

<b>LIST OF ABBREVIATIONS</b>	<b>4</b>
<b>INVESTIGATORS AND COLLABORATORS</b>	<b>5</b>
<b>ACKNOWLEDGEMENTS</b>	<b>5</b>
<b>EXECUTIVE SUMMARY</b>	<b>7</b>
<b>SURVEY OBJECTIVES AND DESIGN</b>	<b>7</b>
<b>SUMMARY OF FINDINGS</b>	<b>8</b>
<i>Gulazyk MNP Program</i>	8
<i>Infant and Young Child Nutrition (IYCN): Knowledge, Attitudes, and Practices</i>	9
<i>Anthropometry</i>	9
<i>Biochemical Indicators</i>	10
<i>Comparison of Nutritional Status of Children 6–29 months between the 2009 and 2013 National Surveys</i>	10
<b>CHAPTER 1: INTRODUCTION</b>	
BACKGROUND	11
SURVEY OBJECTIVES	12
<b>CHAPTER 2: METHODS</b>	
SURVEY POPULATION	13
<i>Sample Size Determination</i>	13
SAMPLING PROCEDURES	14
ON-SITE TRAINING AND SURVEY IMPLEMENTATION	15
<i>Survey Teams</i>	15
<i>Training</i>	15
<i>Preparation for Field Survey</i>	16
<i>Survey Implementation</i>	16
DATA COLLECTION	17
<i>Survey Instrument</i>	17
<i>Anthropometry</i>	17
<i>Hemoglobin Measurement</i>	17
<i>Blood Collection, Processing and Storage</i>	18
<i>Quality Control and Assurance</i>	18
DATA MANAGEMENT AND STATISTICAL ANALYSIS	19
<b>CHAPTER 3: RESPONSE RATES AND CHARACTERISTICS OF RESPONDENTS</b>	
RESPONSE RATES – 41 CHILDREN WERE OUT OF AGE LIMITS AND 2301 WERE ELIGIBLE TO PARTICIPATE.	20
<b>CHAPTER 4: GULAZYK MICRONUTRIENT POWDER (MNP) PROGRAM AND TRENDS IN GULAZYK USE FROM 2011–2013</b>	
BACKGROUND	26
<i>Communications and Social Mobilization</i>	26
<i>Monitoring Gulazyk use</i>	27
GULAZYK USE IN 2013	28
<i>Caregiver Report of Receipt and Use of Gulazyk</i>	28
<i>Experiences Giving Gulazyk</i>	34
<i>Caretaker’s Perceived Effects of Gulazyk</i>	35

---

<i>Coverage of Gulazyk Communications Materials and Activities</i>	36
TRENDS IN GULAZYK USE FROM 2011–2013	37
<i>Background</i>	37
<i>Methods</i>	37
<i>Trends in Gulazyk use</i>	38

## **CHAPTER 5: INFANT AND YOUNG CHILD NUTRITION (IYCN): KNOWLEDGE, ATTITUDES, AND PRACTICES**

BACKGROUND	47
INFANT AND YOUNG CHILD FEEDING PRACTICES	47
KNOWLEDGE AND ATTITUDES REGARDING INFANT AND YOUNG CHILD FEEDING	50
DIETARY AND BREASTFEEDING ADVICE	51

## **CHAPTER 6: ANTHROPOMETRY**

BACKGROUND	53
METHODS	53
RESULTS	55
DISCUSSION	58

## **CHAPTER 7: BIOCHEMICAL INDICATORS FOR MICRONUTRIENT DEFICIENCY AND THE USE OF MICRONUTRIENT SUPPLEMENTS 59**

BACKGROUND	59
EFFECTS OF INFLAMMATION ON NUTRITIONAL BIOMARKERS	59
IRON DEFICIENCY, ANEMIA, AND IRON DEFICIENCY ANEMIA	59
VITAMIN A DEFICIENCY	63
USE OF MICRONUTRIENT SUPPLEMENTS	65

## **CHAPTER 8: COMPARISON OF NUTRITIONAL STATUS OF CHILDREN 6–29 MONTHS AND INFANT YOUNG CHILD NUTRITION KNOWLEDGE, ATTITUDES, AND PRACTICES — 2009 AND 2013 NATIONAL SURVEYS 66**

BACKGROUND	66
METHODS	66
<i>Biochemical indicators</i>	67
RESULTS	69
<i>Anthropometry</i>	70
<i>Biochemical indicators</i>	71
<i>Vitamin A Supplement Use</i>	75
<i>IYCN Knowledge, Attitudes, and Practices</i>	75

## **DISCUSSION 77**

## **REFERENCES 80**

## **APPENDICES 84**

APPENDIX I: QUESTIONNAIRE FOR 2013 NATIONAL NUTRITION SURVEY	85
APPENDIX II: LIST OF CHILD HEMOGLOBIN RESULTS: FOR VILLAGE MEDICAL ATTENDANT	105
APPENDIX III: CONFIDENCE INTERVALS, DESIGN EFFECTS, AND INTRA-CLASS CORRELATION COEFFICIENTS FOR MAJOR INDICATORS, CHILDREN 6–29 MONTHS — KYRGYZSTAN NATIONAL SURVEY, 2013	106
APPENDIX IV: QUESTIONNAIRES FOR 2011 AND 2012 LOT QUALITY ASSURANCE SAMPLING SURVEYS	108

---

APPENDIX V: ALGORITHMS FOR CALCULATING INDICATORS OF INFANT AND YOUNG CHILD FEEDING PRACTICES	116
APPENDIX VI: NATIONAL SURVEY OF NUTRITIONAL STATUS, KYRGYZSTAN 2013: RESULTS FOR THE WHO GLOBAL DATABASE ON CHILD GROWTH AND MALNUTRITION	122
APPENDIX VII: BRIEF SUMMARY QUALITY ASSURANCE FOR MICRONUTRIENT MEASUREMENTS FROM THE 2009 AND 2013 KYRGYZSTAN SURVEYS	124
APPENDIX VIII: QUESTIONNAIRES FOR 2009 NATIONAL NUTRITION SURVEY	125
APPENDIX IX: COMPARISON OF VARIOUS CHARACTERISTICS AMONG CHILDREN (6-24 MONTHS)— KYRGYZSTAN, 2009 AND 2013	148

---

## LIST OF ABBREVIATIONS

<b>AGP</b>	1-acid glycoprotein
<b>CDC</b>	United States Centers for Disease Control and Prevention
<b>CEE</b>	Central and Eastern Europe
<b>CIS</b>	Commonwealth of Independent States
<b>CRP</b>	C - reactive protein
<b>DHS</b>	Demographic and Health Survey
<b>DDPME</b>	Department for Drugs, Procurement and Medical Equipment
<b>Gulazyk</b>	Local name for micronutrient powder
<b>HAZ</b>	Height-for-age, Z-score
<b>Hb</b>	Hemoglobin
<b>Hgb</b>	Hemoglobin
<b>IYCF</b>	Infant and young child feeding
<b>IYCN</b>	Infant and young child nutrition
<b>KSSHPP</b>	Kyrgyz-Swiss-Swedish Health Project
<b>LQAS</b>	Lot Quality Assurance Sampling
<b>MNP</b>	Micronutrient Powder
<b>MOH</b>	Ministry of Health
<b>NSC</b>	National Statistical Committee
<b>RBP</b>	Retinol-binding protein
<b>sTfR</b>	Soluble transferrin receptor protein
<b>UNICEF</b>	United Nations Children's Fund
<b>VAD</b>	Vitamin A Deficiency
<b>VHC</b>	Village Health Committee
<b>WAZ</b>	Weight-for-age, Z-score
<b>WHO</b>	World Health Organization
<b>WHZ</b>	Weight-for-height, Z-score

---

## INVESTIGATORS AND COLLABORATORS

This survey was administered under the Cooperative Agreement between UNICEF and the U.S. Centers for Disease Control and Prevention. Support for this survey was provided by several investigating and collaborating agencies. These include:

- UNICEF Regional Office CEE/CIS
- UNICEF Kyrgyz Republic (UNICEF Kyr)
- U.S. Centers for Disease Control and Prevention (CDC)
- Ministry of Health, Kyrgyz Republic (MoHKyr)
- National Statistics Committee, Kyrgyz Republic (NSC)

The core members of the survey team are as follows:

- Kyrgyz Republic
  - Cholpon Imanalieva (UNICEF Kyr) – Program Coordinator
  - Muktar Minbaev (UNICEF Kyr) – Survey Coordinator
  - Galina Samohleb (NSC) – Fieldwork Coordinator
  - Tursun Mamyrbayeva (MoH) – MoH and Laboratory Coordinator
  - Larisa Praslova (NSC) – Coordinator Data Collection and Management
  - Asel Ismailova (MoH) – Laboratory Field Supervisor
- United States
  - Mary Serdula (CDC) – Senior Scientist
  - Kevin Sullivan (CDC) – Senior Epidemiologist
  - Jennifer N. Lind (CDC) – Epidemic Intelligence Service Officer
  - Ralph D. Whitehead, Jr. (CDC) – Laboratory Advisor

---

## ACKNOWLEDGEMENTS

We would like to acknowledge that the 2010 Talas Oblast, Kyrgyz Republic Follow-up Survey of Nutritional Status in Children 6–24 months of age Report and the 2009 National Survey of the Nutritional Status of Children 6–59 months of age and Their Mothers Report were used as the basis for the development of this report.

Special thanks to Timothy Schaffter and Jonathan Veitch (Representatives of UNICEF Kyrgyz Republic), Vilma Tyler (Nutrition Specialist at the UNICEF Regional Office for CEE/CIS), and Arnold Timmer (Senior Advisor, Micronutrients, Nutrition Section, UNICEF) for their support of the surveys.

We appreciate leadership and support provided by management of the Ministry of Health and the National Statistics Committee of Kyrgyz Republic. Thank you to M.M Karataev (Deputy Minister of Health of the Kyrgyz Republic) and O.A. Abdykalykov (Head of the National Statistics Committee of Kyrgyz Republic) for their support. Sincerest thanks to U.M. Tilikeeva (Chairman of Ethics Committee, Department for Drugs, Procurement and Medical Equipment [DDPME]) and her team, who provided invaluable input into adapting the protocol to protect research participants.

We would like also to thank Dr. Nancy Aburto, Epidemic Intelligence Service Officer (currently with the World Food Program) for her contribution and leadership to the national 2009 survey and Djudsupova Aijan, Program Assistant, for her invaluable input in the organization, coordinating and carrying out of the field research.

Finally, we are grateful to the interviewers, field supervisors and drivers for their participation in the research, and the citizens of the Kyrgyz Republic for their hospitality and participation in this survey.



---

## EXECUTIVE SUMMARY

This report summarizes the findings of the National Nutrition Survey conducted in the Kyrgyz Republic in July and August 2013, and presents the results for the indicators pertaining to the Gulazyk program, infant and young child nutrition, anthropometry, and biochemical markers of nutritional status. To assess changes in breastfeeding and complementary feeding practices and biochemical indicators of nutritional status following implementation of the nationwide combined Infant and Young Child Nutrition (IYCN)/micronutrient powder (MNP) program, this report also compares data from the 2009 and 2013 National Nutrition surveys. The collaborating partners for these surveys included the Ministry of Health (MOH) of the Kyrgyz Republic, the Kyrgyz Republic National Statistical Committee (NSC), United Nations Children's Fund (UNICEF)-Kyrgyzstan, Kyrgyz-Swiss-Swedish Health Project (KSSHP), and the U.S. Centers for Disease Control and Prevention (CDC).

---

## SURVEY OBJECTIVES AND DESIGN

The major objectives of the 2013 Kyrgyz Republic National Survey of Nutritional Status in Children 6–29 months of age were to:

- Assess receipt and use of Gulazyk by children and assess trends (2011-2013)
- Assess Infant and Young Child Feeding practices
- Assess anthropometric status
- Assess the prevalence of anemia, iron deficiency anemia, iron deficiency, vitamin A deficiency in children 6–29 months of age
- Compare the 2009 and 2013 survey results to measure changes in hematologic and biochemical indicators of iron and vitamin A status after adjustment for infection/inflammation status.

Cross-sectional household surveys were conducted in 2009 and 2013. The 2013 survey was designed to collect comprehensive information on a wide range of topics related to nutritional patterns of children 6–29 months of age living in the Kyrgyz Republic (with the exception of the city of Bishkek)<sup>1</sup>. A pre-post survey design was used to measure changes in micronutrient status between the 2009 and 2013 National Surveys. The 2009 and 2013 surveys used similar methods for sampling and data collection. A two-stage cluster sampling design was employed using probability proportional size (PPS) methodology to select 80 clusters of primary health clinics (FAP/FGP) with 30 children aged 6–29 months in each.

Anthropometric measurements were taken and capillary blood specimens were used to measure biochemical indicators of iron status (serum ferritin and soluble transferrin receptor protein), vitamin A status (retinol-binding protein [RBP]), and inflammation status ( $\alpha$ 1-acid glycoprotein [AGP] and C-reactive protein [CRP]).

In addition to the 2009 and 2013 surveys, Lot Quality Assurance Sampling (LQAS) surveys were conducted in 2011 and 2013. These national surveys (with the exception of Bishkek city) were used to assess the implementation and acceptance of the program.

All data analyses for this report were performed by the National Statistics Center of the Kyrgyz Republic.

---

<sup>1</sup> The city of Bishkek was not included because Gulazyk was not distributed there.

---

## SUMMARY OF FINDINGS

### *Gulazyk MNP Program*

In 2009, the Ministry of Health of the Kyrgyz Republic launched an Infant Young and Child Nutrition (IYCN) program which included promotion of home fortification of complementary foods with a micronutrient powder (MNP) containing iron (12.5 mg elemental iron), vitamin A (300 µg), and other micronutrients. Every 2 months, children aged 6 to 23 months were provided 30 sachets of MNPs to be taken on a flexible schedule.

Nearly all (95.0%) of the caretakers interviewed had received at least one package of Gulazyk. When asked how they obtain Gulazyk, 87.9% of caretakers indicated that they got the product from the local health clinic, 6.5% said a health care worker brought Gulazyk to their home, and 5.4% said they obtained Gulazyk through both clinic visits and home visits. The majority of caretakers (73.5%) reported that their child was “currently” taking Gulazyk. When asked about the schedule they use to give Gulazyk to their children, most (45.3%) caretakers indicated that they gave Gulazyk every other day. On average, caretakers reported that they gave 28.2 sachets of Gulazyk to their children in the last two months. The majority of caretakers (80.1%) indicated that they would continue to give Gulazyk when they finished their current package.

Among all children who received Gulazyk, 55.7% of children consumed at least 30 sachets of Gulazyk (one package contains 30 sachets) during the last two months. Among children who were not currently consuming, the most commonly cited reason for not consuming Gulazyk was the child does not like the food when Gulazyk is added (49.9%).

The majority of caretakers (56.4%) indicated that they noticed changes in the color of food to which Gulazyk was added. Among those who noticed a change in color, 67.0% responded that the change in color was not a concern for them or their child. Additionally, 57.7% of caretakers reported that they noticed changes in the taste of food when Gulazyk was added. Among those who noticed a change in taste, 60.1% responded that the change in taste was not a concern for them or their child. When asked about the portion of food into which Gulazyk was mixed that the child consumes, 66.4% said their child consumed the entire portion and 33.6% said their child consumed less than a full portion.

When caretakers were asked if they noticed any positive changes in their children after they started taking Gulazyk, 42.7% had observed positive changes. Among these, 42.6% indicated increased appetite, 33.6% reported that their child overall seems better/healthier, 29.9% of caretakers indicated that the energy level increased, 16.1% indicated more curiosity/intelligence, 10.4% reported that their child gets sick less often, and 9.0% indicated better growth.

Sixty percent of all caretakers reported receiving the Gulazyk brochure. Only 43.0% of respondents had a radio at home; of those 45.6% indicated that they had heard about Gulazyk on the radio. Nearly all caretakers (99.2%) reported having a television; of those 68.8% of respondents indicated that they had heard about Gulazyk on television.

Between 2011 and 2013, clinic records (Green Journal) indicated that the percentage of children

---

who had ever received Gulazyk changed very little in 2011, 2012 and 2013 (93.2%, 90.8% and 92.1%). However, between 2011 and 2013, those who had received Gulazyk within 3 months of the interview decreased significantly from 84.9% to 76.1%. According to caretaker report, the percentage of children who had ever received Gulazyk MNP was about 95% in all three years (95.3%, 95.9% and 95.0%, respectively). The percentage children who were currently receiving Gulazyk decreased was 79.4%, 69.7% and 73.5%, respectively. The percentage of caretakers who reported that their child<sup>2</sup> would continue in the program declined (89.1%, 84.6% and 81.0%, respectively).

### ***Infant and Young Child Nutrition (IYCN): Knowledge, Attitudes, and Practices***

For the first 6 months of age WHO recommends exclusive breastfeeding; at six months of age WHO recommends introduction of solid, semi-solid and soft foods to supplement the breastfed child's diet (WHO, 2001). Appropriate infant and young child feeding (IYCF) practices include altering the frequency, variety, and amount of foods as a child gets older, while continuing breastfeeding until 2 years of age. Among children 6.0 to 23.9 months of age, 99.5% were reported to have ever been breastfed and 40.2% were reported to have been exclusively breastfed for the first six months after childbirth. At 1 year of age, 80.1% were reported to be breastfeeding and at 2 years of age, 26.2% were reported to be breastfeeding. The proportion of infants 6.0–8.9 months of age who were reported to have received appropriate solid, semi-solid or soft foods was 90.9%. The proportion of children 6.0–23.9 months who received a diet with the minimum recommended diversity (4 or more food groups) was 86.8% and the proportion who received a diet with at least the minimum recommended meal frequency was 74.7%.

Among caretakers, 75.6% considered breastfeeding very important for the baby's health and nutrition. Nearly all caretakers (99.6%) reported that babies should be breastfed and 99.7% reported that there were advantages to breastfeeding. Among those who felt a baby should be breastfed, the mean length of time reported that a baby should breastfeed was 23.9 months.

### ***Anthropometry***

A total of 2162 children aged 6.0-29.9 months had weight and age data to calculate length-for-age (stunting), weight-for-length (wasting), and weight-for-age (underweight) z-scores. The 2013 survey found that 11.7% of the children 6.0–29.9 months of age had growth stunting, 2.0% had wasting, 4.8% were underweight, and 3.2% of children were overweight. The prevalence of stunting increased across age through 23.9 months of age, was similar among rural and urban children (11.3 versus 12.6%), and was slightly higher among boys compared to girls (13.1% versus 10.3%).

### ***Biochemical Indicators***

A total of 2156 children had hemoglobin, serum ferritin, and soluble transferrin receptor (sTfR) measurements and 2148 children had retinol binding protein (RBP) measurements. Among all children 6.0–29.9 months of age, 34.2% had iron deficiency as measured by ferritin and 39.3% had iron deficiency as measured by sTfR. After adjustment for altitude, the prevalence of anemia among all children was 32.7%. The prevalence of iron deficiency anemia as measured by ferritin was 18.8% and as measured by sTfR was 21.5%.

When the prevalence of anemia among all children was stratified by age, gender, and place of residence, the prevalence of anemia was higher in children 6 to 23 months (35.8% to 35.0%) compared to children 24.0–29.9 months (24.5%). The prevalence of anemia was 34.5% in males compared to 30.7% in females and was similar in rural and urban areas.

A total of 2148 children had RBP measurements. The prevalence of vitamin A deficiency (VAD) was 15.6%. Among children 6.0-11.9, 12.0-17.9, 18.0-23.9, and 24.0-29.9 months, the prevalence of VAD was

---

<sup>2</sup> Among children who were not aging out of the program prior to running out of sachets.

---

16.0%, 13.7%, 16.1%, and 16.7%, respectively. The prevalence of deficiency was 17.0% among males and 14.1% among females, and was 16.0% in urban and 14.7% in rural areas.

### ***Comparison of Nutritional Status of Children 6–29 months between the 2009 and 2013 National Surveys***

Among children 6 to 29 months (not living in Bishkek), the prevalence of anemia showed a non-significant decline of 6.0 percentage points [PP], from 38.7% at baseline in 2009 to 32.7% at follow-up ( $p=0.116$ ). The prevalence of iron deficiency (as measured by serum ferritin) decreased by 16.4 PP (50.6% vs 34.2%,  $p<0.001$ ); iron deficiency (as measured by sTfR) prevalence decreased by 9.6 PP (48.9% vs 39.3%,  $p=0.007$ ); and iron deficiency anemia prevalence decreased by 8.3% PP (from 31.9% vs 23.6%,  $p=0.014$ ). Among all children, the prevalence of vitamin A deficiency increased by 9.4 PP (6.2% vs 15.6%,  $p<0.001$ ). Between the two surveys, the prevalence of wasting among children 6-29 months remained stable ( $\leq 2\%$ ), while the prevalence of stunting decreased by 7.9 PP from 19.6% to 11.7%,  $p<0.001$ ). IYCN indicators showed either no change or showed improvement.

---

## CHAPTER 1 INTRODUCTION

### BACKGROUND

Anemia is a public health problem in the Kyrgyz Republic. In a 2009 national baseline survey of nutritional status of children 6–59 months of age and their mothers, the prevalence of anemia was 26.0% in children and 23.0% in their mothers (CDC/UNICEF/MOH, 2012). Based on the prevalence in children, anemia constitutes a public health problem of moderate significance according to the World Health Organization (WHO) classification (WHO 2001). To address micronutrient deficiency, in May 2008, the Ministry of Health of Kyrgyzstan (MOH) — in close collaboration with the United Nations Children’s Fund (UNICEF)-Kyrgyzstan and Kyrgyz-Swiss-Swedish Health Project (KSSHP)—developed a nationwide Infant and Young Child Nutrition (IYCN) program, which included a nutrition education program to encourage breastfeeding and appropriate complementary feeding. The program is community-based and relies on active involvement of medical specialists from primary health care level facilities and village health committees (VHCs). The educational campaign uses the infrastructure of the VHC for the dissemination of educational messages to mothers living in rural areas. The VHCs consist of volunteers (mainly women) who learn about specific public health issues and ways to address the issues. The VHCs then share their knowledge with women through peer-to-peer interactions.

In addition to the nationwide IYCN program, in June 2009, the MOH began a pilot program in Talas Oblast, to distribute micronutrient powders (MNP), known locally as Gulazyk, through government primary health care centers, at no cost, to all children 6–23 months of age. The MNP contains iron (12.5 mg elemental iron), vitamin A (300 µg), and other micronutrients. Every 2 months, children aged 6 to 23 months were provided 30 sachets of MNPs to be taken on a flexible schedule. Primary health care providers distributed MNP, instructed caretakers on proper use, and distributed informational materials. A pre-post survey design was used in Talas Oblast to measure change in nutritional status of children before and after implementation of the IYCN/MNP program. A baseline survey was conducted in rural Talas Oblast in June 2008 and a follow-up survey was conducted in July and August 2010. Declines were observed in the prevalence of: anemia, 50.6% versus 43.8% ( $p=0.05$ ); total iron deficiency (either low ferritin or high sTfR), 77.3% versus 63.7% ( $p<0.01$ ); and iron deficiency anemia, 45.5% versus 33.4% ( $p < 0.01$ ). In addition, the 2010 survey found, 66.4% of children were reported to currently consume the MNP.

After the Talas pilot program showed positive results, the IYCN/MNP program was gradually scaled up and had become nationwide (with the exception of the city of Bishkek) by June 2011. In 2011 and 2012, surveys using Lot Quality Assurance Sampling (LQAS) were conducted to assess implementation and acceptance of the nationwide program. LQAS survey results showed nationwide MNP coverage (defined as the caretaker report of consumption of MNP by the child at the time of the survey) to be 76.0% in 2011 and 66.8% in 2012.

A national baseline survey of nutritional status of children 6–59 months of age in the Kyrgyz Republic, and their mothers, was conducted in June and July 2009. In order to measure change in micronutrient status and IYCN practices, a follow-up national survey was conducted in July and August 2013 of children 6–29 months of age.

---

## **SURVEY OBJECTIVES**

The major objectives of the 2013 Kyrgyz Republic National Survey of Nutritional Status in Children 6–29 months of age were to:

- Assess receipt and use of Gulazyk by children and assess trends (2011-2013)
- Assess Infant and Young Child Feeding practices
- Assess anthropometric status
- Assess the prevalence of anemia, iron deficiency anemia, iron deficiency, vitamin A deficiency in children 6–29 months of age
- Compare the 2009 and 2013 survey results to measure changes in hematologic and biochemical indicators of iron and vitamin A status after adjustment for infection/inflammation status.

---

## CHAPTER 2 METHODS

### SURVEY POPULATION

The target demographic for this survey was children 6–29 months of age living in the Kyrgyz Republic (with the exception of the city of Bishkek). Although Gulazyk was only distributed to children 6–23 months of age, we anticipated some micronutrient effects for an additional 6 months (Ip 2009). Children outside of the age range and children whose families moved outside of the designated localities were excluded. In addition, the metropolitan area of Bishkek was excluded from sampling and analysis since the Gulazyk program is not active in the area.

### Sample Size Determination

The objective of the 2013 National Nutrition Survey was to describe the nutritional status of the target population of the MNP program based on a number of indicators including anemia, iron deficiency, iron deficiency anemia, and vitamin A deficiency in children 6-29 months of age. We calculated the necessary sample size in order to be able to detect statistically significant differences from results of the 2009 National Nutrition Survey. We took into account the effects detected in serum biomarkers of iron deficiency and anemia in the pilot study of Gulazyk in Talas in 2010. We calculated the sample sizes for this follow-up study using the following equation:

$$\text{Power} = \Phi \left[ \frac{|p_2 - p_1|}{\sqrt{p_1 q_1 / e_{n_1} + p_2 q_2 / e_{n_2}}} - t_{1 - \frac{\alpha}{2}, m_1 + m_2 - 1} \frac{\sqrt{\bar{p} \bar{q} (1/e_{n_1} + 1/e_{n_2})}}{\sqrt{p_1 q_1 / e_{n_1} + p_2 q_2 / e_{n_2}}} \right]$$

Where:

p1= prevalence/coverage in the baseline survey

q1= 1-p1

n1= sample size in baseline survey

DEFF1= design effect in baseline survey

e<sub>n1</sub>= equivalent sample size in baseline survey calculated as n1/DEFF1

p2= estimated prevalence/coverage in follow-up survey

q2= 1-p2

n2= sample size in follow-up survey

DEFF2= the design effect in the follow-up survey; this is estimated from the baseline survey by:

a) calculating the average number of observations per cluster in the baseline survey (k1) as n1/m1, where m1= number of clusters in baseline survey;

b) calculating ICC1 for the baseline survey as (DEFF1-1)/(k1-1);

c) calculating average number of observations in the follow up survey as k2=n2/m2;

d) finally, calculating the DEFF2 as 1+(k2-1)×ICC1

e<sub>n2</sub>= equivalent sample size in follow-up survey calculated as n2/DEFF2



---

$p = \frac{p_1 \cdot n_1 + p_2 \cdot n_2}{n_1 + n_2}$

$q = 1 - p$

Based on previous survey data from Talas, the sample was designed to detect a 7.7 percentage point change in ferritin deficiency from the baseline of 52% with a power of 80% and a two-sided alpha of 0.05. This also allowed for the detection of a change in elevated sTfR of 9.1 percentage points or more, and a change in the prevalence of iron deficiency anemia (where iron deficiency is defined as low ferritin and/or high sTfR) of 8.1 percentage points. All of these detectable changes were smaller than those detected in the 2008-2010 Talas pilot study (CDC/UNICEF/MOH, 2011).

The sample was nationally representative, consisting of 80 clusters of 30 children each (one cluster had 22 children instead of 30). An anticipated response rate of 85% (based on previous studies) was factored into the sample size calculation.

## SAMPLING PROCEDURES

The Republican Health Information Center provided a list of health clinics (known in Kyrgyzstan as FAPs or FGPs) with the number of children assigned to each FAP/FGP. FAP/FGPs were the Primary Sampling Units (PSUs) in this study. (There were 1,599 clinics (FAP and FGP's) in 2013.) The first stage sampling of 80 PSUs was performed using probability proportional to size (PPS). In this case the probability of PSU selection depended on the number of children aged 6-29 months assigned to the PSU. The methodology (MI/CDC, 2007) followed the steps outlined below:

1. At the first step, we made a list with three columns. The first column had the name of sample unit (FAPs/FGPs), listed in geographical order. The second column had the population size. The third column had the cumulative population calculated by summing the population of each sampling unit to the sampling units preceding it on the list. The last cell of the third column was equal to the total size of population.
2. PSUs were selected by calculating a sampling interval ( $k$ ) which equaled the total population size divided by the number of PSUs being surveyed (in this case 80). A random number between 1 and  $k$  was chosen as the starting point.  $k$  was added cumulatively until the number of clusters desired (in this case 80) was chosen.
3. The first PSU was selected as the PSU which contained the cumulative value (column 3) equal to the random starting point. We then chose the second chosen PSU as that with the cumulative value (column 3) equal to the random starting point + the interval ( $k$ ). The third PSU contained the cumulative value (column 3) equal to the random starting number +  $2k$ . This selection process was repeated until we had 80 PSUs.

Before the beginning of the second stage of sampling, we gathered a list from every selected FAP/FGP of all children born during the period from February 1, 2011 through January 31, 2013. We randomly selected 30 children (one cluster had 22 and another cluster had 29 children instead of 30 children) from the lists from every selected FAP/FGP. Before fieldwork in each PSU, the health care staff members at each FAP/FGP informed all selected children of the day and time they should visit the FAP/FGP for participation in the study. Participation was staggered to prevent a situation where too many children were at the clinic at the same time.

Survey team personnel visited the home of any child on the list that did not arrive at the health clinic as scheduled. All efforts were made to measure all of the participants on the list. No substitutions were made for any reason.

---

## ON-SITE TRAINING AND SURVEY IMPLEMENTATION

### *Survey Teams*

Four field teams collected data for the follow-up survey. Each team consisted of one field supervisor, two anthropometrists/interviewers, one phlebotomist, and one driver. The survey coordinator was responsible for overseeing the implementation of the survey (including ensuring that the timeline was followed) and directly coordinating and overseeing the work of the three major study coordinators (fieldwork coordinator, laboratory coordinator and data management coordinator).

### *Training*

All survey team members received one day of instructions regarding the overall survey objectives and procedures. At the close of training all team members participated in a one-day pilot test (10–20 children per survey team) of all survey procedures including blood collection, centrifugation, and storage. The training included presentations and practical demonstrations on the following topics:

All survey team members

- The consent process and confidentiality issues
- Survey objectives
- An overview of interview techniques

The phlebotomist and supervisors

- Procedures for collecting blood samples
- Handling and shipping of biochemical specimens
- Labeling of biological specimens
- Filling in values on questionnaires
- Universal precautions for handling biological specimen
- Protocol for action in case of exposure to biological specimen

The anthropometrists/interviewers and supervisors

- Reliability and validity issues in collecting information using a questionnaire
- The rationale for each question of the survey
- Anthropometry (measuring height and weight of children)

Immediately after the training sessions, the questionnaire was pilot tested in FGPs not selected for the survey (FGPs# 2–5 in the Ak-Ordo and Archa-Beshik areas). The pilot included 59 children. The interviewers conducted the questionnaire and took anthropometric measurements of the children. The phlebotomists obtained blood samples and the laboratory technicians processed the blood samples. The pilot clarified questions regarding the questionnaire as well as other complex logistical issues that arose in the field.

### *Preparation for Field Survey*

The Ethics Committee under the Department of Drug Provision and Medical Equipment approved the survey protocol, biochemical testing procedures, and transportation itinerary to maintain the cold chain for specimen transport. After approval of the research protocol by the Ethics Committee, an order of the Ministry of Health of Kyrgyz Republic, #425, dated July 23, 2013, outlined the respon-

---

sibilities and obligations of health care facilities in the survey.

Before beginning any fieldwork, the Survey Coordinator in conjunction with the Laboratory Coordinator obtained a letter signed and containing the seal of the MOH providing an introduction to the survey with background, justification and explanation of methodology. The letter contained the contact information for the Survey Coordinator in case there were any questions. Each supervisor also had a copy of the letter. Before the survey began, field supervisors visited the local health organizations in each cluster included in the survey to explain the purpose of the survey. They answered any pending questions, received permission to conduct the survey, and obtained the list of eligible children in each cluster. In each cluster, the clinic heads were provided a schedule of the survey in advance.

Before the survey began, the Fieldwork Coordinator prepared a list of children selected for inclusion to the survey. The list of participants was given to the survey team supervisor and the health worker at the clinic. The health care workers were asked to invite those children and their mothers to the health clinic on the pre-determined day. Survey respondents were notified two to three days in advance about the date and time of their FAP visit for the questionnaire interview, anthropometrical measurements, and blood sampling collection before the arrival of the survey team. The supervisor of each team received a route schedule, cluster numbers and addresses, and location of survey administration.

### **Survey Implementation**

Data collection started on July 31, 2013 and was completed on August 26, 2013. Each team surveyed one health clinic (PSU) each day. The survey team arrived at the clinic approximately 30-60 minutes before the arrival of the first child in order to set up the equipment, prepare questionnaires, and introduce themselves to the medical personnel at the clinic. Upon the arrival of the mother or caretaker at the medical clinic, the team supervisor or another team member explained the purpose of the survey and the methods and procedures, and asked the caregiver for consent for the child.

All children from the predetermined participant list were included in the survey if consent was received. No substitutions from the original randomly selected participant list of children were made. If the selected child did not arrive, the medical worker and a member of the survey team visited the home and invited the child to the health clinic. Transportation to the clinic for these participants was provided if necessary. In the case of refusal, the reason was noted on the household questionnaire.

## **DATA COLLECTION**

### **Survey Instrument**

Once consent was given, one of the interviewers administered the questionnaire. The questionnaire contained modules on: receipt and use of Gulazyk; socio-demographic information; breastfeeding and infant feeding patterns; knowledge, attitude and behaviors regarding breastfeeding and infant feeding; dietary advice; vitamin/supplement use; anthropometric measurements; and blood specimen collection. The questionnaire was written in English and then translated into Kyrgyz and Russian languages (Appendix I).

### **Anthropometry**

Length/height and weight measurements were taken for all children 6–29 months of age included in the survey. The age of the child was calculated based on the difference between the child's birth date and the date of the measurement. For children less than 24 months of age, supine length (lying down) was measured to the nearest 0.1 cm. For children 24–29 months of age, height (standing up) was measured to the nearest 0.1 cm using the same height board. All subjects were measured without shoes or hair accessories because they add artificial height. Except for the first day of measurement when Seca boards were used, all measurement were conducted using a height board manufactured by Shorr

---

Productions (Olney, Maryland, USA).

UNICEF Seca Uniscales were used to measure the body weight of the children. The weight of the children was measured by taring the scale after the mother's weight was recorded, then handing the child to the mother. The child's weight was taken with minimal clothing and without a diaper. The results were recorded on the data collection form.

### **Hemoglobin Measurement**

Hemoglobin (Hgb) was assessed in the clinic using the HemoCue™ 301 photometer. The HemoCue™ 301 instrument has an internal quality control (self-test) and does not have a separate control cuvette or need liquid controls. Laboratory personnel collected capillary blood samples through a finger stick using a retractable lancet. In both the 2009 and 2013 surveys, after the first and second drop, the finger was then wiped clean and the third drop was drawn into a HemoCue cuvette.

All mothers participating in the survey were told the Hgb concentration of their child. The results of the Hgb concentration were also given to the mother in written form. Anyone with an Hgb concentration less than 7.0 g/dL received a referral to return to the medical clinic to see the health care worker. The team supervisor recorded hemoglobin results for all children in each cluster on the "List of Child Hemoglobin Results: For Village Medical Attendant" (Appendix II). The team supervisor gave this form to the medical attendant at the end of each day.

### **Blood Collection, Processing and Storage**

For each participant, capillary blood was collected from a fingerstick, using a lancet, into a Microtainer™ containing the anticoagulant EDTA. To avoid hemolysis of specimens, precautions were taken, such as not milking the finger, not scraping the finger with the Microtainer™ and not allowing specimens to touch the frozen gel packs in the cool box. The phlebotomist used a lancet to prick the ring finger and wiped the first two drops of blood away with an alcohol pad, then collected approximately 250-500 µL of blood directly into a Microtainer™ containing EDTA. The blood was mixed well by gently inverting the Microtainer™ about 10 times. The blood in the Microtainer™ was used to assess Hgb, ferritin, soluble transferrin receptor protein (sTfR), C-reactive protein (CRP), α1-acid glycoprotein (AGP), and retinol-binding protein (RBP).

During all data collection and before leaving the clinic at the end of the day, the survey team ensured that all waste had been put in bio-hazard bags and that lancets were disposed of in the sharps collector.

The capillary blood collected into the Microtainer® containing EDTA was immediately placed by the phlebotomists into a cold box containing frozen gel packs placed at the bottom and were maintained at a temperature between 4-10 °C. Blood samples were transported from the field to the mobile laboratory for processing. At the mobile laboratory, the blood was centrifuged the same day that it was received, and plasma was transferred into labeled 0.2mL PCR tubes and cryovials. After centrifugation and preparation, cryovials were stored in a -20°C freezer at the mobile laboratory until transported to Bishkek. Samples were packed in frozen gel packs and transported to Bishkek in the freezer unit of a refrigerator truck. The temperature within the storage unit declined to -20°C within 3-4 hours of driving. In Bishkek, samples were stored at a temperature of -20°C. Finally, the cryovial boxes containing the 0.2mL PCR tubes were shipped on dry ice to VitA-Iron Tech in Willstaett, Germany for analysis of biochemical indicators.

Analyses of biochemical indicators of iron status (ferritin and sTfR), vitamin A status (retinol binding protein [RBP]) and markers of acute inflammation (CRP and AGP) in blood plasma were conducted by the VitA-Iron Tech research laboratory (Erhardt, 2004). The Department of Drug Provision and Medical Equipment under the Ministry of Health of Kyrgyz Republic granted permission to transport specimens to Germany. For an in-depth discussion of the selection of these indicators and the anticipated effects of inflammation on serum values, please see Chapter 7: Biochemical indicators for micronutrient deficiency and the use of micronutrient supplements.

---

## Quality Control and Assurance

### *External Quality Assurance*

The VitMin Laboratory (Willstaett, Germany) has participated in CDC's external quality assurance program, VITAL-EQA (The Vitamin A Laboratory-External Quality Assurance), since 2006 (Haynes, 2008). The laboratory measures ferritin, sTfR, CRP, and RBP concentrations in plasma using an enzyme-linked immunosorbent assay (ELISA) technique (Erhardt, 2004). The precision and bias were Optimal or Desirable for ferritin, sTfR, and CRP (>90% precision of the VITAL-EQA results, with <0.5% bias). However, the precision and bias shifted 15-20% for RBP (>80-85% precision of the VITAL-EQA results, with 18.3% bias) due to a change in pools used by the VITAL-EQA program. This quality assurance analysis is based on exercises immediately preceding and during the survey (Rounds 20-21).

### *Internal Quality Control*

The VitMin Lab analyzed the survey samples for ferritin, sTfR, CRP, RBP, and AGP using an ELISA technique. The lab routinely tested a single QC pool in 10 different wells randomly distributed in each 384-well plate. The inter-assay coefficients variation (CV) for these analytes were 3.8% for RBP, 3.2% for ferritin, 5.1% for AGP, 3.0% for TfR, and 5.2% for CRP. A CV of about 10% provides acceptable precision using an ELISA technique (Erhardt, 2004; Haynes, 2008). These data indicate that the lab's performance exceeded the acceptable performance expectations while analyzing the survey samples.

## DATA MANAGEMENT AND STATISTICAL ANALYSIS

The data were entered into computers using CPro software. Data entry was in duplicate to ensure quality and consistency of data, and any inconsistencies found between the two entries were investigated and resolved. Data analysis was conducted using SPSS (versions 17 and 19). To calculate the age of each child the child's date of birth was subtracted from the interview date and then divided by 30.40. For both survey years, children younger than 6.0 months of age or older than 29.9 months of age at the time of the interview were excluded.

Confidence intervals and design effects (DEFF) for key variables were calculated based on sampling methods. Probability proportionate to size (PPS) sampling was used, so the data were self-weighted. However, it was necessary to weight the data to account for non-response. The 95% confidence intervals were adjusted to account for the cluster survey design. Bivariate tests of statistical significance were conducted using tests (primarily the Pearson chi-square test), which accounted for the cluster survey design. Design effects, and intra-class correlation coefficients for major indicators can be found in Appendix III.

## CHAPTER 3

# RESPONSE RATES AND CHARACTERISTICS OF RESPONDENTS

### RESPONSE RATES

Figure 3-1 and Table 3-1 show participation in this survey. A total of 2391 children were invited to participate (30 children from each of the 78 clusters, one cluster only had 22 children and one had 29 children) (Figure 3-1). As instructions were to interview all children born between February 1, 2011 and January 31, 2013, 41 children were ineligible for analysis because they were younger than 6.0 months of age or older than 29.9 months of age at the time of the interview and 49 children had moved away from the area permanently; this leaves a total of 2301 children that were eligible to participate. Of these, interviews were not obtained for 138 of the children. Interviews were completed for the remaining 2163 children. Of the children from whom interviews were completed, a blood sample that was sufficient for biochemical analysis was obtained from 2156 (7 children had an insufficient blood sample); of these, a blood sample that was sufficient for analysis of hemoglobin was obtained from 2153 children (3 children had insufficient samples for hemoglobin).

**Figure 3-1: Flow chart of participation in survey – Kyrgyzstan, 2013**




Table 3-1 shows characteristics of the respondents for interviews that were obtained. Of interviews that were obtained, the majority of respondents were mothers (84.3%), with some grandmothers (8.4%) and aunts (3.6%) participating. The majority (74.5%) of the caretakers interviewed lived in rural areas, while 25.5% of respondents lived in urban areas.

**Table 3-1: Results of interviews and characteristics of respondents (unweighted) — Kyrgyzstan, 2013**

Indicator	N	%
<b>Result of interview<sup>a</sup></b>	<b>2301</b>	
Interview obtained		94.0
Interview not obtained		6.0
<b>Reasons interview not obtained</b>		
Refused		0.8
Mother sick		0.1
Child sick		0.3
Mother had to work		0.1
Temporary migration		3.7
Other		1.0
<b>Person interviewed (in relation to child)</b>	<b>2163</b>	
Mother		84.3
Grandmother		8.4
Aunt		3.6
Other		3.7
<b>Place of Residence</b>	<b>2163</b>	
Urban		25.5
Rural		74.5

<sup>a</sup> 41 children not included in analysis due to age out of range; 49 children not included due to having moved permanently.

## DEMOGRAPHIC AND SOCIOECONOMIC CHARACTERISTICS

Demographic and socioeconomic characteristics of surveyed children and their mothers are shown in Tables 3-2 and 3-3. The majority (83.3%) of mothers were under 35 years old. All children were between the ages of 6.0 and 29.9 months, and there was fairly equal participation across the age categories for children. The majority (81.1%) of interviews were conducted in the Kyrgyz language.

The majority (56.3%) of the mothers of surveyed children had completed secondary school, with some going on to technical (11.9%) or higher (21.3%) education (Table 3-4). Most of the mothers were not employed (95.7%). Respondents were asked if their family received universal monthly benefits as a proxy for economic status, with 70.2% reporting that they did.

**Table 3-2 Demographic characteristics of children and their mothers (unweighted) — Kyrgyzstan, 2013**

Characteristic	N	%
Age of mothers	2163	
17-24 years		28.3
25-35 years		55.0
35+ years		16.7
Age of children <sup>a</sup>	2163	
6.0-11.9 months		24.0
12.0-17.9 months		26.3
18.0-23.9 months		24.9
24.0-29.9 months		24.7
Gender of children	2163	
Male		51.0
Female		49.0
Place of Residence	2163	
Urban		25.5
Rural		74.5
Language of interview	2163	
Russian		18.5
Kyrgyz		81.1
Uzbek		0.3

<sup>a</sup> Child's age (in months) was truncated to the integer (representing whole months completed)

**Table 3-3 Socioeconomic characteristics of respondents (unweighted) — Kyrgyzstan, 2013**

Characteristic	N	%
Mother's education (highest level completed) <sup>a</sup>	1823	
Less than secondary		10.5
Secondary		56.3
Technical		11.9
Higher		21.3
Mother's employment status <sup>1</sup>	1823	
Not employed		95.7
Employed / student:		4.3
Family receives universal monthly benefits	2163	
Yes		70.2
No		29.4
Don't know		0.4

<sup>a</sup>Education and employment status were asked only of mothers who were interviewed (n=1823)



---

## CHAPTER 4

# GULAZYK MICRONUTRIENT POWDER (MNP) PROGRAM AND TRENDS IN GULAZYK USE FROM 2011–2013

### BACKGROUND

Through the governmental primary health care (PHC) system in Kyrgyzstan, caretakers of children 6.0–23.9 months old are given, free of charge, one package of 30 Gulazyk sachets every two months. The recommended dosing regimen is to use 30 sachets every two months. Therefore, a child should consume a total of 270 sachets within an 18-month period (starting at 6 months of age when the child begins consuming weaning foods and ending at 24 months). Using a “flexible administration” approach, caretakers are instructed that they can give the Gulazyk according to any schedule they wish, as long as they use all

30 sachets in two months and they do not give more than one sachet per day. Gulazyk was distributed to all areas of the country except for the capital city of Bishkek where children were considered to be at lower risk of micronutrient malnutrition.

### *Communications and Social Mobilization*

There are three main communication channels for the Gulazyk program: primary health care providers, village health committee volunteers, and mass media (CDC 2011).

#### *Primary Health Care Providers*

Primary health care providers (doctors, nurses, and feldshers [nurse practitioners]) are the main conduit through which mothers receive information and counseling on the use of Gulazyk. As the sole distributors of Gulazyk and a trusted source of reliable health information at the community-level, primary health care workers serve as a valuable channel through which Gulazyk use can be promoted in both rural and urban settings. Primary health care providers have been extensively trained on the distribution procedures for Gulazyk and appropriate counseling for caretakers on the use of this product. At the clinic, caretakers receive a Gulazyk flier with usage instructions, as well as a reminder card, which states the date on which they should return for their next package of Gulazyk. Furthermore, a record book is kept at each FAP/FGP to record Gulazyk distribution.

#### *Village Health Committees*

Village Health Committees (VHCs) play an integral role in the interpersonal communication to promote the use of Gulazyk among rural populations. Village Health Committees (which work in nearly 1,500 villages throughout the country) make home visits to families with children 6.0–23.9 months of age in the village to inform and educate mothers about Gulazyk, and encourage them to visit their local health clinic to receive Gulazyk for their children. Members of the Gulazyk Action Group within each Village Health Committee are trained to identify families with children in the target age range, discuss Gulazyk with caretakers, counsel caretakers on its use, and monitor adherence to the intervention. Throughout the program, VHC members refer families to the local health clinic for Gulazyk when a child in the family turns six months of age. In this way, VHCs are a critical channel through which to inform families of the availability of Gulazyk in their village, and encourage them to obtain Gulazyk at the local health clinic. VHC volunteers use illustrated flipcharts and brochures to aid them in discussing key messages with caretakers. Additionally, they provide caretakers with a Gula-

---

zyk-branded children's book, which they can read to their children. VHC volunteers recruit engaged and enthusiastic mothers within the village to serve as "Mother Activists" who help promote and monitor the use of Gulazyk.

### *Mass Media*

Mass media is an important communications conduit for the Gulazyk program in both rural and urban areas. In urban areas (where VHC's are not operating), radio broadcasts are the primary mechanism to encourage caretakers to visit the clinic to receive Gulazyk. Print and radio personnel were invited to attend program advocacy meetings, as well as the campaign kick-off event, and journalists were provided with press kits containing information on the program. During and after the scale-up, radio and television broadcasts were aired nationally 3 to 4 times each year. Local (provincial level) radio broadcasts were aired 1 to 2 times each year.

### *Monitoring Gulazyk use*

Records are kept at each health center (referred to in this report as FAP [Feldsher Obstetrics (Accoucher) Points] and FGP [Family Group Practitioners]) in a book known as the Green Journal. In the Green Journal, local staff record which children receive disbursements of Gulazyk, as well as the quantity and date of receipt and whether caretakers permanently refuse future Gulazyk disbursements.

Using data from the Green Journal, the MOH has conducted an ongoing internal monitoring system. Clinic personnel abstract information from each journal regarding the percentage of children who have ever received Gulazyk and the percentage of children whose caretaker permanently refused to give Gulazyk. This information is collected at the health clinic level and sent to the oblast and national level where it is compiled on a quarterly basis.

In addition to internal monitoring, external monitoring has been conducted through three national surveys in 2011, 2012 and 2013. The 2011 and 2102 surveys were based on Lot Quality Assurance Sampling (LQAS) methods and the 2013 survey was based on population proportional sampling (PPS) methods (see Chapter 2).

This Chapter is divided into two parts: Gulazyk use in 2013 and Trends in Gulazyk use 2011, 2012 and 2013.

## GULAZYK USE IN 2013

### Methods

Data were collected by abstracting information from the Green Journal at the FAP/FGP and by interviewing caretakers. Teams reviewed the Green Journal at each FAP/FGP and recorded if the child had ever been given Gulazyk, the last date that it had been received, and whether the child had received Gulazyk at any point in the previous 3 months (See 2013 Questionnaire, Appendix I). Teams also interviewed caretakers about receipt of Gulazyk, current use, any side effects, where the Gulazyk was obtained, perceived changes in color/taste of food after adding Gulazyk, and plans for future use. Analyses for the 2013 survey are restricted to children 6 to 23 months of age, the age at which children are eligible for the Gulazyk program.

### Caregiver Report of Receipt and Use of Gulazyk

Table 4-1 presents Gulazyk program monitoring indicators related to receipt and use of Gulazyk according to the caregiver report.. Nearly all (95.0%) of the caretakers interviewed had received at least one package of Gulazyk. When asked how they obtain Gulazyk, 87.9% of caretakers indicated that they get the product from the local health clinic, 6.5% said a health care worker brings Gulazyk to their home, and 5.4% said they obtain Gulazyk through both clinic visits and home visits.

Among those who had ever received Gulazyk, the majority of caretakers (73.5%) reported that their child is “currently” taking Gulazyk<sup>3</sup>. It should be noted that the recall period for currently consuming Gulazyk was not defined in the question. Because Gulazyk was given on a flexible schedule it is possible that some children who were adherent (i.e. consumed Gulazyk every day in the previous month, but not currently consuming) may have been reported as not “currently” consuming. When asked about the schedule they use to give Gulazyk to their children, most (45.3%) caretakers indicated that they give Gulazyk every other day. Approximately 37% of the caretakers said they give Gulazyk every day for one month. Very few caretakers reported using another schedule, such as one sachet per day for fifteen days and then taking a break for fifteen days. On average, caretakers reported that they gave 28.2 sachets of Gulazyk to their children in the last two months. The majority of caretakers (80.1%) indicated that they will continue to give Gulazyk when they use up their current package. Among those whose child stopped consuming Gulazyk, caretakers were asked the age when the child stopped consuming Gulazyk; 49.6%, 31.6% and 18.8% reported stopping at 6-11, 12-17 and 18-23 months of age, respectively.

**Table 4-1: Receipt and use of Gulazyk among children 6.0–23.9 months according to the caretaker report— Kyrgyzstan, 2013**

Item	N	%	mean	(95% CI)
Ever received a package of Gulazyk, %	1628			
Yes		95.0		(92.9 - 96.5)
No		4.9		(3.4 - 6.9)
Among those who have ever received a package of Gulazyk:	1537			
How Gulazyk is obtained, %				
From the health clinic		87.9		(83.9 - 91.1)
Medical worker brings it to home		6.5		(4.2 - 9.9)
Both		5.4		(3.5 - 8.1)
Child is currently consuming Gulazyka ,%	1537			
Yes		73.5		(66.8 - 79.3)

<sup>3</sup> The question asked was: “Is this child currently taking Gulazyk?” The time frame was not defined.

No		26.3	(20.6 - 32.9)
<b>Among children not currently consuming Gulazyk:</b>			
<b>Age when child stopped consuming Gulazyk, %</b>	<b>290</b>		
6.0-11.9 months		49.6	(40.3 - 59.0)
12.0-17.9 months		31.6	(23.7 - 40.7)
18.0-23.9 months		18.8	(12.7 - 27)
<b>Frequency of Gulazyk consumption, % (based on the time when child was taking Gulazyk)</b>			
	<b>1537</b>		
Every day for one month		37.2	(31.3 - 43.4)
Every other day		45.3	(37.3 - 53.6)
1 sachet/day for 15 days, break for 15 days		11.9	(6.8 - 20)
Other		3.5	(2.1 - 5.8)
<b>Number of sachets consumed during the last two months, mean</b>	<b>1507</b>	<b>28.2</b>	<b>(25.3 - 31.1)</b>
<b>Will continue giving Gulazyk when current supply runs out, %<sup>a</sup></b>			
	<b>1537</b>		
Yes		80.1	(74.2 - 85.0)
No		12.6	(8.5 - 18.3)
Don't know		6.2	(4.1 - 9.3)

NOTE: CI=confidence interval; 95% confidence intervals adjusted for cluster survey design.

<sup>a</sup>The question asked was "When you run out of Gulazyk, do you plan to continue?"

Table 4-2 shows the use of Gulazyk in the last two months by age, place of residence, and method by which Gulazyk was obtained. We excluded from our calculations children 6-7.9 months of age as they would not have had an opportunity to consume Gulazyk over the previous 2 months. Among all children who received Gulazyk, 56.0% of children consumed at least 30 sachets of Gulazyk (one package) during the last two months. Among children 8.0-11.9 months of age, 60.2% consumed 30 or more sachets (a full package) in the last two months, 28.2% consumed less than 30 sachets, and 11.7% consumed zero sachets. Among children 12.0-17.9 months of age, 59.1% consumed 30 or more sachets in the last two months, 24.8% consumed less than 30 sachets, and 16.2% consumed zero sachets. Among children 18.0-23.9 months of age, 50.1% consumed 30 or more sachets in the last two months, 26.2% consumed less than 30 sachets, and 23.7% consumed zero sachets.

Among urban children, 56.3% consumed  $\geq$  30 sachets compared to 55.3% of rural children. When consumption was stratified by how Gulazyk was obtained, the percentage consuming  $\geq$  30 sachets was 55.2% for those who received Gulazyk at the health clinic, 56% for those who received Gulazyk through a medical worker home visit and 68.8% for those who received from both sources.

**Table 4-2: Among those who had ever received Gulazyk, amount of Gulazyk consumed within a two month period by various characteristics — Kyrgyzstan, 2013**

Characteristic	0 Sachets			< 30 Sachets			≥30 Sachets		
	N	%	(95% CI)	N	%	(95% CI)	N	%	(95% CI)
Age of children <sup>a</sup>	206			403			761		
8.0-11.9 months	28	11.7	(7.2 - 18.4)	123	28.2	(21.4 - 36.1)	187	60.2	(50.5 - 69.1)
12.0-17.9 months	79	16.2	(10.6 - 23.9)	142	24.8	(18.7 - 32.0)	316	59.1	(50.7 - 66.9)
18.0-23.9 months	99	23.7	(16.2 - 33.4)	138	26.2	(20.0 - 33.5)	258	50.1	(42.2 - 57.9)
Place of Residence <sup>a</sup>	206			403			761		
Urban	71	19.9	(12.8 - 29.6)	84	23.8	(17.5 - 31.5)	199	56.3	(47.0 - 65.2)
Rural	135	13.1	(10.1 - 16.8)	319	31.5	(25.8 - 37.9)	562	55.3	(48.6 - 61.9)
How Gulazyk is obtained <sup>a</sup>	206			403			761		
From the health clinic	186	19.0	(13.4 - 26.2)	316	25.8	(20.4 - 32.1)	614	55.2	(48.0 - 62.2)
Medical worker brings to home	14	13.5	(6.8 - 25.1)	65	30.5	(18.2 - 46.4)	80	56.0	(38.4 - 72.3)
Both	6	5.7	(1.7 - 17.8)	22	25.4	(14.5 - 40.6)	67	68.8	(51.0 - 82.4)
Total	206	17.9	(12.6 - 24.6)	403	26.1	(21.2 - 31.7)	761	56.0	(49.2 - 62.6)

<sup>a</sup> CI=confidence interval; 95% confidence intervals adjusted for cluster survey design.

<sup>a</sup> Frequencies add to 100% across rows (horizontally)

<sup>b</sup> Restricted to children 8.0-11.9 months because children < 8.0 months would not have had the opportunity to consume all 30 sachets within the previous two months.

Table 4-3 shows the reasons why caretakers choose not to give Gulazyk to children and the reasons why caretakers will not continue to give Gulazyk when their current package of 30 sachets runs out. Among those not currently consuming Gulazyk, the most commonly cited reasons for not consuming were: the child does not like the food when Gulazyk is added (49.9%), the caretaker perceived that the child experienced side effects/diarrhea (13.7%), the caretaker ran out of Gulazyk (13.7%), Gulazyk makes the food taste bad (8.3%), and the mother/caretaker doesn't like Gulazyk (7.5%). Some caretakers reported they would not give Gulazyk in the future once their current supply runs out, the most commonly cited reasons caretakers gave for not continuing to give Gulazyk were: child does not like the food when Gulazyk is added (72.8%), child experienced side effects (28.4%), children should receive natural vitamins from food (9.5%), and family member doesn't want to use (7.5%).

**Table 4-3: Reasons why caretakers choose not to give Gulazyk to children and why caretakers will not continue to give Gulazyk — Kyrgyzstan, 2013**

Item	%	(95% CI)
<b>Reasons why children are not currently consuming Gulazyk (of those who are not currently consuming Gulazyk) (N=346)<sup>a</sup></b>		
Too difficult to remember to give Gulazyk	3.6	(1.6 - 7.6)
Child experienced side effects/diarrhea	13.7	(9.2 - 19.8)
Child experienced side effects/skin/allergy	3.7	(1.4 - 9.5)
Child experienced side effects/other	0.4	(0.1 - 1.1)
Child does not like food when Gulazyk is added	49.9	(41.7 - 58.2)
Gulazyk makes the food taste bad	8.3	(5.2 - 12.9)
Gulazyk changes the color of the food	2.4	(1.0 - 5.7)
Mother/caretaker doesn't like Gulazyk	7.5	(3.2 - 16.3)
Ran out of Gulazyk	13.7	(8.5 - 21.4)
Children should receive natural vitamins from food	3.8	(1.6 - 8.7)
We have been away from home	5.0	(2.1 - 11.3)
Child has been sick	4.6	(2.0 - 10.5)
Belief that Gulazyk is harmful to the child	0.9	(0.2 - 4.6)
Heard on the TV or radio that Gulazyk is harmful	0.1	(0.0 - 0.7)
Don't know	2.9	(1.5 - 5.6)
<b>Reasons why caretakers will not continue to give Gulazyk when their current package runs out (of caretakers who indicated they will not continue to give Gulazyk when their current package is used up) (N=166)<sup>a</sup></b>		
Too difficult to remember to give Gulazyk	0.6	(0.2 - 2)
Child experienced side effects	28.4	(17.2 - 43.1)
Child does not like food when Gulazyk is added	72.8	(64.4 - 79.8)
Family member doesn't want to use	7.5	(3.6 - 15.2)
Children should receive natural vitamins from food	9.5	(4.4 - 19.5)
Belief that Gulazyk is harmful to the children	0.5	(0.1 - 1.9)
Heard on the TV or radio that Gulazyk is harmful	1.9	(0.4 - 9.6)
When I run out of Gulazyk my child will be >24 months	2.9	(1.0 - 8.2)
Don't know	3.6	(1.3 - 9.7)

NOTE: CI=confidence interval; 95% confidence intervals adjusted for cluster survey design.

<sup>a</sup> Does not add up to 100% because respondents were allowed to provide more than one answer to the question; predefined response options were not read aloud, but rather respondents' individual answers were recorded and categorized.

## Experiences Giving Gulazyk

Table 4-4 reports caretakers' experiences with giving Gulazyk. The majority of caretakers (56.4%) indicated that they notice changes in the color of food to which Gulazyk is added. Among those who noticed a change in color, 67.0% responded that the change in color was not a concern for them or their child. Additionally, 57.7% of caretakers reported that they notice changes in the taste of food when Gulazyk is added. Among those who noticed a change in taste, 60.1% responded that the change in taste was not a concern for them or their child. When asked about the portion of food into which Gulazyk is mixed that the child consumes, 66.4% said their child consumed the entire portion and 33.6% said their child consumed less than a full portion.

**Table 4-4: Experiences consuming Gulazyk among those currently consuming Gulazyk— Kyrgyzstan, 2013**

Item	%	(95% CI)
Have you noticed any changes in the color of the food to which Gulazyk is added? (N=1190)		
Yes	56.4	(48.4 - 64.1)
No	43.6	(35.9 - 51.6)
<i>Among those who noticed a change in color:</i>		
Is this change in color a concern for you or your child?(N=652)		
Yes	33.0	(28.5 - 37.8)
No	67.0	(62.2 - 71.5)
Have you noticed any changes in the taste of the food to which Gulazyk is added? (N=1190)		
Yes	57.7	(50.8 - 64.3)
No	42.3	(35.7 - 49.2)
<i>Among those who noticed a change in taste:</i>		
Is this change in taste a concern for you or your child?(N=662)		
Yes	39.9	(33.7 - 46.4)
No	60.1	(53.6 - 66.3)
Does the child usually consume the entire portion of food into which Gulazyk is mixed, or does he/she consume less than the entire portion? (N=1190)		
Consumes entire portion	66.4	(61.5 - 70.9)
Consumers less than full portion	33.6	(29.0 - 38.4)
Don't know	0.1	(0.0 - 0.4)

NOTE: CI=confidence interval; 95% confidence intervals adjusted for cluster survey design.

## Caretaker's Perceived Effects of Gulazyk

Table 4-5 presents caretakers' reports of positive changes that were experienced by the child after he/she started taking Gulazyk. When caretakers were asked if they noticed any positive changes in their children after they started taking Gulazyk, 42.7% said they had noticed positive changes. Among those who had noticed positive changes, 42.6% indicated increased appetite, 33.6% reported that their child overall seems better/healthier, 29.9% of caretakers indicated that the energy level increased, 16.1% indicated more curiosity/intelligence, 10.4% reported that their child gets sick less often, and 9.0% indicated better growth.<sup>4</sup>

<sup>4</sup> When interpreting results about reported positive changes in characteristics of the child after he/she started taking Gulazyk, it should be noted that, anecdotally, some respondents observed that they had noticed differences, but they reported that these differences might also be attributed to the older age of the child (rather than the Gulazyk).

**Table 4-5: Effects of Gulazyk, as reported by respondents who have ever given Gulazyk to their child of 6.0–23.9 months (based on the time when child was taking Gulazyk) —Kyrgyzstan, 2013**

Item	%	(95% CI)
Have you noticed any positive changes in your child since he/she started taking Gulazyk that you believe are due to Gulazyk? (N=1537)		
Yes	42.7	(38.4 - 47.1)
No	57.3	(52.9 - 61.6)
<i>Among those who noticed positive changes:</i>		
What are the positive changes that you noticed in your child? (N=702) <sup>a</sup>		
More energy	29.9	(23.4 - 37.4)
Better growth	9.0	(5.8 - 13.6)
More curiosity/intelligence	16.1	(11.6 - 21.9)
Improved eyesight	0.8	(0.4 - 1.6)
Gets sick less often	10.4	(6.9 - 15.6)
Increased appetite	42.6	(34.8 - 50.9)
Overall seems better/healthier	33.6	(25.8 - 42.4)
Other	5.1	(3.2 - 7.9)

NOTE: CI=confidence interval; 95% confidence intervals adjusted for cluster survey design.

<sup>a</sup> Does not add up to 100% because respondents were allowed to provide more than one answer to the question; predefined response options were not read aloud, but rather respondents' individual answers were recorded and categorized.

### Coverage of Gulazyk Communications Materials and Activities

Caretakers were asked about their receipt of the Gulazyk communications materials (Table 4-6). Sixty percent of all caretakers reported receiving the Gulazyk brochure. Of those who received the brochure, 58.1% read it. Only 43.0% of respondents had a radio at home. Of those who had a radio, 45.6% indicated that they have heard about Gulazyk on the radio. Nearly all caretakers (99.2%) reported having a television. Among those with a television, 68.8% of respondents indicated that they have heard about Gulazyk on television.

**Table 4-6: Communications coverage as reported by caretakers of children 6.0–23.9 months, — Kyrgyzstan, 2013**

Item	N	%	(95% CI)
Received the Gulazyk brochure	1537		
Yes		60.4	(53.7 - 66.7)
No		38.4	(32.1 - 45.0)
Read the brochure	1537		
Yes		58.1	(51.3 - 64.6)
No		40.6	(33.9 - 47.7)
Has a radio at home	1628		
Yes		43.0	(37.7 - 48.4)
No		57.0	(51.6 - 62.3)
Heard about Gulazyk on the radio (among those who have a radio)	682		
Yes		45.6	(39.7 - 51.7)



---

No		54.3	(48.2 - 60.3)
Has a television	1628		
Yes		99.2	(98.4 - 99.6)
No		0.8	(0.4 - 1.6)
Heard about Gulazyk on television (among those who have a television)	1611		
Yes		68.8	(62.7 - 74.3)
No		30.8	(25.4 - 36.9)

---

*NOTE: CI=confidence interval; 95% confidence intervals adjusted for cluster survey design.*

---

## TRENDS IN GULAZYK USE FROM 2011–2013

### *Background*

This section compares results of the 2011–2012 Lot Quality Assurance Sampling (LQAS) surveys and the 2013 National survey in regards to Gulazyk receipt and use in the population of children aged 6.0–23.9 months of age living in the Kyrgyz Republic (with the exception of the city of Bishkek).

### *Methods*

In 2011 and 2012, surveys which used Lot Quality Assurance Sampling (LQAS) were conducted to assess the implementation and acceptance of the nationwide program. In 2013, the survey used population proportional sampling (PPS) (see Chapter 2). For all three surveys, the metropolitan area of Bishkek was excluded from sampling and analysis because the Gulazyk program was not implemented in Bishkek. LQAS is a sampling methodology that uses small samples from each locality to determine if key indicators of public health programs have been met. While originally designed for quality control of industrially produced goods, this method has been successfully used for monitoring and evaluation of a wide variety of public health programs around the world (Robertson 2006; Valdez 2003; Valdez 1991). At the locality level, LQAS uses cumulative binomial probabilities to determine whether a locality has or has not met a predetermined goal. However, when results are compiled on a national level and weighted for the population of each locality, a numerical national estimate of coverage (defined as the caretaker report of consumption of MNP by the child at the time of the survey) can also be calculated.

In the 2011 and 2012 surveys Lot Quality Assurance Sampling (LQAS) was used. Both surveys were stratified at the district (rayon) level. With the exception of the city of Bishkek, the survey sample was stratified by all districts, the urban territories within each province, and the City of Osh for a total of 48 strata. In each stratum, 24 children aged 6–23 months were selected using systematic sampling from clinic registries. Caretakers of selected children were interviewed in their homes. If the caretakers were not in their homes when the interviewers arrived and a contact phone number was available, the interviewers attempted to conduct the interview by phone. In the 2011 survey, data were collected during July and August, and in the 2012 survey data were collected during September.

In 2013, a survey was conducted to assess MNP use and the micronutrient status of the population. A two-stage cluster PPS sampling design (with no stratification) was used (see Chapter 2). In the 2013 survey, data were collected July and August. The 2013 survey interviews were conducted at the health clinic.

The 2011, 2012, and 2013 questionnaires contained modules on MNP use as recorded in the health clinic (Green Journal) and MNP use as reported by the caretaker (see 2013 questionnaire in Appendix I and 2011 and 2012 questionnaires in Appendix IV). Teams reviewed the clinic registry at each FAP/FGP and recorded Gulazyk had ever been distributed to the child, the last date that it had been received, and whether the child had received Gulazyk at any point in the previous 3 months. In all three surveys, questionnaires were written in English and then translated into the Kyrgyz and Russian languages. Teams also interviewed caretakers about receipt of Gulazyk, current use, any side effects, where the Gulazyk was obtained, perceived changes in color/taste of food after adding Gulazyk, and plans for future use. During the 2013 interviews were conducted at the FAP/FGP; the 2011 and 2012 LQAS interviews were conducted in the home of the child.

Although the survey sampling methods were different, the three survey questionnaires (2011 LQAS, 2012 LQAS, and 2013 national survey) were identical in regard to the questions asked concerning Gulazyk use with two exceptions. The first exception concerned the Russian and Kyrgyz translations of question GU2, where caregivers were asked if the child was currently taking Gulazyk. For the 2013 survey, the translations were changed from “taking” to “consumed” due to confusion on behalf of the caregivers on the difference between taking, as in receiving, and consuming. The second exception concerned the format of the questions on change in color or taste and, among those who had noticed a change, whether the change was of concern. In 2013, separate questions were asked about

---

change in color and/or taste and concerns about color and/or taste (see Appendix I: GU8, GU8a, GU9 and GU9a) whereas in the 2011 and 2012 surveys, the question on change in color or taste were combined into one questions (see Appendix IV: GU9 and GU9a).

Data analysis was conducted using SPSS (versions 17 and 19). For the 2011 and 2012 surveys only interviews conducted in the home were included in the analysis. Bivariate statistical testing was used to determine the significance of differences between the 2011, 2012 and 2013 surveys. The 2011 and 2012 surveys were weighted to account for stratification (district) and non-response. The 2013 survey was weighted for non-response only (the sample was not stratified). To compare the Kyrgyzstan 2011, 2012, and 2013 assessments, approximate p-values were calculated for an overall chi-square and test-for-trend. A spreadsheet was developed where the prevalence and confidence limits from each of the three assessments were entered. For each assessment, the effective sample size was calculated based on the point estimate and confidence limits (Kish, 1965). Based on the effective sample size, an overall chi-square and test-for-trend p-values were calculated. (Rosner, 2010). P-values for the Pearson chi-square test are presented to indicate whether there are statistically significant differences in these indicators among the 2011, 2012 and 2013 surveys. The Pearson chi-square tests were run using methods that account for the design of the surveys (standard errors were adjusted for the cluster survey design). The overall chi-square can be interpreted as to whether or not there was a statistically significant association in the estimates across the three years. The test-for-trend can be interpreted as whether or not there was a statistically significant linear increase or decrease in estimates across the three years.

### **Trends in Gulazyk use**

Table 4-7 compares trends in national key indicator results among children 6.0–23.9 months of age from 2011–2013. Between 2011 and 2013, Green Journal records indicated that the percentage of children who had ever received Gulazyk was slightly over 90% in all survey years and remained stable ( $p=0.252$ ). The percentage children who had received Gulazyk within 3 months of the interview declined by 8.8% ( $p < 0.001$ ). According to caretaker report, the percentage of children who had ever received Gulazyk was about 95% during all three survey years. Among children who had ever received Gulazyk, the percentage children who were currently receiving Gulazyk decreased 5.9% (79.4% versus 73.5%,  $p < 0.001$ ). Caretakers reporting a perceived change in the color or taste of food increased 20.2% (46.4% versus 66.6%,  $p < 0.001$ )<sup>5</sup>. Among those who had noticed a change in the taste or color, there was little change in the percentage who were concerned (31.6% versus 31.9%,  $p=0.421$ ) about the change in color or taste. Among children who had ever received Gulazyk and who were not going to age out of the program prior to their next scheduled receipt of the product, care takers reported an 8.1% decrease (89.1% versus 81.0%,  $p=0.003$ ) in their plans to continue use of Gulazyk in the future.

---

<sup>5</sup> It should be noted that questions on change in color and/or taste were not asked in the same way across all survey years. The changes in questionnaire format may account for some of the differences in response.

**Table 4-7: National Key Indicator Results— among children 6.0–23.9 months of age, Kyrgyzstan, 2011–2013**

	2011 <sup>a</sup>			2012 <sup>a</sup>			2013 <sup>b</sup>			Overall p-value <sup>c</sup>	Test-for-trend p-value
	N	%	95% CI	N	%	95% CI	N	%	95% CI		
<b>Green Journal Indicators</b>											
Ever received Gulazyk	1023	93.2	(90.9-94.9)	995	90.8	(88.4-92.6)	1628	92.1	(88.7 - 95.5)	0.252	0.304
Received Gulazyk within 3 months of interview	1023	84.9	(81.9-87.6)	995	72.8	(69.1-76.3)	1628	76.1	(68.9 - 82.1)	0.000	0.000
<b>Questionnaire Indicators (per caretaker report)</b>											
Ever received Gulazyk	1023	95.3	(93.3-96.7)	995	95.9	(94.3-97.1)	1628	95.0	(92.9 - 96.5)	0.707	0.818
Currently consuming Gulazyk <sup>d</sup>	974	79.4	(76.1-82.4)	949	69.7	(65.9-73.2)	1537	73.5	(66.8 - 79.3)	0.000	0.004
Noticed a change in taste or color of food <sup>d,e</sup>	972	46.4	(42.5-50.3)	936	57.9	(54.3-61.5)	1535	66.6	(60.3 - 72.3)	0.000	0.000
Concerned about change in taste or color <sup>d,e</sup>	515	31.6	(26.5-37.2)	471	27.8	(23.3-32.8)	1535	31.9	(27.6 - 36.7)	0.421	0.834
Plan to continue to give Gulazyk <sup>f</sup>	974	89.1	(86.6-91.2)	949	84.6	(81.6-87.3)	1270	80.1	(74.2 - 85.0)	0.003	0.001

NOTE: CI=confidence interval

<sup>a</sup> Results aggregated across all localities, using population to weight results from each locality within national total (excluding Bishkek city).

<sup>b</sup> 95% confidence intervals adjusted for cluster survey design

<sup>c</sup> Bivariate tests of statistical significance were conducted using the Pearson chi-square test, accounting for the cluster survey design.

<sup>d</sup> Among children who ever received Gulazyk.

<sup>e</sup> Separate questions on color and taste were asked in 2013 whereas they were combined into one question in 2011 and 2012.

<sup>f</sup> Out of those children currently consuming Gulazyk and who were not aging out of the program prior to running out of sachets

**Table 4-8: Other Indicators Related to Gulazyk Receipt and Use — among children 6.0–23.9 months of age who had ever received Gulazyk, Kyrgyzstan, 2011–2013**

	2011 <sup>a,b</sup>			2012			2013			Overall p-value	Test-for-trend p-value
	N	%	Mean 95% CI	N	%	Mean 95% CI	N	%	Mean 95% CI		
<b>Where Gulazyk obtained by family, %</b>											
Health clinic	974	53.9	(49.9-57.8)	949	43.8	(39.9-47.7)	1537	87.9	(84.4 – 90.8)	0.000	0.000
Home health worker		37.4	(33.6-41.3)		34.7	(30.9-38.6)		6.5	(4.2 – 10.0)	0.000	0.000
Both of above		8.5	(6.8-10.6)		20.8	(17.8-24.1)		5.4	(3.6 – 7.9)	0.000	0.918
Other		0.2	(0.0-1.2)		0.8	(0.2-2.3)		0.2	(0.0 – 1.1)	0.107	0.996
<b>Schedule of consumption, %</b>											
Every day for one month	974	27.2	(23.6-31.0)	949	24.7	(21.4-28.2)	1537	37.2	(31.3 - 43.4)	0.001	0.029
Every day for 15 days, Alternating		14.5	(12.0-17.5)		6.6	(4.8-9.1)		11.9	(6.8 – 20.0)	0.000	0.003
Every other day		49.9	(45.9-53.9)		55.8	(51.8-59.8)		45.3	(37.3 - 53.6)	0.031	0.766
Other schedule		7.7	(5.7-10.4)		10.1	(8.1-12.5)		3.5	(2.1 - 5.8)	0.000	0.035
Never consumed		0.1	(0.0-0.3)		1.0	(0.5-2.0)		1.6	(0.9 - 2.7)	0.000	0.000
Don't know		0.6	(0.2-1.7)		1.8	(0.9-3.5)		0.7	(0.2 - 2.1)	0.063	0.717
<b>Number of sachets of Gulazyk consumed by child in last 2 months (mean #)</b>											
	972	23.9	(22.3-25.5)	936	33.7	(31.4-36.1)	1507	28.2	(25.3 - 31.1)	0.000	0.000

NOTE: CI=confidence interval

<sup>a</sup>Results aggregated across all localities, using population to weight results from each locality within national total (excluding Bishkek city).

<sup>b</sup>95% confidence intervals adjusted for cluster survey design

<sup>c</sup>Bivariate tests of statistical significance were conducted using the Pearson chi-square test, accounting for the cluster survey design.

**Table 4-9: Reasons given for not using Gulazyk, among children 6.0–23.9 months of age not currently consuming Gulazyk, Kyrgyzstan, 2011–2013**

	2011 <sup>a</sup>			2012 <sup>a</sup>			2013 <sup>b</sup>			Overall p-value <sup>d</sup>	Test-for-trend p-value
	N	%	95% CI	N	%	95% CI	N	%	95% CI		
Too difficult to remember to give Gulazyk	203	1.0	(0.4-2.8)	302	3.5	(1.6-7.6)	346	3.6	(1.7 - 7.1)	0.018	0.008
Child experienced side effects (total)		20.6	(15.5-26.9)		29.3	(23.3-36.2)		17.6	(11.3 - 26.4)	0.036	0.931
Diarrhea		20.6	(15.5-26.9)		26.0	(20.2-32.8)		13.7	(9.3 - 19.7)	0.013	0.110
Skin/allergy		0.0	N/A		1.1	(0.4-3.5)		3.7	(1.4 - 9.5)		
Other side effect(s)		0.0	N/A		4.0	(2.2-7.2)		0.4	(0.1 - 1.2)		
Perceived change in food due to Gulazyk (total)		10.4	(6.4-16.4)		17.4	(12.4-23.8)		52.9	(45.0 - 60.7)	0.000	0.000
Child does not like food when Gulazyk added		9.7	(5.8-15.6)		14.1	(9.6-20.2)		49.9	(41.8 - 58.0)	0.000	0.000
Gulazyk makes food taste bad		0.3	(0.0-2.0)		3.0	(1.3-7.0)		8.3	(5.2 - 12.9)	0.000	0.000
Gulazyk changes the color of the food		1.3	(0.4-4.3)		2.9	(1.2-6.9)		2.4	(1.0 - 5.7)	0.397	0.323
Mother/caretaker doesn't like Gulazyk		3.2	(1.5-6.8)		7.2	(4.3-11.9)		7.5	(3.3 - 16.0)	0.105	0.054
Health care worker told family to stop using Gulazyk		0.1	(0.0-1.0)		0.2	(0.0-1.4)		0.0	N/A		
Ran out of Gulazyk		45.9	(38.0-54.0)		32.8	(26.5-39.9)		13.7	(8.6 - 21.2)	0.000	0.000
Believe children should get natural vitamins from food		2.2	(0.8-5.6)		0.7	(0.2-2.4)		3.8	(1.6 - 8.7)	0.012	0.357
Family has been away from home		2.0	(0.9-4.6)		1.0	(0.3-2.8)		5.0	(2.1 - 11.2)	0.014	0.185
Child has been sick		11.5	(7.3-17.5)		4.4	(2.4-8.0)		4.6	(2.0 - 10.5)	0.008	0.011
Don't know		1.5	(0.4-4.9)		1.9	(0.7-5.2)		2.9	(1.5 - 5.4)	0.463	0.227
Other		0.0	N/A		25.3	(20.3-31.0)		10.3	(6.0 - 17.0)		

NOTE: CI=confidence interval

<sup>a</sup> Results aggregated across all localities, using population to weight results from each locality within national total (excluding Bishkek city).

<sup>b</sup> Percentages do not add to 100% because respondent could choose more than one option.

<sup>c</sup> 95% confidence intervals adjusted for cluster survey design

<sup>d</sup> Bivariate tests of statistical significance were conducted using the Pearson chi-square test, accounting for the cluster survey design.

---

Table 4-8 compares trends in other indicators related to Gulazyk receipt and use results among children 6.0–23.9 months of age from 2011–2013. Between 2011 and 2013, the majority of respondents obtained Gulazyk from a health clinic, with a 34.0% increase (53.9 versus 87.9%,  $p<0.001$ ) in respondents obtaining Gulazyk from a health clinic. Across the three years, a schedule of every other day was the most frequently followed regimen, followed by every day for one month and every day for 15 days, alternating. Between 2011 and 2013, the percentage of respondents reporting a schedule of every other day consumption decreased 4.6% (49.9% versus 45.3%,  $p=0.031$ ) while the percentage giving Gulazyk every day for one month increased 10.0% (27.2% versus 37.2%,  $p=0.001$ ). Finally, the mean number of sachets consumed in the previous 2 months increased from 23.9 in 2011 to 28.2 in 2013 ( $p<0.001$ ).

Table 4-9 presents trends in reasons for not using Gulazyk, among children not currently consuming Gulazyk according to caretakers from 2011–2013. In 2011, the most frequently cited reason for not currently using Gulazyk was “Ran out of Gulazyk” (45.9%) whereas in 2013 the most frequently cited reason (49.9%) was “child does not like food when Gulazyk added”. There was a 3% decrease (20.6% versus 17.6%,  $p=0.036$ ) from 2011 to 2013 in caretakers citing that the child experiencing side effects as the reason for not using Gulazyk.

---

## DISCUSSION

Overall, program monitoring indicators collected during all three surveys showed positive results. Almost all caretakers had heard of Gulazyk and had received at least one package of 30 sachets of Gulazyk. These results show success in the initial outreach efforts within communities to raise awareness of the availability of MNP and draw caretakers to local health clinics where they can obtain Gulazyk. Additionally, a majority of caretakers reported that their children were currently taking Gulazyk at the 2011, 2012 and 2013. However, the prevalence of reported current use decreased by 5.9% from 2011 to 2013. This slight decrease in current use could reflect the fact that as programs move into the maintenance phase, it can be more difficult to sustain long-term interest and motivation among caretakers.

In the 2013 survey two thirds of caretakers reported a change in the taste or color of food compared with about half of caretakers in 2011. Among caretakers who discontinued use of Gulazyk, the percentage reporting that their child did not like the taste of food when Gulazyk was added increased from 10% in 2011 to 50% in 2013. The reasons for these increases are unclear. A possible reason for the change in taste or color is inappropriate preparation and serving time; however, we did not collect information on preparation and could not address whether preparation methods changed over the time period. There was no change to the formulation (ferrous fumarate was coated during all years), manufacturing technology or packaging over the survey time periods (personal communication, Pritesh Shetty, Piramal) Furthermore, Piramal, the distributor, tested samples from the Gulazyk batches supplied to UNICEF in 2010 and 2012<sup>6</sup> and neither showed any abnormal enhanced metallic taste (personal communication, Pritesh Shetty [Piramal], 2014). Persons from the MOH noticed that some children refused to consume Gulazyk during the training; they also noticed an increase in complaints about taste of Gulazyk in 2013, but attributed this to negative press publicity resulting from parliamentary debate concerning future funding of the Gulazyk program. It should be noted that in 2013, all caretakers of children who were currently consuming Gulazyk were asked separate questions about change in color and change in taste when Gulazk was added to the food while in 2011 and 2012, caretakers were asked one combined question about change in color and/or taste (see Appendix I and Appendix IV). Change in question format may have led to an apparent increase in the percentage of all caretakers who noted a change in color or taste (Serdula 1992, Serdula, 1995). However, change in format would not account for the increase seen in the percentage of caretakers who said they stopped Gulazyk because the “child does not like food when Gulazyk added” because identical questions were asked about reasons for discontinuation in all survey years.

A limitation of these surveys is that some information on Gulazyk use were self-reported by caretakers; however, for several questions, information on Gulazyk use was also abstracted from the clinic registry (green journal). The figures for ever receiving Gulazyk and current use obtained through clinic registry were consistently similar to those found through the interviews with caretakers, which can be interpreted as a positive sign that self-reporting was accurate.

During all survey years, caretaker satisfaction and acceptance of Gulazyk was high. However, there was a slight decline in the proportion of caretakers who said they would continue to give Gulazyk from 9 in 10 caretakers in 2011 to 8 in 10 in 2013. Outreach through three complementary communications channels – health care professionals, VHC volunteers, and mass media – may have been an important factor in ensuring satisfactory knowledge, attitudes and practices among caretakers. Coverage of communications materials and activities is an area of program performance that may require programmatic adjustments. When caretakers who had ever received MNP were asked whether they received the MNP communications materials in 2013, it was found that only 60% received the instructional brochure. In regard to mass media outreach, slightly over two thirds of the population had heard about Gulazyk through television. Program administrators will need to make programmatic adjustments to improve the coverage of community outreach activities and distribution of MNP communications materials.

---

<sup>6</sup> The 2012 shipment was distributed in both 2012 and 2013.



---

## CHAPTER 5

# INFANT AND YOUNG CHILD NUTRITION (IYCN): KNOWLEDGE, ATTITUDES, AND PRACTICES

### BACKGROUND

WHO recommends exclusive breastfeeding for the first six months and then at six months introduction of solid, semi-solid and soft foods is recommended to supplement the child's diet (WHO, 2001). Appropriate infant and young child feeding (IYCF) practices include altering the frequency, variety, and amount of foods as a child gets older, while continuing breastfeeding until 2 years of age.

### INFANT AND YOUNG CHILD FEEDING PRACTICES

The *WHO Indicators for Assessing Infant and Young Child Feeding Practices* (2008) were used to measure infant and young child feeding practices (Table 5-1). Because children under 6 months of age were not included in the survey, we could not use the standard WHO indicators for exclusive breastfeeding and age-appropriate breastfeeding. Instead we used maternal recall to estimate these indicators (Table 5-2). The algorithm for calculating indicators of feeding practices is found in Appendix V. In the 2009 baseline survey only mothers answered the IYCN indicator questions, whereas in 2013, all caretakers answered these questions. For consistency, the 2013 IYCN analysis of indicators was restricted to only those interviews where the mother was the respondent, with the exception of dietary and breastfeeding advice received which reflects that of all caretakers.

**Table 5-1: Definitions of the WHO indicators for infant and young child feeding practices (WHO, 2008)**

Early initiation of breastfeeding*: <b>Proportion of children born in the last 23.9 months who were put to the breast within one hour of birth</b>  <b>Children born in the last 23.9 months who were put to the breast within one hour of birth</b> <b>Children born in the last 23.9 months</b>  * included only children 6-23.9 months
Ever breastfed*: <b>Proportion of children born in the last 23.9 months who were ever breastfed</b>  <b>Children born in the last 23.9 months who were ever breastfed</b> <b>Children born in the last 23.9 months</b>  * included only children 6-23.9 months
Continued breastfeeding at 1 year: <b>Proportion of children 12-15.9 months of age who are fed breast milk</b>  <b>Children 12-15.9 months of age who received breastmilk during the previous day</b> <b>Children 12-15.9 months of age</b>
Continued breastfeeding at 2 years: <b>Proportion of children 20-23.9 months of age who are fed breast milk</b>  <b>Children 20-23.9 months of age who received breastmilk during the previous day</b> <b>Children 20-23.9 months of age</b>

Введение твердой, полутвердой или мягкой пищи: **Соотношение детей в возрасте 6-8.9 месяцев, получающих твердую, полутвердую или мягкую пищу**

**Дети в возрасте 6-8.9 мес., получивших накануне твердую, полутвердую или мягкую пищу**  
**Дети в возрасте 6-8.9 месяцев**

Introduction of solid, semi-solid or soft foods: **Proportion of infants 6-8.9 months of age who receive solid, semi-solid or soft foods**

**Infants 6-8.9 mo of age who received solid, semi-solid or soft foods during the previous day**  
**Infants 6-8.9 months of age**

Minimum dietary diversity: **Proportion of children 6-23.9 months of age who receive foods from  $\geq 4$  food groups**

**Children 6-23.9 mo of age who received foods from  $\geq 4$  food groups during the previous day**  
**Children 6-23.9 months of age**

Minimum meal frequency: **Proportion of breastfed and non-breastfed children 6-23.9 months of age who receive solid, semi-solid, or soft foods (but also including milk feeds for non-breastfed children) the minimum number of times or more**

**Breastfed children 6-23.9 mo of age who received solid, semi-solid or soft foods the minimum number of times or more during the previous day**  
**Breastfed children 6-23.9 months of age**

and

**Non-breastfed children 6-23.9 mo of age who received solid, semi-solid or soft foods or milk feeds the minimum number of times or more during the previous day**  
**Non-breastfed children 6-23.9 months of age**

Minimum acceptable diet: **Proportion of children 6-23.9 months of age who receive a minimum acceptable diet (apart from breast milk)**

**Breastfed children 6-23.9 months of age who had at least the minimum dietary diversity and the minimum meal frequency during the previous day**  
**Breastfed children 6-23.9 months of age**

and

**Non-breastfed children 6-23.9 months of age who received at least 2 milk feedings and had at least the minimum dietary diversity and the minimum meal frequency during the previous day**  
**Non-breastfed children 6-23.9 months of age**

Milk feeding frequency for non-breastfed children: **Proportion of non-breastfed children 6-23.9 months of age who receive at least 2 milk feedings**

**Non-breastfed children 6-23.9 months of age who received at least 2 milk feedings during the previous day**  
**Non-breastfed children 6-23.9 months of age**

**Table 5-2: Definitions of other infant and young child feeding indicators used**

<p>Exclusive breastfeeding under 6 months: <b>Proportion of children 6-23.9 months of age whose mothers reported that they were exclusively breastfed (no other liquids, milks, or other foods) until 6 months of age</b></p> <p style="text-align: center;"><b>Children who were exclusively breastfed until 6 months of age</b> <b>Children 6-23.9 months of age</b></p>
<p>Age appropriate breastfeeding (children 6-23.9 months): <b>Proportion of children 6-23.9 months of age who are appropriately breastfed</b></p> <p style="text-align: center;"><b>Children 6-23.9 months of age who received breast milk, as well as solid, semi-solid or soft foods, during the previous day</b> <b>Children 6-23.9 months of age</b></p>

The estimates of indicators for infant and young child feeding practices are summarized in Table 5-3. Among children, 85.1% were reported to have been breastfed within the first hour after birth (early initiation of breastfeeding). Among children, 99.5% were reported to have ever been breastfed and 40.2% had been exclusively breastfed during the first 6 months of life. A total of 80.1% of children continued breastfeeding at 1 year and 26.2% continued breastfeeding at 2 years.

Appropriate introduction of solid, semi-solid or soft foods was reported in 90.9% of all children. The criteria for minimum dietary diversity (four or more food groups) was met by 86.8% of children and the criteria for minimum meal frequency (three or more times a day) was met by 74.7% of children. Combining the indicators of minimum dietary diversity and minimum meal frequency resulted in 67.2% of children reaching the criteria for minimum acceptable diet. Age appropriate breastfeeding (proportion of children 6 to 23.9 months of age who were appropriately breastfed) was reported for 62.8% of children. Adequate milk feeding frequency (at least 2 milk feedings a day) was reported for 65.4% of non-breastfed children.

**Table 5 3: Prevalence of infants achieving each Infant Young Child Nutrition (IYCN) indicator for appropriate feeding practices among children aged 6–23.9 months as reported by mothers — Kyrgyzstan, 2013**

IYCN Indicator	N	%	(95% CI)
Early initiation of breastfeeding	1416	85.1	(82.1 - 88.1)
Exclusive breastfeeding under 6 months	1410	40.2	(34.9 – 45.6)
Ever breastfed	1416	99.5	(98.9 - 100.1)
Continued breastfeeding at 1 year	348	80.1	(72.7 - 87.5)
Continued breastfeeding at 2 years	295	26.2	(20.8 - 31.5)
Appropriate introduction of solid, semi-solid or soft foods	212	90.9	(86.2 - 95.7)
Consuming minimum dietary diversity	1416	86.8	(84.1 – 89.6)
Consuming minimum meal frequency	1416	74.7	(70.9 - 78.5)
Consuming minimum acceptable diet	1416	67.2	(63.1 – 71.3)
Age-appropriate breastfeeding	1416	62.8	(59.3 - 66.3)
Adequate milk feeding frequency for non-breastfed children	637	65.4	(56.4 - 74.4)

NOTE: CI=confidence interval; 95% confidence intervals adjusted for cluster survey design.

## KNOWLEDGE AND ATTITUDES REGARDING INFANT AND YOUNG CHILD FEEDING

Table 5-4 presents knowledge and attitudes of mothers regarding breastfeeding and feeding of their babies. Among caretakers, 75.6% considered breastfeeding very important for a baby's health and nutrition. Almost all (99.6%) of the mothers felt a baby should be breastfed and 97.7% felt there were advantages to breastfeeding. When asked how long a baby should be breastfed, on average mothers reported that children ideally should be breastfed for the first 23.9 months. Caretakers reported children should be given other liquids (tea, water, animal milk, etc.) at the age (mean) of 5.2 months and children should start eating other foods at the age (mean) of 7.4 months.

**Table 5 4: Knowledge and attitudes regarding infant and young child feeding as reported by all caretakers — Kyrgyzstan, 2013**

Пункт	N	%, средний	(95% ДИ)
Importance of breastfeeding, %	2163		
Very important		75.6	(68.8 – 81.3)
Important		24.2	(18.6 - 31.0)
Somewhat important		0.1	(0.0 - 0.4)
Not important		0.02	(0.0 - 0.2)
Baby should be breastfed, %	2163		
Yes		99.6	(99.0 – 99.8)
No		0.1	(0.0 - 0.2)
<i>Among those who felt a baby should be breastfed:</i>			
Length of time a baby should be breastfed (in months), mean	2154	23.9	(22.9 – 24.8)
Age at which a baby should start drinking other liquids like tea, water, milk, etc. (in months), mean	2163	5.2	(4.7 - 5.6)
Age at which a baby should start eating foods like porridge, cereal, bulymak, etc. (in months), mean	2163	7.4	(6.9 - 7.8)
Advantages to breastfeeding, %	2163		
Yes		97.7	(96.1 - 98.6)
No		2.3	(1.4 – 3.9)

NOTE: CI=confidence interval; 95% confidence intervals adjusted for cluster survey design.

## DIETARY AND BREASTFEEDING ADVICE

Caretakers were asked about the advice they received from medical professionals regarding breastfeeding and feeding their baby which is summarized in Table 5-5. Overall, 87.8% of caretakers reported receiving advice on breastfeeding from a doctor, nurse, midwife, or feldsher. Among those who had received advice, on average, caretakers reported that medical professionals recommended breastfeeding for 23.3 months and to exclusively breastfeed for approximately 7.2 months.

**Table 5-5: Dietary and breastfeeding advice received by all caretakers — Kyrgyzstan, 2013**

Item	N	%, mean	(95% CI)
Doctor, nurse, midwife, or feldsher gave advice on breastfeeding, %	2163		
Yes		87.8	(84.8 - 90.3)
No		6.5	(4.8 - 8.6)
Don't know		5.7	(4.0-8.0)
<i>Among those who received advice on breastfeeding from a doctor, nurse, midwife, or feldsher:</i>			
Length of time advised to breastfeed without giving other liquids or solids (in months), mean	1698	7.2	(6.7 – 7.7)
Age at which advised to stop breastfeeding (in months), mean	1223	23.3	(22.5 – 24.2)

NOTE: CI=confidence interval; 95% confidence intervals adjusted for cluster survey design.

---

## CHAPTER 6 ANTHROPOMETRY

### BACKGROUND

This chapter documents the anthropometric measurements of children in Kyrgyzstan and compares them to international standards for growth.

### METHODS

Anthropometric indicators of length/height-for-age, weight-for-age, and weight-for-length/ height were determined for all children 6 to 29.9 months (Table 6-1). Shorr boards were used to measure height/length and UNICEF Seca Uniscales were used to measure weight. For children  $\geq 24$  months standing height was measured. For children  $< 24$  months supine length, rather than height, was measured<sup>7</sup>. The age of the child was calculated based on the difference between the child's birth date and the date of the measurement divided by 30.4.

**Table 6-1: Interpretation of anthropometric indicators for children**

Reference: The WHO growth curves for nutritional status will be used to interpret the anthropometric data of the children (WHO, 2006). This system is based on parameters of height for children from six different countries, where children receive proper nutrition and health care in a hygienic environment. The present system is appropriate to use for all population groups. Healthy and well-nourished children from most countries have growth patterns similar to the parameters of this system.

Z-scores: The anthropometric indices used for evaluating the nutritional status of children include height-for-age, weight-for-age, and weight-for-height. These indices are interpreted using classifications based on Z-scores (standard deviation units from the reference median). The WHO recommends that a Z-score cut-off point of  $< -2$  be used to classify low height-for-age (stunting), low weight-for-age (underweight), and low weight-for-height (wasting) for estimating the prevalence of malnutrition (WHO, 1995). A Z-score cut off of  $< -3$  indicates severe wasting, stunting or underweight. The reference Z-score distribution for each index has a mean of 0.0 and a standard deviation of 1.0. A Z-score cut-off of  $+2$  should be used to classify high weight-for-height for estimating the prevalence of overweight (also a form of malnutrition). A Z-score of  $-2$  corresponds to the 2.3rd percentile of the reference distribution, while a Z-score of  $2$  corresponds to the 97.7th percentile on the reference distribution. Thus, with any of the indicators, a prevalence less than or equal to 2.3% is regarded as the surveyed population being free from malnutrition based on that indicator.

Height-for-age: A low height-for-age indicates growth stunting, which reflects a long term deficit of nutritional status and/or a history of illness and disease such as diarrhea and acute respiratory infection. On a population level, a high prevalence of stunting is usually associated with poor socioeconomic conditions and a greater risk for frequent and/or early exposure to adverse environmental conditions such as illness and inadequate nutrition. A decrease in the prevalence of stunting usually parallels improvements in economic conditions (WHO, 1995).

---

<sup>7</sup> For the purpose of this report, the term height will be used to indicate either length or height.

Weight-for-age: This index is a composite of height-for-age and weight-for-height. On a cross-sectional basis, weight-for-age is less useful than height-for-age or weight-for-height in defining nutritional status. In most populations where there are few children with low weight-for-height, the weight-for-age status provides essentially the same information as height-for-age.

Weight-for-height: Low weight-for-height, or wasting, is an indicator of acute under-nutrition and is often the result of severe food shortages and/or prolonged illness.

#### Data Quality:

Data Cleaning: The records with potentially erroneous data were excluded from analysis based on the following standard Z-score cutoffs (WHO, 1995):

Height-for-age, Z-score (HAZ) <-6.0 or >6.0

Weight-for-age, Z-score (WAZ) <-6.0 or >5.0

Weight-for-height, Z-score (WHZ) <-5.0 or >5.0

Data quality: The Standard Deviation (SD) of the Z-score (unweighted) provides information on the spread of the distribution and the quality of the anthropometric measurements done for a survey. In the reference population, the standard deviation (S.D.) of the Z-score distribution for height-for-age and weight-for-height is 1.0. A Z-score S.D. that is lower than 0.9 indicates that the distribution is more homogeneous or one with less variation compared to the reference distribution. A Z-score S.D. >1.0 and <1.2 indicates that the distribution has a wider spread than the reference. A Z-score S.D. <0.80 or >1.3 is suggestive of inaccurate anthropometric measurements and/or inaccurate age information (WHO, 1995).

*NOTE: Supine length, rather than height, is measured for children under 24 months of age.*

The prevalence of anthropometric indicators was interpreted using the WHO classification presented in Table 6-2. Anthropometric results for the WHO global data base on child growth and malnutrition can be found in Appendix VI.

**Table 6 2: WHO classification for low anthropometric values according to public health significance for children < 5 years of age (WHO, 1995)**

Anthropometric Index	Low	Medium	High	Very High
Wasting (WHZ < -2)	<5.0%	5.0-9.9%	10.0-14.9%	≥15.0%
Stunting (HAZ < -2)	<20.0%	20.0-29.9%	30.0-39.9%	≥40.0%
Underweight (WAZ < -2)	<10.0%	10.0-19.9%	20.0-29.9%	≥30.0%

## RESULTS

A total of 2162 children aged 6.0-29.9 months had height/length, weight, and age data to calculate height-for-age (stunting), weight-for-height (wasting), and weight-for-age (underweight) z-scores. Participants were excluded from analysis after applying the standard z-score cutoffs (WHO, 1995): height-for-age (7 children), weight-for-height (2 children), and weight for age (0 children). After excluding the observations with invalid z-scores, the standard deviations for all anthropometry indicators suggest that measurements were accurately taken because they fall within the acceptable range (Mei 2007, WHO 1995; Table 6-3).

**Table 6-3: Distribution of anthropometry measurements among children aged 6.0-29.9 months — Kyrgyzstan, 2013**

Anthropometric Indicator	N	Mean	SD
Height -for-age	2155	-0.59	1.27
Weight-for-height	2160	0.17	1.04
Weight-for-age	2162	-0.17	1.06

Note: SD=standard deviation; SD's are unweighted. Mean estimates are weighted for non-response. Anthropometric values based on WHO growth reference curves (WHO, 2006).

The prevalence of wasting was 2.0% and the prevalence of underweight was 4.8% among children. Overweight children comprised 3.2% of the population (Table 6-4).

**Table 6-4: Prevalence of wasting, underweight, and overweight for children 6.0-29.9 months of age — Kyrgyzstan, 2013**

Anthropometric Indicator	N	%	(95% CI)
Wasting (weight-for-height Z < -2)	2160	2.0	(1.1 - 2.8)
Underweight (weight-for-age Z < -2)	2162	4.8	(3.7 - 5.9)
Overweight (weight-for-height Z > +2)	2160	3.2	(1.8 - 4.5)

NOTE: CI=confidence interval; percent estimates weighted for non-response and 95% CI's adjusted for survey design. Anthropometric values based on WHO growth reference curves (WHO, 2006).

Stunting was identified in 11.7% of the children aged 6-29.9 months (Table 6-5). When stratified by gender, 10.3% of girls were stunted, while 13.1% of boys were stunted. The prevalence of stunting increased across age categories through 23.9 months of age, and was similar among rural children and urban children.

**Table 6-5: Stunting (height -for-age Z<-2) by age, gender, and place of residence for children 6.0-29.9 months of age — Kyrgyzstan, 2013**

Characteristic of Child	N	% Stunted	(95% CI)
<b>Age Group (months)</b>			
6.0-11.9 months	515	5.0	(2.3 - 7.6)
12.0-17.9 months	568	8.6	(5.5 - 11.6)
18.0-23.9 months	539	17.1	(12.5 - 21.7)
24.0-29.9 months	533	15.5	(10.9 - 20.1)
<b>Gender</b>			
Male	1099	13.1	(9.8 - 16.4)
Female	1056	10.3	(7.4 - 13.1)
<b>Place of Residence</b>			
Urban	549	11.3	(8 - 14.6)
Rural	1606	12.6	(10.5 - 14.7)
<b>Total</b>	<b>2155</b>	<b>11.7</b>	<b>(9.3 - 14.1)</b>

NOTE: CI=confidence interval; percent estimates weighted for non-response and 95% CI's adjusted for survey design. Anthropometric values based on WHO growth reference curves (WHO, 2006).



Wasting was identified in 2.0% of the children aged 6-29.9 months (Table 6-6). When stratified by gender, 1.0% of girls were found to have wasting, while 3.0% of boys had wasting. The prevalence of wasting decreased across age categories, and was similar among urban and rural children.

**Table 6-6: Wasting (weight-for-height  $Z < -2$ ) by age, gender, and place of residence for children 6.0-29.9 months of age — Kyrgyzstan, 2013**

Characteristic of Child	N	% with Wasting	(95% CI)
<b>Age Group (months)</b>			
6.0-11.9 months	520	2.7	(0.8 - 4.5)
12.0-17.9 months	568	2.5	(0.8 - 4.2)
18.0-23.9 months	539	2.3	(0.5 - 4.2)
24.0-29.9 months	533	0.5	(0.1 - 0.8)
<b>Gender</b>			
Male	1101	3.0	(1.4 - 4.5)
Female	1059	1.0	(0.3 - 1.7)
<b>Place of Residence</b>			
Urban	552	2.0	(0.8 - 3.2)
Rural	1608	2.0	(1.3 - 2.7)
<b>Total</b>	<b>2160</b>	<b>2.0</b>	<b>(1.1 - 2.8)</b>

NOTE: CI=confidence interval; percent estimates weighted for non-response and 95% CI's adjusted for survey design. Anthropometric values based on WHO growth reference curves (WHO, 2006).

Underweight was identified in 4.8% of the children aged 6-29.9 months (Table 6-7). When stratified by gender, 4.3% of girls were underweight, while 5.3% of boys were underweight. The prevalence of underweight increased slightly across age categories through 23.9 months of age, and was similar among urban and rural children.

**Table 6-7: Underweight (weight-for-age  $Z < -2$ ) by age, gender, and place of residence for children 6.0-29.9 months of age — Kyrgyzstan, 2013**

Characteristic of Child	N	% Underweight	(95% CI)
<b>Age Group (months)</b>			
6.0-11.9 months	520	3.4	(0.9 - 6)
12.0-17.9 months	569	3.7	(1.9 - 5.6)
18.0-23.9 months	539	6.3	(3.4 - 9.3)
24.0-29.9 months	534	5.6	(3.1 - 8.1)
<b>Gender</b>			
Male	1102	5.3	(3.3 - 7.3)
Female	1060	4.3	(2.8 - 5.8)
<b>Place of Residence</b>			
Urban	552	5.0	(3.5 - 6.5)
Rural	1610	4.3	(3.2 - 5.4)
<b>Total</b>	<b>2162</b>	<b>4.8</b>	<b>(3.7 - 5.9)</b>

NOTE: CI=confidence interval; percent estimates weighted for non-response and 95% CI's adjusted for survey design. Anthropometric values based on WHO growth reference curves (WHO, 2006).

---

## DISCUSSION

The prevalence of stunting (11.7%), wasting (2.0%), and underweight (4.8%) among children are all classified as low according to the WHO criteria for public health significance (Table 6-2) (WHO, 1995). Although the WHO criteria apply to population estimates for children less than 59 months, this national nutrition survey included only children 6-29.9 months of age. In general, the prevalence of stunting increases with age, which agrees with other studies (Rah, 2009; UN, 2003]. Because the prevalence of stunting has been shown to be higher in children 24-59 compared to younger children in Kyrgyzstan (CDC/UNICEF/MOH, 2012), it is likely that the prevalence of stunting in all preschool children would be higher than that measured in children 6-29 months (CDC, 2012).

---

## CHAPTER 7

# BIOCHEMICAL INDICATORS FOR MICRONUTRIENT DEFICIENCY AND THE USE OF MICRONUTRIENT SUPPLEMENTS

### BACKGROUND

This chapter highlights 1) the results from biochemical tests to estimate the extent of iron and vitamin A deficiencies and 2) the use of micronutrient supplements.

### METHODS

In order to account for the effect of inflammation on ferritin and RBP levels, the acute phase indicators CRP and AGP were measured for each child. Inflammation was considered present if either indicator was elevated (CRP > 5.0 mg/L or AGP > 1.0g/L). All results are presented for the total population and for the population without inflammation.

### EFFECTS OF INFLAMMATION ON NUTRITIONAL BIOMARKERS

Ferritin and retinol, including its proxy, retinol binding protein (RBP), are acute-phase reactants. Whereas ferritin is elevated during infection/inflammation, retinol and RBP are depressed. Thus, if inflammation is not taken into account, iron deficiency, as measured by serum ferritin, will be underestimated and vitamin A deficiency, as measured by retinol or RBP, will be overestimated. In this survey, both C-reactive protein (CRP) and  $\alpha$ 1-glycoprotein acid (AGP) are used as markers for inflammation. CRP is an acute phase protein that is often used as a marker for acute inflammation, and AGP is used as a marker for chronic inflammation (Thurnham 2003, Thurnham 2010).

### IRON DEFICIENCY, ANEMIA, AND IRON DEFICIENCY ANEMIA

Iron deficiency is the leading cause of anemia, yet not all cases of anemia are caused by iron deficiency and iron deficiency does not necessarily develop into anemia. On the population level a prevalence of iron deficiency is on average 2-5 times higher than the prevalence of iron deficiency anemia (WHO, 2001). The magnitude of iron deficiency can be assessed by several biochemical indicators, including ferritin, soluble transferrin receptor protein (sTfR), and hemoglobin.

In our survey, iron deficiency was defined as (1) decreased ferritin concentration in plasma or (2) increased sTfR levels. In children, ferritin levels in plasma below 12 $\mu$ g/L or sTfR levels greater than 8.3 mg/L indicate iron deficiency (WHO, 2001; Erhardt, 2004). Total iron deficiency was defined as presence of either low ferritin or high sTfR. The prevalence of anemia was determined from hemoglobin levels collected from capillary blood samples using a Hemocue<sup>®</sup> photometer (Hb 301, HemoCue AB, Angelholm, Sweden). Cut-off values for anemia depend on the age and sex of the person and the altitude where the person lives (WHO/UNU/UNICEF, 2001). Anemia in children was defined as hemoglobin < 11.0 g/dL after adjusting for altitude (see details below). Iron deficiency anemia was defined as having both: 1) a low hemoglobin value and 2) low plasma ferritin or high sTfR.

For determining the prevalence of anemia in the population, adjustments for altitude were necessary to account for a reduction in oxygen saturation of blood and a subsequent increase in hemoglobin (Hb) values (WHO/UNU/UNICEF, 2001). The adjustment for altitude was done using the following formula (Sullivan, 2008):

$$\text{Hb adjustment} = -0.032 \times [\text{altitude (m)} \times 0.0032808] + 0.022 \times [(\text{altitude (m)} \times 0.0032808)^2]$$

where the Hb adjustment was the value subtracted from each individual's observed hemoglobin level and then compared to the cut-off values for sea level. If the altitude where the individual lives is

<1000 meters, adjustment of the hemoglobin level is not needed (Sullivan, 2008).

A summary of cut off levels for biochemical indicators used to estimate iron load in the blood are presented in Table 7-1.

**Table 7 1: Biochemical indicators for identification of iron deficiency, anemia, and inflammation**

Indicators	Children	Iron status
Plasma ferritin	<12 µg/L <sup>a</sup>	Iron deficiency (defined by ferritin)
sTfR	>8.3 mg/L <sup>b</sup>	Iron deficiency (defined by sTfR)
Hemoglobin	< 11.0 g/dL <sup>c</sup>	Anemia
CRP	>5 mg/L <sup>d</sup>	Inflammation present
AGP	>1.0 g/L <sup>d</sup>	Inflammation present

Note: sTfR= soluble transferrin receptor protein; CRP=C-reactive protein; AGP=a1-glycoprotein acid.

<sup>a</sup>WHO, 2001

<sup>b</sup>Erhardt, 2004

<sup>c</sup>After adjusting for altitude (WHO, 2001).

<sup>d</sup>Thurnham, 2010

Iron deficiency, anemia, and iron deficiency anemia were calculated for all participants (with and without evidence of inflammation), as well as for those participants without evidence of inflammation (i.e. after exclusion of those with high CRP and/or AGP) (Table 7-2). The prevalence of iron deficiency as measured by ferritin was 34.2% (39.7% among children without inflammation) and as measured by sTfR was 39.3% (38.9% among children without inflammation). The prevalence of total iron deficiency was 48.0% (49.8% among children without inflammation). After adjustment for altitude, the prevalence of anemia was 32.7% (29.5% among children without inflammation). The prevalence of iron deficiency anemia as measured by ferritin was 18.8% (20.4% among children without inflammation) and as measured by sTfR was 21.5% (19.8% among children without inflammation). The prevalence of total iron deficiency anemia was 23.6% (22.4% among children without inflammation).

**Table 7 2: Prevalence of iron deficiency and anemia among children aged 6.0–29.9 months, stratified by presence of inflammationa —Kyrgyzstan, 2013**

Iron Status of Children	All Participants		Participants Without Inflammation <sup>a</sup>	
	n	% (95% CI)	n	% (95% CI)
<b>Iron deficiency</b>				
Low ferritin (<12.0 µg/L)	2156	34.2 (31.3 - 37.2)	1436	39.7(36.4 - 43.1)
High sTfR (>8.3 mg/L)	2156	39.3 (35.6 - 43.1)	1436	38.9 (34.6 - 43.1)
Total iron deficiency <sup>b</sup>	2156	48.0 (44.7 - 51.3)	1436	49.8 (46.2 - 53.4)
<b>Anemia (Hb&lt; 11.0 g/dL)<sup>c</sup></b>				
	2153	32.7 (28.9 - 36.4)	1434	29.5 (25.8 - 33.1)
<b>Iron deficiency anemia</b>				
Low ferritin (<12.0 µg/L)	2153	18.8 (15.6 - 22.0)	1434	20.4 (16.6 - 24.2)
High sTfR (>8.3 mg/L)	2153	21.5 (17.9 - 25.2)	1434	19.8 (15.8 - 23.9)
Total iron deficiency anemia <sup>d</sup>	2153	23.6 (20.1 - 27.1)	1434	22.4 (18.7 - 26.1)

NOTE:CI=confidence interval; 95% CI's adjusted for survey design.

Abbreviations: sTfR= soluble transferrin receptor protein; Hb=hemoglobin.

<sup>a</sup> Inflammation not present (low C-reactive protein [CRP≤5 mg/L] and low a1-glycoprotein acid [AGP≤1.0 g/L]).

<sup>b</sup> Total iron deficiency defined as having either low plasma ferritin (<12.0 µg/L) or high sTfR (>8.3 mg/L).

<sup>c</sup>Adjusted for altitude.

<sup>d</sup>Iron deficiency anemia defined as having a Hb level < 11.0 g/dL and low plasma ferritin (<12 µg/L) or high sTfR (>8.3 mg/L)

The prevalence of anemia among all children and among those without evidence of inflammation is stratified by age, gender, and place of residence (Table 7-3).

The World Health Organization classifies anemia as a problem of public health significance based on prevalence estimates from hemoglobin values (WHO/UNU/UNICEF, 2001). Among all children, 32.7% of all children were anemic and 23.6% had iron deficiency anemia. The prevalence of anemia in this population is considered a moderate public health problem according to the WHO criteria (WHO/UNU/UNICEF, 2001).

**Table 7 3: Prevalence of anemia among children aged 6.0–29.9 months, stratified by age, gender, residence, and presence of inflammation — Kyrgyzstan, 2013**

Characteristics of Child	All Participants		Participants Without Inflammation <sup>b</sup>	
	n	% with Anemia (95% CI)	n	% with Anemia (95% CI)
<b>Age of children</b>				
6.0-11.9 months	519	35.6 (27.8 - 43.4)	354	33.6 (25.4 - 41.7)
12.0-17.9 months	567	35.8 (30.6 - 41.0)	380	33.4 (27.5 - 39.2)
18.0-23.9 months	537	35.1 (30.2 - 40.0)	353	29.2 (23.7 - 34.8)
24.0-29.9 months	530	24.5 (17.7 - 31.3)	347	21.7 (15.6 - 27.8)
<b>Gender of children</b>				
Male	1096	34.5 (29.2 - 39.8)	723	31.5 (26.3 - 36.7)
Female	1057	30.8 (26.3 - 35.2)	711	27.4 (23.0 - 31.8)
<b>Place of Residence</b>				
Urban	551	32.5 (27.3 - 37.7)	361	29.6 (24.5 - 34.7)
Rural	1602	33.0 (29.9 - 36.1)	1073	29.2 (25.7 - 32.6)
<b>Total</b>	<b>2153</b>	<b>32.7 (28.9 - 36.4)</b>	<b>1434</b>	<b>29.5 (25.8 - 33.1)</b>

NOTE:CI=confidence interval; 95% CI's adjusted for survey design.

<sup>a</sup>Adjusted for altitude. Anemia was defined as having a hemoglobin < 11.0 g/dL.

<sup>b</sup>Inflammation not present (low C-reactive protein [CRP≤5 mg/L] and low α1-glucoprotein acid [AGP≤1.0 g/L]).

## VITAMIN A DEFICIENCY

Vitamin A is an essential nutrient required for the immune system, cell function and growth, and epithelial maintenance (WHO, 2009). Kyrgyzstan distributed vitamin A capsules to young children for several years; however the vitamin A capsule distribution program was discontinued based on the low prevalence of deficiency (4.2%) found among children 6-59 months in 2009 (CDC/UNICEF/MOH, 2012).

The most common biochemical indicator used to assess the prevalence of VAD in a population is plasma retinol. A plasma retinol concentration <0.70 µmol/L indicates mild or sub-clinical VAD (WHO, 2009). In the 2013 Kyrgyzstan Survey, retinol binding protein (RBP) was used as a proxy measure of vitamin A status. The CDC nutrition laboratory developed a correlation index comparing plasma retinol to RBP in a sample of children from Kyrgyzstan in order to validate the use of RBP as an indicator of vitamin A status. The RBP cut-off of 0.71 µmol/L provided the best sensitivity and specificity compared with plasma retinol of <0.70 µmol/L (CDC/UNICEF, 2010). Therefore, the RBP cut-off of <0.71 µmol/L was used to indicate vitamin A deficiency among children. The correlation agreed with past research, showing that RBP behaves like serum retinol and can be used as an indicator of vitamin A status (Gorstein, 2008).

Unlike ferritin, which is elevated during infection/inflammation, retinol binding protein is a negative acute phase reactant, meaning it decreases during infection/inflammation (Thurnham 2003; Thurnham 2010). To account for the presence of inflammation, the prevalence of vitamin A deficiency was calculated for all participants (with and without evidence of inflammation) as well as for those participants without evidence of inflammation (i.e. after exclusion of those with high CRP and/or AGP). Table 7-4 below shows the prevalence of vitamin A deficiency among all children and among those without evidence of inflammation, stratified by age, gender, and place of residence. Among all children (n=2148), the prevalence of VAD was 15.6% (95% CI: 13.1, 18.1). Among children without presence of inflammation (n=1436), the prevalence of VAD dropped to 7.8% (95% CI: 5.7, 9.8). The prevalence of vitamin A deficiency (VAD) was 15.6%. Among children 6.0 -11.9, 12.0-17.9, 18.0-23.9, and 24.0-29.9 months, the prevalence of VAD was 16.0%, 13.7%, 16.1%, and 16.7%, respectively. The prevalence of deficiency was 17.0% among males and 14.1% among females, and was 16.0% in urban and 14.7% in rural areas.

In interpreting the results of vitamin A deficiency it is important to recognize that compared to the VITAL-EQA program mean, the precision and bias were Minimal or Unacceptable for RBP (>80-85% precision of the VITAL-EQA results, with 18.3% bias (see Appendix VII). The RBP concentrations for the EQA samples measured by the VitMin Lab were 15 to 20% higher when compared to the target values. Thus, the actual prevalence of vitamin A deficiency could likely be higher than the prevalence in this report. Unfortunately, there is no accepted quantitative method for adjusting the values according to imprecision and bias

The WHO classifies the level of public health significance of VAD based on the prevalence of VAD among preschool-age children in a population. According to the classification, prevalence of VAD of  $\geq 10\%$  to  $< 20\%$  would be classified as a moderate public health problem (WHO, 2009). Again, it should be noted that the WHO criteria apply to population estimates for all children less than 71 months of age. Because the prevalence of VAD has been shown to decrease with increasing age in Kyrgyzstan (CDC/UNICEF/MOH, 2012), it is likely that the prevalence of VAD deficiency in all preschool children in Kyrgyzstan would be lower than that measured in children 6-29 months (CDC, 2012).

**Table 7 4: Prevalence of vitamin A deficiency (VAD) among children aged 6.0–29.9 months, stratified by age, gender, residence, and presence of inflammation — Kyrgyzstan, 2013**

Characteristics of Child	n	All Participants % with VAD (95% CI)	n	Participants Without Inflammation <sup>a</sup> % with VAD (95% CI)
<b>Age of children</b>				
6.0-11.9 months	517	16.0 (11.1 - 21.0)	354	9.5 (5.3 - 13.8)
12.0-17.9 months	568	13.7 (9.9 - 17.5)	381	4.0 (1.6 - 6.5)
18.0-23.9 months	537	16.1 (11.2 - 20.9)	354	8.2 (3.2 - 13.2)
24.0-29.9 months	526	16.7 (12.2 - 21.3)	347	9.6 (5.1 - 14.1)
<b>Gender of children</b>				
Male	1093	17.0 (13.1 - 20.9)	724	7.7 (4.0 - 11.4)
Female	1055	14.1 (11.1 - 17.1)	712	7.8 (5.5 - 10.1)
<b>Place of Residence</b>				
Urban	549	16.0 (12.4 - 19.5)	361	7.7 (4.8 - 10.6)
Rural	1599	14.7 (12.9 - 16.6)	1075	7.9 (6.3 - 9.5)
<b>Total</b>	<b>2148</b>	<b>15.6 (13.1 - 18.1)</b>	<b>1436</b>	<b>7.8 (5.7 - 9.8)</b>

NOTE: VAD=Vitamin A Deficiency (RBP  $< 0.71 \mu\text{mol/L}$ ); CI=confidence interval; 95% CI's adjusted for survey design. <sup>a</sup>Inflammation not present (low C-reactive protein [CRP $\leq 5 \text{ mg/L}$ ] and low  $\alpha 1$ -glucoprotein acid [AGP $\leq 1.0 \text{ g/L}$ ]).

---

## USE OF MICRONUTRIENT SUPPLEMENTS

Among children surveyed whose mother was interviewed (n=1823), 34.6% (95% CI: 29.5, 40.2) had ever received an anemia diagnosis from a health care worker. Among children with a history of anemia (n=573), 62.6% (95% CI: 56.7, 68.2) were reported to have taken iron syrup or drops. Among all children, 21.7% (95% CI: 17.9, 25.9) were reported to have received iron tablets or syrup to improve anemia status.

---

## CHAPTER 8

# COMPARISON OF NUTRITIONAL STATUS OF CHILDREN 6–29 MONTHS AND INFANT YOUNG CHILD NUTRITION KNOWLEDGE, ATTITUDES, AND PRACTICES — 2009 AND 2013 NATIONAL SURVEYS

### BACKGROUND

This chapter compares results of the 2009 and 2013 surveys in regard to anthropometry, hemoglobin, and other biochemical indicators in the population of children aged 6–29 months living in the Kyrgyz Republic, as well as infant and young child nutrition (IYCN) knowledge, attitudes, and practices.

### METHODS

Both the 2009 and 2013 surveys used similar methodology for sample selection and measurement of anthropometry, hemoglobin, biochemical indicators, and IYCN indicators. In both the 2009 and 2013 surveys, the sample population was selected using population proportionate to size (PPS) methods, and information was collected from 66 and 80 clusters, respectively. Clusters were formed based on the catchment areas for health centers (based on geographic location of residence, all children are assigned to a primary health care center). The sampling frame for the 2009 survey was comprised of children 6–59 months and their mothers and was stratified by rural and urban residence. For each stratum, 33 clusters were selected for the survey with 30 children invited to participate in each cluster. The sampling frame for the 2013 survey consisted of children 6–29 months of age living in the Kyrgyz Republic (with the exception of the city of Bishkek). Although micronutrient powder was only distributed to children 6–24 months of age, we anticipated that the effects of the Gulazyk on iron status may have persisted 6 months of age after discontinuation (Ip 2009). To increase our sample size for the 2009 survey comparison, we included children 24 to 29 months of age. The 2013 sample was not stratified; 80 clusters of 30 children each (instead of 30 children, one cluster had 22 children and another had 29 children) were selected for the 2013 survey. Detailed methods for the 2013 survey including sample size calculations are described in Chapter 2 of this report. Detailed methods for the 2009 report have been previously published (CDC/UNICEF/MOH, 2012).

For the 2009 and 2013 surveys, data were collected during June and July, and July–September, respectively. Caretakers of selected children were invited to come to their primary health care center on a predetermined day. If children did come, survey personnel visited the home. In the 2009 survey, blood was collected from both mothers and children; therefore, mothers were actively recruited to accompany their children. In the 2013 survey, blood was collected only from children and mothers were not actively recruited to accompany their children. The 2009 and 2013 questionnaires were similarly worded. Both contained modules on sociodemographic information and the use of micronutrient powders (see Appendix I and VIII). Both questionnaires were written in English and then translated into the Kyrgyz and Russian languages. Comparisons between 2009 and 2013 data are made based on children 6–29 months of age, excluding children residing in the city of Bishkek and limiting the 2009 age range to children 6–29 months.

The anthropometric equipment and methods used to measure children were the same for the 2009 and 2013 surveys. During both surveys, Shorr boards were used to measure weight/length and UNICEF Seca Uniscales were used to measure weight. For children  $\geq 24$  months standing height was measured. For children  $< 24$  months supine length, rather than height, was measured<sup>8</sup>. The same instructor conducted training for both surveys. The age in months of the participant was calculated by

---

<sup>8</sup> For the purpose of this report, the term height will be used to indicate either length or height.



---

subtracting the date of measurement from the birth date and dividing by 30.4. The WHO growth reference (2006) was used to calculate anthropometric indicators (Z-scores) of height-for-age, weight-for-age, and weight-for-height.

Records with potentially erroneous data were excluded from the analysis based on standard Z-score cutoffs developed by WHO (WHO, 1995) (see Chapter 6). In the 2009 survey, anthropometric measurement (weight and height) were taken on 576 participants 6-29 months of age and living outside of Bishkek City. Of these, 3 participants were excluded based on out of range Z-scores. Of the 2162 participants from whom anthropometric measurements were taken in the 2013 survey, 7 participants were excluded based on out of range Z-scores.

A measure of the quality of the anthropometric measurements is the unweighted SD of the Z-score distribution (WHO, 1995). For the 2009 survey, among children 6-29 months of age and living outside of Bishkek City, the unweighted SD's for both WHZ and WAZ fall within the acceptable range for data quality; however, the SD for HAZ falls outside the acceptable range and measurements may be slightly inaccurate (most likely height/length measurements or inaccurate age determination) [HAZ: SD 1.331; WHZ: SD 0.986; WAZ: SD 1.018] (CDC/UNICEF/MOH, 2012). Although the SD for HAZ is wider than expected based on 1995 WHO, these guidelines were based on an earlier growth reference and it has been shown that higher standard deviations may be expected based on the 2006 WHO growth reference (Mei, 2007). For the 2013 survey, the unweighted SD's for all anthropometry indicators fall within the acceptable range [HAZ: SD 1.27; WHZ: SD 1.04; WAZ: SD 1.06] (see Chapter 6 of this report).

### **Biochemical indicators**

In both the 2009 and 2013 surveys, hemoglobin was assessed in the field using the HemoCue<sup>®</sup> photometric instrument (Model 301, HemoCue AB, Angelholm, Sweden). Laboratory personnel collected capillary blood samples through a finger stick using a single-use retractable lancet. After the first two drops were wiped clean, the third drop was drawn into a HemoCue<sup>®</sup> cuvette. Afterwards, 500  $\mu$ L of blood was collected in a Microtainer<sup>®</sup>. Analysis of biochemical indicators was conducted using the "sandwich assay" at the Erhardt VitA-Iron Tech research laboratory (Erhardt, 2004). The biochemical indicators measured were iron status (serum ferritin, soluble transferrin receptor [sTfR]) and inflammation status (C-reactive protein [CRP] and  $\alpha$ 1-glycoprotein acid [AGP]).

Anemia was defined as an altitude-adjusted hemoglobin concentration of <11.0 g/dL (Sullivan 2008, WHO 2001). Total iron deficiency was defined as either decreased serum ferritin concentration (<12  $\mu$ g/L) or increased sTfR levels (>8.3 mg/L). Iron deficiency anemia was defined as having both a low hemoglobin value and either low serum ferritin or high sTfR. Inflammation was considered present if either indicator was elevated (CRP>5.0 mg/L or AGP>1.0 g/L) (Thurnman 2003, Thurnman 2010).

RBP was used as an indicator of vitamin A status (Gorstein 2008). The CDC Nutrition Laboratory compared RBP and plasma retinol on a subsample of participants in the 2009 survey; RBP concentration less than 0.71  $\mu$ mol/L was determined as the cut-off for vitamin A deficiency (personal communication, Rosemary Schleicher).

### **External Quality Assurance**

Analysis of biochemical indicators was conducted using the "sandwich assay" at the Erhardt VitMin Laboratory (Erhardt, 2004). The laboratory measured ferritin, soluble transferrin receptor (sTfR), 1-acid glycoprotein (AGP), C-reactive protein (CRP), and retinol binding protein (RBP) concentrations in plasma using an enzyme-linked immunosorbent assay (ELISA) technique. During both surveys, the laboratory (Willstaett, Germany) participated in CDC's VITAL-EQA (Vitamin A Laboratory-External Quality Assurance) program, an external quality assurance program that assesses laboratory performance during the course of analyzing survey samples (<http://www.cdc.gov/labstandards/vitaleqa.html>; Haynes, 2008) (see Appendix VII). For the 2009 survey, the precision and bias were Optimal or Desirable for all of the above indicators (>90% precision of the VITAL-EQA results, with <0.5% bias). For the 2013 survey, the precision and bias were Optimal or Desirable for ferritin, sTfR, and CRP (>90% precision of the VITAL-EQA results, with <0.5% bias). However, the precision and bias shifted 15-20%

---

for RBP (>80-85% precision of the VITAL-EQA results, with 18.3% bias) due to a change in pools used by the VITAL-EQA program. In regard to internal quality control, the inter-assay coefficients of variation (CV) were 3.0% in for the 2009 survey and 3.8% for the 2013 survey. A CV of about 10% provides acceptable precision during an ELISA technique (Erhardt, 2004; Haynes, 2008).

The questions on infant and young child feeding were the same in 2009 and 2013. Questions on feeding practices were derived from a consensus meeting on indicators and held in 2007 (WHO, 2007) and the WHO Indicators for Assessing Infant and Young Child Feeding Practices (2008). Because children under 6 months of age were not included in the survey, we could not use the standard indicators for exclusive breastfeeding and age-appropriate breastfeeding. Instead we used maternal recall to estimate these indicators see Chapter 5, Table 5-1). In the 2009 baseline survey only mothers answered the infant feeding questions, whereas in 2013, all caretakers answered these questions. For consistency, the 2013 infant feeding analysis was also restricted to only those interviews where the mother was the respondent.

As the 2013 survey included only children 6-29 months living outside of Bishkek City, in the 2009 survey only children 6-29 months living outside Bishkek City were included in the analysis. Because children were first enrolled in the IYCF/MNP program either at the first well child check-up after they reached 6 months of age or when the health care worker visited them at home, we expected that children in the 6-11 months age category would have had a shorter degree of exposure to the micronutrient powder program than older children (i.e. they could not have been in the program for one full year). Children 24-29 months were no longer eligible for the micronutrient powder program, but we expected that MNP would have affected iron status for an additional 6 months after discontinuation (Ip 2009). Therefore, analyses were stratified by age (6-11, 12-17, 18-23 and 24-29 months of age). Because serum ferritin and RBP are acute-phase reactants, all biochemical results are presented for the total population and for the population without inflammation.

Data analysis was conducted using SPSS (version 20.0, USA). Bivariate statistical testing was used to determine the significance of differences in percent deficient between the 2009 and 2013 surveys. The 2009 survey was weighted to account for stratification (rural, urban) and non-response. The 2013 survey was weighted for non-response only (the sample was not stratified). P-values for the Pearson chi-square test are presented to indicate whether there are statistically significant differences in these indicators between the 2009 and 2013 surveys. The Pearson chi-square tests were run using methods that account for the design of the two surveys (standard errors were adjusted for the cluster survey design).

## RESULTS

Comparisons between 2009 and 2013 data are made based on children 6–29 months of age, excluding children residing in the city of Bishkek. Table 8-1 compares various characteristics of the children living in the Kyrgyz Republic who were surveyed in 2009 and 2013. The proportion of males and females was similar. For both surveys, participation was fairly even across the four age groups. In the 2009 survey nearly all mothers were present for the interview (99.9%) whereas in 2013, 86.2% of mothers were present. The majority of mothers in both surveys had completed at least a secondary education. The prevalence of inflammation was higher in the 2013 survey compared to the 2009 survey: 22.3% of children had inflammation in the 2009 survey and 33.9% had inflammation in 2013.

**Table 8 1: Characteristics of children 6–29 months and their mothers — Kyrgyzstan, 2009 and 2013**

Characteristics	2009		2013		p- value <sup>c</sup>
	n <sup>b</sup>	% (95% CL)	n <sup>b</sup>	% (95% CL)	
<b>Age of children</b>	576		2156		0.822
6-11 months		24.7 (20.5 – 29.5)		22.7 (20.6 - 25)	
12-17 months		23.5 (20.1 – 27.2)		26.5 (24 - 29.3)	
18-23 months		25.4 (21.8 – 29.5)		25.1 (22.9 - 27.5)	
24-29 months		26.3 (22.1 - 31.0)		25.6 (23.5 - 27.7)	
<b>Gender of children</b>	576		2156		0.946
Male		50.5 (45.8 - 55.2)		50.7 (47.3 - 54.1)	
Female		49.5 (44.8 – 54.2)		49.3 (45.9 - 52.7)	
<b>Relation of person interviewed to child<sup>d</sup></b>	576		2156		
Mother		99.9 (99.2 - 100)		86.2 (83.8 – 88.3)	0.000
Grandmother		0.1 (0.0 – 0.8)		7.7 (6.2 – 9.6)	
Other				6.1 (4.4 – 8.4)	
<b>Mother's education<sup>e</sup> (highest level completed)</b>	575		1823		0.038
Less than secondary		8.0 (4.6 – 13.7)		14.1 (8.9 - 21.7)	
Secondary		55.8 (49.8 – 61.6)		47.5 (42.7 - 52.5)	
Technical		16.0 (12.1 – 21.0)		12.6 (9.9 - 15.8)	
Higher education		20.1 (16.3 - 24.5)		25.8 (21.1 - 31.1)	
<b>Inflammation status of children</b>	576		2156		0.000
Have inflammation		22.3 (18.9 - 26.2)		33.9 (30.8 - 37.1)	
No inflammation <sup>f</sup>		77.7 (73.8 – 81.1)		66.1 (62.9 - 69.2)	

NOTE: CI=confidence interval; 95% CI's adjusted for survey design.

<sup>a</sup>Calculated for children who had valid measurements for anthropometry, hemoglobin, and/or other biochemical indicators. (If a child had a valid measurement for at least one of these indicators they were included in this description of the sample).

<sup>b</sup>n's are unweighted denominators; subgroups that do not sum to the total have missing data

<sup>c</sup>Bivariate tests of statistical significance were conducted using the Pearson chi-square test, accounting for the cluster survey design.

<sup>d</sup>In the 2009 survey, mothers were strongly encouraged to be present; in the 2013 survey either mothers or caretakers could be present for the interview.

<sup>e</sup>Mothers education was only reported by mothers who were present for the interview.

<sup>f</sup>Inflammation not present (low C-reactive protein [CRP≤5 mg/L] and low α1-glucoprotein acid [AGP≤1.0 g/L]).

### Anthropometry

Table 8-2 compares the prevalence of wasting, stunting, and underweight among children who were surveyed in 2009 and 2013. Between the 2009 and 2013 surveys, the prevalence of wasting (weight-for-height Z <-2.0) remained stable at ≤2.0% (p-value: 0.895). The prevalence of stunting (height-for-age Z <-2.0) decreased from 19.6% to 11.7% (p-value: 0.000). The largest percentage decline was in the 24 to 29 month age group where the percentage of stunting declined from 37.1% to 15.5%. It should also be noted that age pattern for stunting differed between the two surveys. In 2009 the prevalence of stunting doubled from 19.6% at 18-23 months to 40.3% at 24 to 29 months of age; whereas in 2013, the prevalence of stunting was similar across these same age groups. The prevalence of underweight (weight-for-age Z <-2.0) was similar between the two surveys, with 4.1% of children underweight in 2009 and 4.8% in 2013 (p-value: 0.701).

**Table 8 2: Anthropometric characteristics by age among children 6–29 months — Kyrgyzstan, 2009 and 2013**

Characteristics	2009		2013		p- value <sup>a</sup>
	N	% (95% CI)	N	% (95% CI)	
<b>Wasted</b> (weight-for-height Z<-2.0)	502	1.9 (0.7 – 3.1)	2160	2.0 (1.1 - 2.8)	0.893
6-11 months	131	1.9(0.0 – 4.1)	520	2.7 (0.8 - 4.5)	0.568
12-17 months	120	2.8(0.0 – 6.1)	568	2.5 (0.8 - 4.2)	0.866
18-23 months	123	0.9(0.0 - 2.7)	539	2.3 (0.5 - 4.2)	0.238
24-29 months	128	2.1(0.0 – 4.5)	533	0.5 (0.1 - 0.8)	0.168
<b>Stunted</b> (height-for-age Z<-2.0)	502	19.6 (15.7 – 23.5)	2155	11.7 (9.3 - 14.1)	0.001
6-11 months	131	6.9(2.8 – 11.2)	515	5.0 (2.3 - 7.6)	0.451
12-17 months	121	16.9(9.9 - 23.8)	568	8.6 (5.5 - 11.6)	0.032
18-23 months	123	18.5(10.2 – 26.9)	539	17.1 (12.5 - 21.7)	0.772
24-29 months	127	37.1(28.4 – 45.8)	533	15.5 (10.9 - 20.1)	0.000
<b>Underweight</b> (weight-for-age Z<-2.0)	505	4.1 (2.1 – 6.0)	2162	4.8 (3.7 - 5.9)	0.537
6-11 months	132	2.6(0.0 - 5.4)	520	3.4 (0.9 - 6)	0.671
12-17 months	122	3.8(0.1 - 7.5)	569	3.7 (1.9 - 5.6)	0.962
18-23 months	123	1.7(0.0 – 4.3)	539	6.3 (3.4 - 9.3)	0.015
24-29 months	128	8.2(3.1 – 13.3)	534	5.6 (3.1 - 8.1)	0.367

NOTE:CI=confidence interval; 95% CI's adjusted for survey design.

Anthropometric values based on WHO growth reference curves (WHO, 2006).

aBivariate tests of statistical significance were conducted using the Pearson chi-square test, accounting for the cluster survey design.

### Biochemical indicators

Table 8-3 compares 2009 and 2013 prevalence figures for anemia, iron deficiency, and vitamin A deficiency among surveyed children 6–29 months of age living in the Kyrgyz Republic. The prevalence figures are presented for all survey participants, and for those without inflammation. Among all participants, the prevalence of anemia was 38.7% in 2009 and 32.7% in 2013 (p-value: 0.116). Among participants without inflammation, the prevalence of anemia decreased from 37.4% to 29.4% (p-value: 0.038).

Iron deficiency prevalence was analyzed separately based on each of the two biomarkers for deficiency, serum ferritin and sTfR. Among all participants, the prevalence of iron deficiency (defined by low ferritin) decreased from 50.6% to 34.2% (p-value: <0.001). Among those without inflammation, the prevalence of iron deficiency (defined by low ferritin) decreased from 56.7% to 39.7% (p-value: <0.001). Among all participants, the prevalence of iron deficiency (defined by high sTfR) decreased from 48.9% to 39.3% (p-value: 0.007). Among those without inflammation, the prevalence of iron deficiency (defined by high sTfR) decreased from 48.0% to 38.9% (p-value: 0.021). Among all participants 6-29 months of age, the prevalence of iron deficiency anemia (defined as low hemoglobin and either low ferritin or high sTfR) decreased from 31.9% to 23.6% (p-value: 0.014). Among those without inflammation, the prevalence of iron deficiency anemia (defined using the same criteria) decreased from 31.7% to 22.4% (p-value: 0.009). Among all participants, the prevalence of vitamin A deficiency increased from 6.2% to 15.6%; p-value: <0.001). Among those without inflammation the prevalence of vitamin A deficiency increased from 3.3% to 7.8%; p-value: 0.001). (Appendix IX shows the anthropometric and biochemical findings of children 6 to 24 months in 2009 and 2013).

### Table 8 3: Prevalence of anemia, iron deficiency, and vitamin A deficiency by age among chil-

**dren 6–29 months —**

Characteristics	All Participants				Participants without Inflammation				
	2009		2013		2009		2013		
	N	% (95% CI)	N	% (95% CI)	N	% (95% CI)	N	% (95% CI)	
<b>Anemia<sup>a</sup></b>									
6-11 months	576	38.7(32.2 - 45.2)	2156	32.7(28.9 - 36.4)	447	0.116	1436	29.4(25.7 - 33.1)	<b>0.038</b>
12-17 months	137	40.7(29.9 - 51.5)	519	35.6(27.8 - 43.4)	112	0.450	354	33.6(25.4 - 41.7)	<b>0.831</b>
18-23 months	135	42.9(30.5 - 55.3)	568	35.8(30.5 - 41.0)	101	0.299	381	33.3(27.5 - 39.1)	<b>0.402</b>
24-29 months	148	37.4(29.5 - 44.7)	538	35.0(30.1 - 39.9)	111	0.601	354	29.2(23.6 - 34.7)	<b>0.039</b>
	156	34.3(24.6 - 43.9)	531	24.5(17.7 - 31.2)	123	0.102	347	21.7(15.6 - 27.8)	<b>0.046</b>
<b>Low ferritin (&lt;12 µg/L)</b>									
6-11 months	576	50.6(45.8 - 55.4)	2156	34.2(31.3 - 37.2)	447	0.000	1436	39.7(36.4 - 43.1)	<b>0.000</b>
12-17 months	137	44.9(35.1 - 54.7)	519	29.9(23.7 - 36.1)	112	0.012	354	34.2(25.5 - 42.9)	<b>0.041</b>
18-23 months	135	56.5(46.2 - 66.8)	568	34.9(29.0 - 40.8)	101	0.000	381	44.3(37.6 - 50.9)	<b>0.018</b>
24-29 months	148	51.3(42.5 - 60.1)	538	35.3(30.1 - 40.6)	111	0.002	354	38.8(33.2 - 44.4)	<b>0.000</b>
	156	49.9(41.9 - 57.7)	531	36.3(30.8 - 41.9)	123	0.006	347	41.1(34.1 - 48.1)	<b>0.006</b>
<b>High sTfR (&gt;8.3 mg/L)</b>									
6-11 months	576	48.9(43.1 - 54.7)	2156	39.3(35.6 - 43.1)	447	0.007	1436	38.9(34.6 - 43.1)	<b>0.021</b>
12-17 months	137	53.2(44.3 - 62.1)	519	38.5(33.0 - 44.1)	112	0.006	354	37.7(30.8 - 44.6)	<b>0.031</b>
18-23 months	135	54.9(45.5 - 64.4)	568	43.7(36.8 - 50.5)	101	0.060	381	43.7(35.7 - 51.6)	<b>0.174</b>
24-29 months	148	48.2(39.5 - 56.9)	538	38.4(32.7 - 44.2)	111	0.065	354	36.6(29.4 - 43.7)	<b>0.050</b>
	156	40.2(30.8 - 49.6)	531	36.4(30.5 - 42.3)	123	0.499	347	37.3(30.5 - 44.2)	<b>0.571</b>
<b>Iron deficiency anemia<sup>c</sup></b>									
6-11 months	576	31.9(26.3 - 37.5)	2156	23.6(20.1 - 27.1)	447	0.014	1436	22.4(18.7 - 26.1)	<b>0.009</b>
12-17 months	137	32.4(22.8 - 41.9)	519	24.1(16.9 - 31.4)	112	0.173	354	22.3(14.3 - 30.3)	<b>0.312</b>
18-23 months	135	38.1(27.3 - 49.0)	568	25.7(20.8 - 30.6)	101	0.041	381	26.4(19.9 - 32.9)	<b>0.184</b>
24-29 months	148	30.0(22.8 - 37.2)	538	26.0(21.8 - 30.3)	111	0.346	354	22.9(18.3 - 27.4)	<b>0.026</b>
	156	27.8(19.1 - 36.4)	531	18.6(12.6 - 24.7)	123	0.087	347	17.8(11.1 - 24.5)	<b>0.078</b>
<b>Vitamin A deficiency<sup>e</sup></b>									
6-11 months	576	6.2(4.2 - 8.2)	2148	15.6(13.1 - 18.1)	447	0.000	1436	7.8(5.7 - 9.8)	<b>0.001</b>
12-17 months	137	9.2(4.8 - 13.6)	517	16.0(11.1 - 20.9)	112	0.007	354	9.5(5.3 - 13.8)	<b>0.155</b>
18-23 months	135	4.0(0.7 - 7.3)	568	13.7(10.0 - 17.5)	101	0.000	381	4.0(1.6 - 6.5)	<b>0.590</b>
24-29 months	148	7.2(2.8 - 11.5)	537	16.1(11.2 - 20.9)	111	0.000	354	8.2(3.2 - 13.2)	<b>0.039</b>
	156	4.5(1.0 - 7.9)	526	16.7(12.2 - 21.3)	123	0.000	347	9.6(5.1 - 14.1)	<b>0.008</b>

NOTE:CI=confidence interval; 95% CI's adjusted for survey design.

<sup>a</sup>Inflammation not present (low C-reactive protein [CRP≤5 mg/L] and low α1-glucoprotein acid [AGP≤1.0 g/L]).

<sup>b</sup>Anemia: hemoglobin <11.0 g/dL adjusted for altitude.

<sup>c</sup>Iron deficiency anemia: Hemoglobin <11.0 g/dL and low plasma ferritin (<12 µg/L) or high sTfR (>8.3 mg/L).

<sup>d</sup>Bivariate tests of statistical significance were conducted using the Pearson chi-square test, accounting for the cluster survey design.

<sup>e</sup>Vitamin A deficiency, defined as RBP <0.71 µmol/L.



## Vitamin A Supplement Use

In the 2009 survey (before the vitamin A capsule program was discontinued), among children 6-29 months of age whose mother were surveyed (n=575), 94.2% (95% CI: 91.7 – 96.0) had ever received a vitamin A capsule. Among children who had received a vitamin A capsule, 66.8% (95% CI: 55.4 – 76.5) had received a capsule within 2 months of the survey.

## IYCN Knowledge, Attitudes, and Practices

Table 8-4 compares changes in IYCN practices among children 6–23.9 months between the 2009 and 2013 surveys. Although the placement of the infant feeding module was different in the two surveys, the questions were worded identically. In the 2009 survey only mothers answered the infant feeding questions whereas in 2013, all caretakers answered the questions. For analytic purposes, the infant feeding analysis was restricted to interviews where the mother was the respondent. The definitions for IYCN indicators can be found in Tables 5.1 and 5.2. The algorithm for analysis was the same in both survey years (Appendix V).

Among children 6–23.9 months of age, the prevalence of children breastfed within the first hour after birth increased from 71.1% to 85.1% (p-value: <0.001); the prevalence of children ever breast fed was similar (98.1% vs 99.5%, respectively). An increase was observed in the prevalence of children who were exclusively breastfed during the first 6 months of life (22.1% vs. 40.2%, respectively, p-value: <0.001). The prevalence of children breastfeeding at 1 year was (72.4% vs. 80.1%, p-value: 0.320) and, at 2 years was 13.3% vs.26.2 %, p-value: 0.028.

Appropriate introduction of solid, semi-solid or soft foods increased from 79.0% to 90.9% (p-value: 0.051). The prevalence of children who met the criteria for minimum dietary diversity (four or more food groups) increased from 74.5% to 86.8% (p-value: 0.004) and the prevalence of children who met criteria for minimum meal frequency increased from 64.9% to 74.7% (p-value: 0.012). After combining the indicators of minimum dietary diversity and minimum meal frequency the prevalence of children reaching the criteria for minimum acceptable diet increased from 51.3% to 67.2% (p-value: < 0.001 ). Among children who were appropriately breastfed, the prevalence of age-appropriate breast feeding increased from 50.0% to 62.8% (p-value: <0.001). The prevalence of adequate milk feeding frequency (at least 2 milk feedings a day among non-breast fed children was 62.6% to 65.4%, respectively (p-value: 0.684).

**Table 8 4: Changes in IYCN practices among children 6-23.9 months as reported by their mothers — Kyrgyzstan, 2009 and 2013**

	2009		2013		p- value <sup>a</sup>
	N	%, mean (95% CI)	N	%, mean (95% CI)	
<b>IYCN Indicators, % (Children 6–23.9 months)</b>					
Early initiation of breastfeeding	419	71.1 (64.5 – 77.6)	1416	85.1 (82.1 - 88.1)	0.000
Exclusive breastfeeding under 6 months	410	22.1 (15.7 – 28.5)	1410	40.2 (34.9 – 45.6)	0.000
Ever breastfed	419	98.1 (96.3 - 99.9)	1416	99.5 (98.9 - 100.0)	0.144
Continued breastfeeding at 1 year	94	72.4 (59.0 – 85.7)	348	80.1 (72.7 - 87.5)	0.320
Continued breastfeeding at 2 years	96	13.3 (3.1 – 23.5)	295	26.2 (20.8 - 31.5)	0.028
Appropriate introduction of solid, semi-solid or soft foods	51	79.0(68.1 – 90.0)	212	90.9 (86.2 - 95.7)	0.051
Consuming minimum dietary diversity	419	74.5 (66.6 – 82.4)	1416	86.8 (84.1 – 89.6)	0.004

---

Consuming minimum meal frequency	419	64.9 (58.3 – 71.4)	1416	74.7 (70.9 - 78.5)	0.012
Consuming minimum acceptable diet	419	51.3 (43.6 – 58.9)	1416	67.2 (63.1 – 71.3)	0.000
Age-appropriate breastfeeding	419	50.0 (44.5 – 55.4)	1416	62.8 (59.3 - 66.3)	0.000
Adequate milk feeding frequency for non-breastfed Children	232	62.6 (52.5 – 72.8)	637	65.4 (56.4 - 74.4)	0.684

NOTE:CI=confidence interval; 95% CI's adjusted for survey design. Definitions for indicators can be found in Table 5.1 and 5.2.

<sup>a</sup>Bivariate tests of statistical significance were conducted using the Pearson chi-square test, accounting for the cluster survey design

---

## DISCUSSION

In June 2009, the government of the Kyrgyz Republic introduced an integrated IYCN/MNP program. Thus far, two nationally representative surveys (with the exclusion of Bishkek) to assess the micronutrient status of Kyrgyz children were conducted in June 2009 and August/September 2013. Among all children, statistically significant declines were observed in the prevalence of iron deficiency (as measured by either serum ferritin or sTfR) and iron deficiency anemia. Among children without evidence of inflammation, a similar magnitude of decline in prevalence of anemia, iron deficiency and iron deficiency anemia was observed. The percentage point declines observed between the 2009 and 2013 surveys was similar to those observed before and after a pilot home fortification program in the Talas oblast where declines were observed in the prevalence of anemia (50.6% versus 43.8%,  $p=0.05$ ); total iron deficiency (either low ferritin or high sTfR) (77.3% versus 63.7%,  $p<0.01$ ); and iron deficiency anemia (45.5% versus 33.4%,  $p<0.01$ ) (Serdula 2013).

Between 2009 and 2013, the prevalence of wasting among children remained stable while the prevalence of stunting decreased from 19.6% to 11.7%. The decrease in the prevalence of stunting from 2009 to 2013 was surprising because in the Talas Oblast pilot program, the prevalence of stunting increased from 10.7% to 17.0% (CDC, 2011). The largest difference in stunting prevalence between the 2009 and 2013 surveys was seen among children 24 to 29 months of age. It should also be noted that the age pattern of stunting differed between the two surveys. In 2009, the prevalence of stunting nearly doubled from 18 to 23 months to 24 to 29 months whereas in 2013 the prevalence of stunting was similar across these age categories. Although the anthropometric equipment and training was the same in the 2009 and 2013 surveys and the variance (SD) in measurements was similar, it is still possible that the decrease in stunting may be due to differences in measurement or ascertainment of age.

Unexpectedly, the prevalence of vitamin A deficiency was significantly higher at follow-up compared to the baseline. Given the positive laboratory bias observed by comparison with the VITAL-EQA standard, the actual prevalence of vitamin A deficiency could likely be higher than the reported prevalence in 2013. Unfortunately, there is no accepted quantitative method for adjusting the values. The laboratory internal QC however is acceptable (3.8% for the 2013 survey), where a CV of about 10% provides acceptable precision during an ELISA technique (Erhardt, 2004; Haynes, 2008). The reasons for the increase in prevalence of vitamin A deficiency remain unclear as previous micronutrient powder interventions have shown either no change or an improvement in vitamin A deficiency. The previous pilot intervention in one district of the Kyrgyz Republic showed no significant impact of home fortification on vitamin A status (Serdula 2013).

Because of the low prevalence of vitamin A deficiency in the 2009 survey, the Ministry of Health discontinued its routine vitamin A capsule distribution program. Although Vitamin A capsule distribution programs have been found to only transiently (<2 months) increase the distribution of serum retinol (Palmer 2102), the capsule distribution program may still have had an effect on retinol levels at baseline. In the 2009 survey, of children 6 to 29 months, 94% were reported to have received vitamin A capsules, 66.8% of whom had received capsules in the previous 2 months. Nonetheless, because vitamin A was included in the MNP, discontinuation of the capsule program would not have been expected to result in an increased prevalence of vitamin A deficiency.

IYCN indicators showed a general improvement. During both survey years, breastfeeding was initiated for almost all children. Comparing the 2009 and 2013 surveys, the prevalence of exclusive breastfeeding under 6 months increased nearly doubled from 22% to 40% and early initiation of breastfeeding increased from 71% to 85%. It is possible that at least some of these differences may have resulted from the IYCN program. However, it should also be noted that, although the IYCN questions were worded identically, the placement of the infant feeding module was different in the two surveys.



---

In any case, causal inferences cannot be made in comparing the two cross-sectional surveys. Although the demographic characteristics were similar between the baseline and follow-up surveys, without a control population, it is not possible to account for secular changes. During the four years between surveys, changes independent of the micronutrient program could have affected population micronutrient and anthropometric status. Government instability and decreased access to food caused by the increase in food prices which began in 2010, may have influenced the micronutrient status in the population (Ortiz, 2011). Furthermore, we cannot distinguish the effects of the nutrition education program from the micronutrient powder program. Strengths of our design are that indicators of iron and vitamin A status were measured and that inflammatory markers were measured to adjust for potential differences in inflammation status of the population in the pre- and post-surveys.

The Kyrgyz Republic will be among the first countries in the world to document improvement in iron status before and after implementing an integrated IYCN/MNP home fortification program on a national scale (with the exception of Bishkek). Our pre and post surveys showed a reduction in iron deficiency in the Kyrgyz Republic after this national-scale micronutrient powder program was implemented through a primary health care system in combination with an IYCN health education program and extensive community mobilization.

---

## REFERENCES

1. CDC, UNICEF, MOH Kyrgyz Republic. Assessment of Nutritional Status of Children 6-24 Months of Age and Their Mothers, Rural Talas Oblast, Kyrgyzstan, 2008 Report. Bishkek: U.S. Centers for Disease Control and Prevention, UNICEF-Kyrgyzstan, Ministry of Health of the Kyrgyz Republic, National Statistics Committee of the Kyrgyz Republic. 2010.
2. CDC, UNICEF, MOH Kyrgyz Republic. Follow-up survey of nutritional status in children 6-24 months of age, Talas Oblast, Kyrgyz Republic, 2010 Report. Bishkek: U.S. Centers for Disease Control and Prevention, UNICEF-Kyrgyzstan, Ministry of Health of the Kyrgyz Republic, National Statistics Committee of the Kyrgyz Republic. 2011.
3. CDC, UNICEF, MOH Kyrgyz Republic. National Lot Quality Assurance Sampling (LQAS) Survey of Gulazyk Program: A Micronutrient Powder-Home Fortification Program for children 6–24 months of age, Kyrgyzstan, 2011 Report. Bishkek: U.S. Centers for Disease Control and Prevention, UNICEF-Kyrgyzstan, Ministry of Health of the Kyrgyz Republic, National Statistics Committee of the Kyrgyz Republic.
4. CDC, UNICEF, MOH Kyrgyz Republic. National survey of the nutritional status of children 6-59 months of age and their mothers, Kyrgyzstan, 2009 Report. Bishkek: U.S. Centers for Disease Control and Prevention, UNICEF-Kyrgyzstan, Ministry of Health of the Kyrgyz Republic, National Statistics Committee of the Kyrgyz Republic. 2012.
5. deBenoist B, Martines J, Goodman T. Vitamin A supplementation and the control of vitamin A deficiency: Conclusions. *Food and Nutrition Bulletin*. 2001. 22(3):335-340.
6. Erhardt JG, Estes JE, Pfeiffer CM, Biesalski HK, Craft NE. Combined measurement of ferritin, soluble transferrin receptor, retinol binding protein, and C-reactive protein by an inexpensive, sensitive, and simple sandwich enzyme-linked immunosorbent assay technique. *Journal of Nutrition*. 2004.134(11):3127-32.
7. Gorstein JL, Dary O, Pongtorn, Shell-Duncan B, Quick T, Wasanwisut E. Feasibility of using retinol-binding protein from capillary blood specimens to estimate serum retinol concentrations and the prevalence of vitamin A deficiency in low-resource settings. *Pub Health Nutrition*. 2008 May;11(5):513-20.
8. Haynes BM, Schleicher RL, Jain RB, Pfeiffer CM. The CDC VITAL-EQA program, external quality assurance for serum retinol, 2003-2006. *Clin Chim Acta*. 2008 Apr;390:90-9.
9. Ip H, Hyder SMZ, Haseen F, Rahman M, Zlotkin SH. Improved adherence and anaemia cure rates with flexible administration of micronutrient Sprinkles: a new public health approach to anaemia control. *Eur J Clin Nutr*. 2009;63:165-172.
10. Kish L. *Survey Sampling*. 1965. Wiley: New York.
11. Kyrgyz Republic, Demographic and Health Survey, 1997. Research institute of Obstetrics and Pediatrics, Ministry of Health of the Kyrgyz Republic, Macro intl. Inc., Calverton, Maryland, USA, 259p.
12. Mei Z., Grummer-Strawn LM. Standard deviation of anthropometric Z-scores as a data quality assessment tool using the 2006 WHO growth standards: a cross country analysis. *Bull World Health Organization*. 2007. 85:441-8.

- 
13. MI, CDC. Micronutrient Initiative and the CDC: Indicators and methods for cross-sectional surveys of vitamin and mineral status of populations. Micronutrient Initiative and Centers for Disease Control and Prevention. 2007.
  14. Ortiz I, Chai J, Cummins M. Escalating Food Prices: The threat to poor households and policies to safeguard a Recovery for All. United Nations Children's Fund (UNICEF), New York, 2011 [http://www.unicef.org/socialpolicy/files/Escalating\\_Food\\_Prices.pdf](http://www.unicef.org/socialpolicy/files/Escalating_Food_Prices.pdf)
  15. Palmer AC, West KP Jr, Dalmiya N, Schultink W. The use and interpretation of serum retinol distributions in evaluating the public health impact of vitamin A programmes. *Public Health Nutr.* 2012 Jul;15(7):1201-15. doi: 10.1017/S1368980012000560. Epub 2012 Mar 9. Rah JH, Christian P, Abu Ahmed S, Arju UT, Labrique AB, Rashid M. Predictors of stunting and thinness in post-menarcheal adolescent girls in rural Bangladesh. *Public Health Nutrition.* 2009. 12(12):2400-9.
  16. Robertson SE, Valadez JJ. Global review of health care surveys using lot quality assurance sampling (LQAS), 1984-2004. *Social science & medicine.* Sep 2006;63(6):1648-1660.
  17. Rosner B. *Fundamentals of Biostatistics*, 7th Ed. 2010. Brooks/Cole, Boston, MA.
  18. Serdula MK, Lundeen E, Nichols EK, et al. Effects of a large scale micronutrient powder and young child feeding education program on the micronutrient status of children 6-24 months of age in the Kyrgyz Republic. *Eur J Clin Nutr.* 2013 March. doi:10.1038/ejcn.2013.67
  19. Serdula MK, Mokdad AH, Byers T, Siegel PZ. Assessing alcohol consumption: beverage-specific vs. grouped beverage questions. *J Studies Alcohol* 1999;60:99-102.
  20. Serdula MK, Mokdad AH, Pamuk ER, Williamson DF, Byers T. Effects of question order on estimates of the prevalence of attempted weight loss. *Am J Epidemiol* 1995;142:64-7.
  21. Serdula MK, Byers T, Coates R, Mokdad A, Simoes E, Eldridge LE. Assessing intakes of high-fat foods: the effect of grouping foods into single questions. *Epidemiology* 1992;3:503-508.
  22. Sommer A and Davidson FR, Assessment and control of Vitamin A deficiency: The Anancy Accords, *J Nutr* 132: 2845S-2850S, 2002.
  23. Sullivan KM, Mei Z, Grummer-Strawn L, Parvanta I. Haemoglobin adjustments to define anaemia. *Tropical Medicine and International Health.* 2008. 13(10):1267-71.
  24. Thurnham DI, McCabe GP, Northrop-Clewes C, Nestel P. Effects of subclinical infection on plasma retinol concentrations and assessment of prevalence of vitamin A deficiency: a meta-analysis. *Lancet.* 2003;362:2052-58.
  25. Thurnham DI, McCabe LD, Haldar S, Wieringa FT, Northrop-Clewes C, McCabe GP. Adjusting plasma ferritin concentrations to remove the effects of subclinical inflammation in the assessment of iron deficiency: a meta-analysis. *Am J Clin Nutr* 2010;92:546-55.
  26. UN, Working Group on Nutrition of School-Age Children. Overview of the
  27. Working Group accomplishments. United Nations Standing Committee on Nutrition (UN-SCN). 2003. [06/18/2013]; Available from: [http://www.unscn.org/en/working\\_groups/sac/sessions/30th\\_report.php](http://www.unscn.org/en/working_groups/sac/sessions/30th_report.php)
  28. Valadez JJ. *Assessing child survival programs in developing countries: testing lot quality assurance sampling.* Boston: Harvard University Press; 1991.
-

- 
29. Valadez JJ, Weiss W, Leburg C, Davis R. Assessing Community Health Programs, A Trainer's Guide; Using LQAS for Baseline Surveys and Regular Monitoring. St Albans: Teaching-aids At Low Cost; 2003.
  30. WHO. Global Database on Child Growth and Malnutrition. 2011. World Health Organization: Geneva. Available from: [www.who.int/nutgrowthdb/about/introduction/en/index5.html](http://www.who.int/nutgrowthdb/about/introduction/en/index5.html)
  31. WHO. Global prevalence of vitamin A deficiency in populations at risk 1995-2005. 2009. World Health Organization: Geneva.
  32. WHO. Indicators for assessing infant and young child feeding practices: Conclusions of a consensus meeting held 6-8 November 2007 in Washington D.C., USA. 2008, WHO: Geneva.
  33. WHO. Indicators for assessing infant and young child feeding practices – part I: definition. 2008. World Health Organization: Geneva.
  34. WHO. Indicators for assessing infant and young child feeding practices – part 2: measurement. 2008. World Health Organization: Geneva.
  35. WHO. Indicators for assessing vitamin A deficiency and their application in monitoring and evaluating intervention programmes. 1996. World Health Organization: Geneva.
  36. WHO, UNU, UNICEF. Iron deficiency anaemia assessment, prevention, and control - A guide for programme managers. 2001. World Health Organization United Nations University, UNICEF: Geneva.
  37. WHO. Physical status: The use and interpretation of anthropometry: Report of a WHO expert committee. WHO Technical Report Series No. 854. 1995. World Health Organization: Geneva.
  38. WHO. The optimal duration of exclusive breastfeeding. Report of an Expert Consultation. 2001, World Health Organization: Geneva.
  39. WHO. Multicentre Growth Reference Study Group. WHO Child Growth Standards: Length/height-for-age, weight-for-age, weight-for-length, weight-for-height, and body mass index-for-age: Methods and development. 2006. World Health Organization: Geneva.
  40. WHO. Guideline: Vitamin A supplementation in infants and children 6–59 months of age. 2011. World Health Organization: Geneva.

---

## APPENDICES

- Appendix I: Questionnaires for 2013 National Nutrition Survey
- Appendix II: List of Child Hemoglobin Results: For Village Medical Attendant
- Appendix III: Confidence Intervals, Design Effects, and Intra-Class Correlation Coefficients for Major Indicators, children 6–29 months — Kyrgyzstan National Survey, 2013
- Appendix IV: Questionnaires for 2011 and 2012 Lot Quality Assurance Sampling
- Appendix V: Algorithms for Calculating Indicators of Feeding Practices
- Appendix VI: National Survey of Nutritional Status, Kyrgyzstan 2013: Results for the WHO Global Database on Child Growth and Malnutrition
- Appendix VII: Brief Summary Quality Assurance for Micronutrient Measurements from the 2009 and 2013 Kyrgyzstan Surveys
- Appendix VIII: Questionnaires for 2009 National Nutrition Survey
- Appendix IX: Comparison of Various Characteristics among Children 6–24 months

# APPENDIX I

## QUESTIONNAIRE FOR 2013 NATIONAL NUTRITION SURVEY

English

2013 NATIONAL NUTRITION AND DIET SURVEY			
		May 6, 2013 - DRAFT	
Fill the following information before beginning the interview.			
HH1. Cluster number	<input type="text"/>	HH4. Supervisor's code	<input type="text"/>
HH2. Interviewer code	<input type="text"/>	HH5. Data entry operator code	<input type="text"/>
HH3. Day/Month/Year of interview:		HH6. Oblast code	<input type="text"/>
<input type="text"/>	<input type="text"/>	02 Issyk - kul	05 Batken
d	d	03 Jalalabad	06 Osh Oblast
		04 Naryn	07 talas
			08 Chui
			11 Bishkek city
			21 Osh City
		HH7. Rayon code	<input type="text"/>
HH8. Result of interview	<input type="text"/>	HH9. Result of anthropometry	<input type="text"/>
Completed	1	Completed	1
Refused	2	Not completed	2
Partially completed	3		
Not available to interview	4		
HH10. Result of capillary sample	<input type="text"/>	HH11. Location of data collection	<input type="text"/>
Completed	1	Clinic	1
Not completed	2	Home	2
Partially completed	3	Partial in clinic / partial in home	3
HH12. In what language was the interview conducted?	<input type="text"/>	ID Label - CHILD	
Kyrgyz	1	Affix child label here	
Russian	2		
Uzbek	3		
Other (specify) _____	4		
HH13. Does this caretaker have more than one child in this survey?	<input type="text"/>		
yes	1		
no	0		
HH14. Child's Date of Birth	<input type="text"/>	If birthday is before February 1, 2011 or after January 31, 2013, the child cannot participate. Explain this to the mother and thank her for coming.	
d	d	→	
		If birthday falls between February 1, 2011 and January 31, 2013 => G1	
**For questions G1-G3, obtain information from <i>Green Journal (GJ)</i> in clinic		G3. What is the last date that the child received Gulazyk (according to the GJ)?	
G1. Has child ever received Gulazyk?	<input type="text"/>	<input type="text"/>	
yes	1	d	
no (or not listed in GJ)	0	d	
G2. Has child received Gulazyk within 3 months of today's date?	<input type="text"/>	m	
yes	1	m	
no (or not listed in GJ)	0	y	
If no data is collected on child: Find the following information (NR1-NR5) from the clinic, the medical worker, the VHC volunteer or by going to the child's home. If data is collected on a child: <b>SKIP TO CONSENT FORM</b>			
NR1. What is the gender of the child?	<input type="text"/>	NR4. Does the mother work/study outside of the home?	<input type="text"/>
1=male 2=female		1= yes 0 = no 8 = don't know/can't find out	
NR2. What is the ethnicity of the mother?	<input type="text"/>	NR5. Reason for not coming to the interview?	<input type="text"/>
Kyrgyz	1	family moved from village permanently	1
Russian	2	mother refused	2
Kazakh	3	family (husband, mother-in-law, etc.) refused	3
Uzbek	4	mother sick	4
Tajik	5	child sick	5
Uigher	6	mother had to work	6
Other (specify) _____	7	was not invited by the health clinic	7
Don't know	8	temporary migration	8
NR3. How many brothers/sisters does the child have?	<input type="text"/>	Other (specify) _____	9
(put 88 if don't know/cannot find out)		Don't know/can't find out	10
Note: If no data is collected on child, form concludes here (after completing NR1-NR5). Interviewer should move on to the next child.			

We are from the Ministry of Health and the National Statistics Committee. We are working on a joint project concerned with child diet, nutrition and health. I would like to talk to you about this and record your answers to some questions that I have. This interview will take approximately 20 minutes. After the interview, we will weigh and measure your baby and take a small blood sample the finger of your baby. From this sample we will be able to inform you if your baby has anemia. The only direct benefit to you is the knowledge of your baby's anemia status. The risks are small and consist of the possible discomfort caused by pricking the finger in order to draw the blood sample. The discomfort will only be temporary and will not be very great. All of the information we obtain will remain strictly confidential and nobody will know that the information is yours, however, your hemoglobin results will be shared with your village health clinic. You have the right to choose whether or not you would like to participate, and there are no consequences if you decide you would rather not participate. At any time during the interview, weight and length measurement, or blood sample collection you may decide that you no longer wish to participate, and we will stop the interview. If you agree to participate, I would like you to sign this form. May we begin?

Would you please sign?

If permission is given, ask respondent to sign here and begin the interview.

Agreed to participate?   
 yes 1  
 no 0

(Signature) \_\_\_\_\_  
 Permission for child's participation

(Name) \_\_\_\_\_

Signature of interviewer confirming that the respondent agreed to participate and signed this form:

Signature _____		Date _____									
HH15. What is your name?  (last, first, middle initial)	HH20. What is your relationship to (child's name): mother 1 grandmother 2 <b>1→GU1</b> aunt 3 other _____ 6 (specify)										
HH16. What is the name of the child?  (last, first, middle initial)	HH21. Why is (child's name)'s mother not here today?  working/studying 1 <input type="checkbox"/> sick 2 did not want to attend 3 family did not allow 4 busy at home/with housework 5 other _____ 7 (specify) don't know 8										
HH17. What is the gender of the child <input type="checkbox"/> 1= male 2=female											
HH18. How many brothers/sisters does the child have? <input type="checkbox"/> <input type="checkbox"/> (put 88 if don't know/cannot find out)											
HH19. What is the mother's date of birth? <table border="1" style="width: 100%; text-align: center;"> <tr> <td>d</td><td>d</td><td>m</td><td>m</td><td>y</td><td>y</td><td>y</td><td>y</td> </tr> </table> (put 8888 for year if don't know/remember)	d	d	m	m	y	y	y	y			
d	d	m	m	y	y	y	y				

Gulazyk Module		GU
<p>GU1. Have you ever received a package of Gulazyk like this for (child's name)? (show Gulazyk sachets)</p> <p style="text-align: right;">[ ]</p> <p>1=yes 0 = no 8= don't know <b>0→C3</b></p>	<p>GU6. How often does/did (child's name) consume Gulazyk?</p> <p style="text-align: right;">[ ] <b>5→GU11</b></p> <p>Every day for one month 1</p> <p>1 sachet per day for 15 days, then a break for 15 days 2</p> <p>Every other day 3</p> <p>Other schedule _____ 4 (specify)</p> <p>Never yet received 5</p> <p>Don't know 8</p>	
<p>GU2. Is (child's name) currently taking Gulazyk?</p> <p style="text-align: right;">[ ]</p> <p>1=yes 0 = no 8= don't know <b>1→GU5</b></p>	<p>GU7. How many sachets of Gulazyk has (child's name) consumed within 2 months?</p> <p>88 if don't know/remember [ ] [ ]</p> <p>99 if haven't yet taken for 2 months</p> <p>00 if none</p>	
<p>GU3. Why isn't (child's name) currently taking Gulazyk? <i>Don't read out! Circle each answer mentioned.</i></p> <p>A. Too difficult to remember to give Gulazyk</p> <p>B. Child experienced side effects/diarrhea</p> <p>R. Child experienced side effects/skin/allergy</p> <p>S. Child experienced side effects/other</p> <p>(specify) _____</p> <p>C. Child does not like food when Gulazyk is added</p> <p>D. Gulazyk makes the food taste bad</p> <p>E. Gulazyk changes the color of the food</p> <p>F. Mother/caretaker doesn't like Gulazyk</p> <p>G. Health care worker told me to stop using Gulazyk</p> <p>H. Ran out of Gulazyk</p> <p>I. Children should receive natural vitamins from food</p> <p>L. We have been away from home</p> <p>N. Child has been sick</p> <p>O. Don't know</p> <p>P. Belief that Gulazyk is harmful to the child</p> <p>Q. Heard on the TV or radio that Gulazyk is harmful</p> <p>T. Child is &gt;24 months</p> <p>U. Other _____ (specify)</p>	<p>A</p> <p>B</p> <p>R</p> <p>S</p> <p>C</p> <p>D</p> <p>E</p> <p>F</p> <p>G</p> <p>H</p> <p>I</p> <p>L</p> <p>N</p> <p>O</p> <p>P</p> <p>Q</p> <p>T</p> <p>U</p>	<p>GU8. Have you noticed any changes in the color of the food to which Gulazyk is added?</p> <p style="text-align: right;">[ ] <b>0→GU9</b></p> <p>1=yes 0 = no</p> <p>GU8a. Is this change in color a concern for you or your child?</p> <p style="text-align: right;">[ ]</p> <p>1=yes 0 = no</p> <p>GU9. Have you noticed any changes in the taste of the food to which Gulazyk is added?</p> <p style="text-align: right;">[ ] <b>0→GU10</b></p> <p>1=yes 0 = no</p> <p>GU9a. Is this change in taste a concern for you or your child?</p> <p style="text-align: right;">[ ]</p> <p>1=yes 0 = no</p>
<p>GU4. At what age in months did (child's name) stop consuming Gulazyk?</p> <p style="text-align: right;">[ ] [ ]</p> <p>88 if don't know/remember m m</p> <p>99 if still consuming</p> <p>77 if never started</p>	<p>GU10. Does/Did (child's name) usually consume the entire portion of food into which Gulazyk is/was mixed, or does he/she consume less than the entire portion?</p> <p style="text-align: right;">[ ]</p> <p>Consumes entire portion 1</p> <p>Consumes less than full portion 2</p> <p>Don't know 8</p>	
<p>GU5. How did you obtain Gulazyk for (child's name)?</p> <p>I go to the health clinic 1</p> <p>Medical worker brings Gulazyk to my home 2</p> <p>Both (of above answers) 3</p> <p>Other _____ 8 (specify)</p>	<p>GU11. When you run out of Gulazyk sachets do you plan to continue?</p> <p style="text-align: right;">[ ] <b>1 or 2→E1</b></p> <p>1=yes 0 = no 2= child is &gt;24 months 8= don't know</p> <p>GU12. What are the reasons that you will probably not continue to give Gulazyk? <b>Do NOT read each item, circle all answers mentioned</b></p> <p>A. Too difficult to remember to give Gulazyk</p> <p>B. Child experienced side effects</p> <p>C. Child does not like food when Gulazyk is added</p> <p>D. Family member doesn't want to use</p> <p>E. Health care worker told me to stop using Gulazyk</p> <p>F. Children should receive natural vitamins from food</p> <p>G. When I run out of Gulazyk my child will be &gt;24 months</p> <p>H. Belief that Gulazyk is harmful to children</p> <p>I. Heard on TV or radio that Gulazyk is harmful</p> <p>J. Other (specify) _____</p> <p>K. Don't know</p>	



Effects Module		E																																																			
E1. Have you noticed any positive changes in your child since he/she started taking Gulazyk that you believe are due to Gulazyk? <div style="text-align: right; margin-right: 50px;"> <input style="width: 50px; height: 20px;" type="text"/>              1=yes 0 = no <span style="float: right;">0→E3</span> </div>	E3. Have you recommended Gulazyk to other families? <div style="text-align: right; margin-right: 50px;"> <input style="width: 50px; height: 20px;" type="text"/>              1=yes 0 = no 8= don't know           </div>																																																				
E2. What are the positive changes that you noticed in your child? <b>Do NOT read each item, circle all answers mentioned</b> <table style="width: 100%; border-collapse: collapse;"> <tr><td>More energy</td><td style="text-align: center;">A</td></tr> <tr><td>Better growth</td><td style="text-align: center;">B</td></tr> <tr><td>More curiosity/intelligence</td><td style="text-align: center;">C</td></tr> <tr><td>Improved eyesight</td><td style="text-align: center;">D</td></tr> <tr><td>Gets sick less often</td><td style="text-align: center;">E</td></tr> <tr><td>Increased appetite</td><td style="text-align: center;">F</td></tr> <tr><td>Overall seems better/healthier</td><td style="text-align: center;">G</td></tr> <tr><td>No positive changes noticed</td><td style="text-align: center;">H</td></tr> <tr><td>Other _____</td><td style="text-align: center;">I</td></tr> </table> <p style="text-align: center; margin-top: 5px;">(specify)</p>	More energy	A	Better growth	B	More curiosity/intelligence	C	Improved eyesight	D	Gets sick less often	E	Increased appetite	F	Overall seems better/healthier	G	No positive changes noticed	H	Other _____	I																																			
More energy	A																																																				
Better growth	B																																																				
More curiosity/intelligence	C																																																				
Improved eyesight	D																																																				
Gets sick less often	E																																																				
Increased appetite	F																																																				
Overall seems better/healthier	G																																																				
No positive changes noticed	H																																																				
Other _____	I																																																				
Communications Module		C																																																			
C1. Did you receive this brochure about Gulazyk? <i>(show brochure)</i> <div style="text-align: right; margin-right: 50px;"> <input style="width: 50px; height: 20px;" type="text"/>              1=yes 0 = no 8= don't know           </div>	C4. Have you heard about Gulazyk on the radio? <div style="text-align: right; margin-right: 50px;"> <input style="width: 50px; height: 20px;" type="text"/>              1=yes 0 = no 8= don't know           </div>																																																				
C2. Did you read the brochure? <div style="text-align: right; margin-right: 50px;"> <input style="width: 50px; height: 20px;" type="text"/>              1=yes 0 = no 8= don't know           </div>	C5. Do you have a television? <div style="text-align: right; margin-right: 50px;"> <input style="width: 50px; height: 20px;" type="text"/>              1=yes 0 = no 8= don't know  <b>0: if mother→WM1 if NOT mother→BF1</b> </div>																																																				
C3. Do you have a radio? <div style="text-align: right; margin-right: 50px;"> <input style="width: 50px; height: 20px;" type="text"/>              1=yes 0 = no 8= don't know <span style="float: right;">0→C5</span> </div>	C6. Have you heard about Gulazyk on the television? <div style="text-align: right; margin-right: 50px;"> <input style="width: 50px; height: 20px;" type="text"/>              1=yes 0 = no 8= don't know           </div>																																																				
If respondent is mother→WM1 If respondent is NOT mother: SKIP TO BF1																																																					
Woman's Module		WM																																																			
WM1. What is your ethnicity? <table style="width: 100%; border-collapse: collapse;"> <tr><td>Kyrgyz</td><td style="text-align: center;">1</td><td style="text-align: right;"><input style="width: 50px; height: 20px;" type="text"/></td></tr> <tr><td>Russian</td><td style="text-align: center;">2</td><td></td></tr> <tr><td>Kazakh</td><td style="text-align: center;">3</td><td></td></tr> <tr><td>Uzbek</td><td style="text-align: center;">4</td><td></td></tr> <tr><td>Tajik</td><td style="text-align: center;">5</td><td></td></tr> <tr><td>Uigher</td><td style="text-align: center;">6</td><td></td></tr> <tr><td>Other (specify) _____</td><td style="text-align: center;">7</td><td></td></tr> <tr><td>Don't know</td><td style="text-align: center;">8</td><td></td></tr> </table>	Kyrgyz	1	<input style="width: 50px; height: 20px;" type="text"/>	Russian	2		Kazakh	3		Uzbek	4		Tajik	5		Uigher	6		Other (specify) _____	7		Don't know	8		WM5. Do you currently work or study outside the home (for example, as an employee, business owner, laborer in fields, etc.)? <div style="text-align: right; margin-right: 50px;"> <input style="width: 50px; height: 20px;" type="text"/>              1 = yes 0 = no <span style="float: right;">0→BF1</span> </div>																												
Kyrgyz	1	<input style="width: 50px; height: 20px;" type="text"/>																																																			
Russian	2																																																				
Kazakh	3																																																				
Uzbek	4																																																				
Tajik	5																																																				
Uigher	6																																																				
Other (specify) _____	7																																																				
Don't know	8																																																				
WM2. What is the highest level of school you completed? <table style="width: 100%; border-collapse: collapse;"> <tr><td>Never attended</td><td style="text-align: center;">0</td><td style="text-align: right;"><input style="width: 50px; height: 20px;" type="text"/></td></tr> <tr><td>Primary (1-4 grades)</td><td style="text-align: center;">1</td><td></td></tr> <tr><td>Incomplete secondary (5-9)</td><td style="text-align: center;">2</td><td></td></tr> <tr><td>Complete secondary</td><td style="text-align: center;">3</td><td></td></tr> <tr><td>Technical school</td><td style="text-align: center;">4</td><td></td></tr> <tr><td>Higher</td><td style="text-align: center;">5</td><td></td></tr> <tr><td>Religious curriculum</td><td style="text-align: center;">6</td><td></td></tr> <tr><td>Don't know</td><td style="text-align: center;">8</td><td></td></tr> </table>	Never attended	0	<input style="width: 50px; height: 20px;" type="text"/>	Primary (1-4 grades)	1		Incomplete secondary (5-9)	2		Complete secondary	3		Technical school	4		Higher	5		Religious curriculum	6		Don't know	8		WM6. What type of work or study do you do? <table style="width: 100%; border-collapse: collapse;"> <tr><td>laborer (in the fields)</td><td style="text-align: center;">1</td><td style="text-align: right;"><input style="width: 50px; height: 20px;" type="text"/></td></tr> <tr><td>vender of food, fruit, homemade goods or other</td><td style="text-align: center;">2</td><td></td></tr> <tr><td>employee in a business</td><td style="text-align: center;">3</td><td></td></tr> <tr><td>business owner</td><td style="text-align: center;">4</td><td></td></tr> <tr><td>professional (nurse, doctor, teacher, pharmacist, etc.)</td><td style="text-align: center;">5</td><td></td></tr> <tr><td>student</td><td style="text-align: center;">6</td><td></td></tr> <tr><td>other (specific) _____</td><td style="text-align: center;">7</td><td></td></tr> <tr><td>Don't know</td><td style="text-align: center;">8</td><td></td></tr> </table>		laborer (in the fields)	1	<input style="width: 50px; height: 20px;" type="text"/>	vender of food, fruit, homemade goods or other	2		employee in a business	3		business owner	4		professional (nurse, doctor, teacher, pharmacist, etc.)	5		student	6		other (specific) _____	7		Don't know	8				
Never attended	0	<input style="width: 50px; height: 20px;" type="text"/>																																																			
Primary (1-4 grades)	1																																																				
Incomplete secondary (5-9)	2																																																				
Complete secondary	3																																																				
Technical school	4																																																				
Higher	5																																																				
Religious curriculum	6																																																				
Don't know	8																																																				
laborer (in the fields)	1	<input style="width: 50px; height: 20px;" type="text"/>																																																			
vender of food, fruit, homemade goods or other	2																																																				
employee in a business	3																																																				
business owner	4																																																				
professional (nurse, doctor, teacher, pharmacist, etc.)	5																																																				
student	6																																																				
other (specific) _____	7																																																				
Don't know	8																																																				
WM3. Are you currently married? <div style="text-align: right; margin-right: 50px;"> <input style="width: 50px; height: 20px;" type="text"/>              1= yes 0= no <span style="float: right;">0→WM5</span> </div>	WM7. How many hours a day do you USUALLY work or study outside of the home? (put 88 if don't know) <div style="text-align: right; margin-right: 50px;"> <input style="width: 50px; height: 20px;" type="text"/> </div>																																																				
WM4. What is the highest level of school your spouse completed? <table style="width: 100%; border-collapse: collapse;"> <tr><td>Never attended</td><td style="text-align: center;">0</td><td style="text-align: right;"><input style="width: 50px; height: 20px;" type="text"/></td></tr> <tr><td>Primary (1-4 grades)</td><td style="text-align: center;">1</td><td></td></tr> <tr><td>Incomplete secondary (5-9)</td><td style="text-align: center;">2</td><td></td></tr> <tr><td>Complete secondary</td><td style="text-align: center;">3</td><td></td></tr> <tr><td>Technical school</td><td style="text-align: center;">4</td><td></td></tr> <tr><td>Higher</td><td style="text-align: center;">5</td><td></td></tr> <tr><td>Religious curriculum</td><td style="text-align: center;">6</td><td></td></tr> <tr><td>Don't know</td><td style="text-align: center;">8</td><td></td></tr> </table>	Never attended	0	<input style="width: 50px; height: 20px;" type="text"/>	Primary (1-4 grades)	1		Incomplete secondary (5-9)	2		Complete secondary	3		Technical school	4		Higher	5		Religious curriculum	6		Don't know	8		WM8. Who USUALLY feeds (child's name) while you are outside of the home? <table style="width: 100%; border-collapse: collapse;"> <tr><td>The mother (takes the child with her)</td><td style="text-align: center;">1</td><td style="text-align: right;"><input style="width: 50px; height: 20px;" type="text"/></td></tr> <tr><td>Baby's grandmother</td><td style="text-align: center;">2</td><td></td></tr> <tr><td>Baby's sisters/brothers</td><td style="text-align: center;">3</td><td></td></tr> <tr><td>Baby's father</td><td style="text-align: center;">4</td><td></td></tr> <tr><td>Other family member</td><td style="text-align: center;">5</td><td></td></tr> <tr><td>Baby sitter</td><td style="text-align: center;">6</td><td></td></tr> <tr><td>Day care / children's garden</td><td style="text-align: center;">7</td><td></td></tr> <tr><td>Other (specify) _____</td><td style="text-align: center;">8</td><td></td></tr> <tr><td>Don't know</td><td style="text-align: center;">9</td><td></td></tr> </table>		The mother (takes the child with her)	1	<input style="width: 50px; height: 20px;" type="text"/>	Baby's grandmother	2		Baby's sisters/brothers	3		Baby's father	4		Other family member	5		Baby sitter	6		Day care / children's garden	7		Other (specify) _____	8		Don't know	9	
Never attended	0	<input style="width: 50px; height: 20px;" type="text"/>																																																			
Primary (1-4 grades)	1																																																				
Incomplete secondary (5-9)	2																																																				
Complete secondary	3																																																				
Technical school	4																																																				
Higher	5																																																				
Religious curriculum	6																																																				
Don't know	8																																																				
The mother (takes the child with her)	1	<input style="width: 50px; height: 20px;" type="text"/>																																																			
Baby's grandmother	2																																																				
Baby's sisters/brothers	3																																																				
Baby's father	4																																																				
Other family member	5																																																				
Baby sitter	6																																																				
Day care / children's garden	7																																																				
Other (specify) _____	8																																																				
Don't know	9																																																				

Breastfeeding and Infant Feeding		BF
Now I would like to ask you some questions about the breastfeeding and feeding of (child's name)		
BF1. Was (child's name) ever breastfed? 1=yes 0=no 8=Don't know	<input type="text"/>	0 or 8 → BF3
BF2. Approximately, how long after birth was (child's name) first put to the breast? Immediately (< 1 hour after birth) 0 During first 24 hours 1 Between 24 - 48 hours 2 > 48 hours 3 Don't know/remember 8	<input type="text"/>	
The next few questions are about the first time (child's name) was fed something other than breastmilk.		
BF3. How old was [child's name] in months when (he/she) was first fed animal milk, powdered milk or formula? <i>(if less than 1 month put 00, if NEVER fed milk, powdered milk or formula put 99, if don't know put 88)</i>	<input type="text"/> <input type="text"/> m m <i>(round down to nearest whole month)</i>	
BF4. The next question is about liquids. Please include all liquids such as animal milk, powdered milk, formula, juice, water, sugar or fruit water, tea, or anything else that (child's name) might have been given. How old was (child's name) in months when he/she was first given any liquid, even tea, other than breastmilk? <i>(if less than 1 month put 00, if NEVER fed anything other than breastmilk put 99 if don't know put 88)</i>	<input type="text"/> <input type="text"/> m m <i>(round down to nearest whole month)</i>	
BF5. The next question is about solid or semi-solid foods. Please include all solids such as porridge, rice, cereal, bulymak or anything else that (child's name) might have been given. How old was (child's name) in months when he/she was first fed any solid food? <i>(if less than 1 month put 00, if NEVER fed anything other than breastmilk put 99 if don't know put 88)</i>	<input type="text"/> <input type="text"/> m m <i>(round down to nearest whole month)</i>	
		Now think about everything (child's name) has drunk or eaten since this time yesterday. Don't forget snacks and eating or drinking during the night or things (child's name) ate with someone other than yourself.
		BF6. Since this time yesterday, was (child's name) fed any of the following items? <i>(read each item aloud and record response before continuing)</i> 1=yes 0=no 8= don't know
		a Breastmilk <input type="text"/>
		b Animal milk, yogurt, kefir, cheese, etc. <input type="text"/>
		c infant formula or powdered milk <input type="text"/> <i>(probe: what was the name?)</i>
		Brand name? _____
		d haricot, pea or nuts <input type="text"/>
		e kasha, potatoes, noodles, beet <input type="text"/>
		f meat, fish, poultry, liver/organ meat <input type="text"/>
		g eggs <input type="text"/>
		h carrots, pumpkin, tomatoes <input type="text"/>
		i other fruit or vegetable (spinach, dried apricots, cucumbers, etc.) <input type="text"/>
		j bread or biscuit <input type="text"/>
		k baby cereal/food which was purchased <input type="text"/> Brand name? _____
		BF7. Since this time yesterday, how many times was (child's name) fed: <i>(if more than 7 put 7. If don't know put 8)</i> <i>("fed" means any meal or snack, excluding trivial amounts)</i>
		a any solid, semisolid, or soft food such as porridge, cereal, meat, vegetables, cookies, fruit, etc. <input type="text"/>
		b Breastmilk <input type="text"/>
		c animal milk, powdered milk or formula <input type="text"/>
		d anything from a bottle <input type="text"/>
		BF8. Has (child's name) stopped breastfeeding? 1=yes 0=no <input type="text"/> 0 → AB1
		BF9. At what age in months did (child's name) stop breastfeeding? <i>(put 88 if don't know/can't remember)</i> <input type="text"/> <input type="text"/> m m
Attitude, Behavior Module		AB
We are interested in knowing what mothers think about breastfeeding and feeding of their babies. I would like to ask you what you think about breastfeeding and feeding of your baby. Remember there are no right or wrong answers to any of these questions. We just want to know what you think about these topics.		
AB1. Using this scale, how would you describe the importance of breastfeeding for a baby's health and nutrition? <input type="text"/> <i>(show the scale and note the number that corresponds to the answer)</i>	<input type="text"/>	
AB2. In your opinion, should a baby be breastfed? 1=yes 0=no 8= don't know	<input type="text"/>	0 or 8 → AB4
AB3. In your opinion, how long in months should a baby be breastfed? <i>(note 00 if &lt; 1 m; 88 if don't know)</i>	<input type="text"/> <input type="text"/> m m	
		AB4. In your opinion, at what age in months should a baby start drinking other liquids like tea, water, milk, etc.? <input type="text"/> <input type="text"/> <i>(note 00 if &lt; 1 m; 88 if don't know)</i> m m
		AB5. In your opinion, at what age in months should a baby start eating foods like porridge, cereal, bulymak, etc.? <input type="text"/> <input type="text"/> <i>(note 00 if &lt; 1 m; 88 if don't know)</i> m m
		AB6. Some people think there are advantages to breastfeeding while some people do not. In your opinion, are there advantages to breastfeeding? <input type="text"/> 1=yes 0=no 8= don't know
Dietary Advice Module		DA
When a family has a new baby, many people give advice on breastfeeding and feeding the baby. I want to ask just about the advice you have received, it doesn't matter if it is advice you followed or not. I am just interested in what people have told you and who you have heard it from.		
DA1. Did a doctor, nurse, midwife or feldsher give you advice on breastfeeding? 1=yes 0=no 8= don't know	<input type="text"/>	0 or 8 → VS1
DA2. For how long (in months) did a doctor, nurse, midwife	<input type="text"/> <input type="text"/> m m	00 if < 1 m
		DA3. At what age (in months) did a doctor, nurse, midwife or feldsher advise you to stop breastfeeding? <input type="text"/> <input type="text"/> m m

or feldsher advise you to breastfeed without giving other liquids or solids? <input type="text"/> m <input type="text"/> m <i>(Put 00 if &lt; 1 m; 88 if don't know/remember; 99 if they did not give advice on length or did not specify exact length)</i>	88 if don't know/remember 99 if they did not give advice on length or did not specify exact length)
<b>Vitamins/Supplements Module</b> <span style="float: right;"><b>VS</b></span>	
I am now going to ask some questions about vitamins and supplements your baby might have taken. Some people take these supplements and some don't and that is okay.	
VS1. Has (child's name) ever taken a Vitamin A capsule like this one? <input type="text"/> Show 100,000IU for 6-11 month old <b>0→VS3</b> Show 200,000IU for 12-29 month old <b>8→VS3</b> 1=yes 0 = no 8= don't know	VS6. How long ago (in months) did (child's name) stop taking the iron syrup or drops? <input type="text"/> m <input type="text"/> m <i>(mark 00 if &lt;1 month; mark 88 if don't know/remember)</i>
VS2. How long ago (in months) did (child's name) take the most recent vitamin A capsule? <input type="text"/> m <input type="text"/> m <i>(note 00 if &lt; 1 m; note 88 if don't know/remember)</i>	VS7. For how long (in months) did (child's name) consume, or has (child's name) been consuming, iron syrup or drops? <input type="text"/> m <input type="text"/> m <i>(note 00 if &lt;1 month; note 88 if don't know/remember)</i>
VS3. Have you ever been told by a doctor or nurse that (child's name) had anemia? <input type="text"/> 1=yes 0 = no 8= don't know	VS8. Have you, or someone else, ever given (child's name) any of these other vitamin or mineral supplements? <i>(read the list and mark each answer)</i>
VS4. Has (child's name) ever taken iron syrup or drops like this? <input type="text"/> <i>(show iron syrup and drops)</i> <b>0→VS8</b> 1=yes 0 = no 8= don't know <b>8→VS8</b>	a Vitamin D <input type="text"/> b Fish oil <input type="text"/> c Multi-vitamins <input type="text"/> d Zinc <input type="text"/> e Other _____ <input type="text"/> <i>(specify)</i>
VS5. Is (child's name) currently taking iron syrup or drops? <input type="text"/> 1=yes 0 = no 8= don't know <b>1→VS7</b>	1=yes 0 = no 8= don't know
<b>Household Characteristics Module</b> <span style="float: right;"><b>HC</b></span>	
I would now like to ask you a few questions about your home and those who live in it.	
HC1. Does your family currently receive the universal monthly benefit? <input type="text"/> 1=yes 0=no 8= don't know	

<b>Anthropometry</b> <span style="float: right;"><b>AN</b></span>	
Now I am going to measure the length and weight of your baby.	
AN1. Were anthropometrics taken from the child? <input type="text"/> 1 = yes 0= no <b>1→AN3</b>	AN4. Child's length/height (cm) <input type="text"/> <input type="text"/> <input type="text"/> . <input type="text"/>
AN2. Why not? 1 = refused (cried, kicked, etc.) 3=not present 2 = mother/guardian refused 4 other (specify)	AN5. Was length/height of child measured lying down or standing up? <input type="text"/> 1 = Lying down (children less than 24 months of age) 2 = Standing up (children 24-29 months of age)
AN3. Child's weight (kg) <input type="text"/> <input type="text"/> . <input type="text"/>	

<b>Blood Sample Module</b> <span style="float: right;"><b>BS</b></span>	
The last thing we will do today is take a small sample of blood from the finger of your baby. This might cause a little discomfort from the stick but we will be able to tell you if your baby has anemia.	
BS1. Was a capillary sample obtained from the child? <input type="text"/> 1=yes 0 = no <b>1→BS3</b>	BS4. Hemoglobin concentration from Hemocue <input type="text"/> <input type="text"/> . <input type="text"/> g/dL <i>(put 88.8 if not measured/don't know)</i>
BS2. Why not? 1 = refused (cried, kicked, etc.) 4 = technical difficulties 2 = mother/guardian refused 5 = other (specify) 3 = not present <b>Go to →BS5</b>	BS5. ID Label - <b>CHILD</b> Affix child label for blood here
BS3. Approximately how many microliters of blood were collected in the microtainer? <input type="text"/> <input type="text"/> <input type="text"/>	

Signature of site supervisor confirming child was referred to primary health care provider for treatment if hemoglobin <8.0 g/dL

\_\_\_\_\_  
 Signature Date

Don't forget to provide the mother with the hemoglobin measurement results for her child.  
 Ask mother to sign here to confirm receipt of hemoglobin measurement results and referral (if appropriate).

\_\_\_\_\_  
 Signature Date

Signature of site supervisor confirming they have checked the questionnaire and it is complete:

\_\_\_\_\_  
 Signature Date

Interviewer Comments:  
 \_\_\_\_\_  
 \_\_\_\_\_

2013 НАЦИОНАЛЬНОЕ ИССЛЕДОВАНИЕ ПО ПИТАНИЮ															
6 мая, 2013 - Предварительный вариант															
<i>Заполните следующую информацию перед началом интервью.</i>															
НН1. Номер кластера <input style="width: 50px;" type="text"/>		НН4. Код супервайзера <input style="width: 50px;" type="text"/>													
НН2. Код интервью <input style="width: 50px;" type="text"/>		НН5. Код оператора введения данных <input style="width: 50px;" type="text"/>													
НН3. День/Месяц/Год проведения интервью: <table style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 15%; border: 1px solid black; text-align: center;">д</td> <td style="width: 15%; border: 1px solid black; text-align: center;">д</td> <td style="width: 15%; border: 1px solid black; text-align: center;">м</td> <td style="width: 15%; border: 1px solid black; text-align: center;">м</td> <td style="width: 15%; border: 1px solid black; text-align: center;">г</td> <td style="width: 15%; border: 1px solid black; text-align: center;">г</td> </tr> <tr> <td style="text-align: center;">1</td> <td style="text-align: center;">3</td> <td colspan="4"></td> </tr> </table>		д	д	м	м	г	г	1	3					НН6. Код области 02 Иссык-Куль    05 Баткен    08 Чуй 03 Жалалабад    06 Ошская область    21 Город Ош 04 Нарын    07 Талас	
д	д	м	м	г	г										
1	3														
НН7. Код района <input style="width: 100px;" type="text"/>															
НН8. Результат интервью Проведено <input type="checkbox"/> 1 Отказ <input type="checkbox"/> 2 Частично проведено <input type="checkbox"/> 3 Отсутствие интервьюируемого <input type="checkbox"/> 4		НН9. Результат антропометрии Проведено <input type="checkbox"/> 1 Не проведено <input type="checkbox"/> 2													
НН10. Результат анализа крови Проведено <input type="checkbox"/> 1 Не проведено <input type="checkbox"/> 2 Частично проведено <input type="checkbox"/> 3		НН11. Место сбора данных Медучреждение <input type="checkbox"/> 1 На дому <input type="checkbox"/> 2 Частично в медучр., частично на дому <input type="checkbox"/> 3													
НН12. На каком языке проводилось интервью? Кыргызском <input type="checkbox"/> 1 Русском <input type="checkbox"/> 2 Узбекском <input type="checkbox"/> 3 Другом (укажите) <input style="width: 50px;" type="text"/> 4		Бирка с идентификационным номером ребенка <i>Прикрепить номер ребенка здесь</i>													
НН13. Имеет ли данное лицо, обеспечивающее уход за ребенком и участвующее в исследовании, более одного ребенка? Да <input type="checkbox"/> 1 Нет <input type="checkbox"/> 0															
НН14. Дата рождения ребенка <table style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 15%; border: 1px solid black; text-align: center;">д</td> <td style="width: 15%; border: 1px solid black; text-align: center;">д</td> <td style="width: 15%; border: 1px solid black; text-align: center;">м</td> <td style="width: 15%; border: 1px solid black; text-align: center;">м</td> <td style="width: 15%; border: 1px solid black; text-align: center;">г</td> <td style="width: 15%; border: 1px solid black; text-align: center;">г</td> </tr> </table> → если ребенок родился <b>до 1 февраля 2011 г. или после 31 января 2013 г.</b> , ребенок не может принимать участие в исследовании. Объясните это матери и поблагодарите ее за визит. если день рождения приходится между <b>1 февраля 2011г. и 31 января 2013г.</b> => <b>G1</b>				д	д	м	м	г	г						
д	д	м	м	г	г										
**Для вопросов G1-G3 возьмите информацию из Зеленого Журнала (ЗЖ) в поликлинике Г1. Получал ли когда-либо ребенок Гулязык? да <input type="checkbox"/> 1 нет (или не отмечено в ЗЖ) <input type="checkbox"/> 0		G3. Когда последний раз ребенок получал Гулязык (согласно ЗЖ)? <table style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 15%; border: 1px solid black; text-align: center;">д</td> <td style="width: 15%; border: 1px solid black; text-align: center;">д</td> <td style="width: 15%; border: 1px solid black; text-align: center;">м</td> <td style="width: 15%; border: 1px solid black; text-align: center;">м</td> <td style="width: 15%; border: 1px solid black; text-align: center;">г</td> <td style="width: 15%; border: 1px solid black; text-align: center;">г</td> </tr> </table>		д	д	м	м	г	г						
д	д	м	м	г	г										
Г2. Получал ли ребенок Гулязык в течение 3 месяцев начиная с этого дня? да <input type="checkbox"/> 1 нет (или не отмечено в ЗЖ) <input type="checkbox"/> 0															
Если на ребенка отсутствует информация: Получите следующую информацию во время посещения ребенка на дому. Если информация на ребенка имеется: <b>ПЕРЕХОДИТЕ К ФОРМЕ НА ДАЧУ СОГЛАСИЯ</b>															
NR1. Каков пол ребенка? 1=муж    2=жен		NR4. Работает/учится ли мать за пределами места проживания? 1= да    0 = нет    8 = неизвестно/невозможно узнать													
NR2. Какова национальность матери? Кыргызка <input type="checkbox"/> 1 Русская <input type="checkbox"/> 2 Казашка <input type="checkbox"/> 3 Узбечка <input type="checkbox"/> 4 Таджичка <input type="checkbox"/> 5 Уйгурка <input type="checkbox"/> 6 Другое (укажите) <input style="width: 50px;" type="text"/> 7 Не знаю <input type="checkbox"/> 8		NR5. Причина неучастия в интервью? семья навсегда уехала из села <input type="checkbox"/> 1 отказ матери <input type="checkbox"/> 2 отказ семьи (мужа, свекрови и т.д.) <input type="checkbox"/> 3 болезнь матери <input type="checkbox"/> 4 болезнь ребенка <input type="checkbox"/> 5 занятость матери на работе <input type="checkbox"/> 6 отсутствие приглашения из поликлиники <input type="checkbox"/> 7 временная миграция <input type="checkbox"/> 8 другое (укажите) <input style="width: 50px;" type="text"/> 9 неизвестно/невозможно узнать <input type="checkbox"/> 10													
NR3. Сколько братьев/сестер у ребенка? <input style="width: 50px;" type="text"/> <i>(поставьте 88 если неизвестно/невозможно узнать)</i>															
Прим: Если на ребенка нет информации, форма заканчивается здесь (после заполнения NR1-NR5). Переходите к опросу следующего ребенка															
Мы из Министерства здравоохранения и Натстаткома. Мы работаем в рамках проекта по питанию и здоровью детей. Мне бы хотелось поговорить с вами об этом и записать ваши ответы на вопросы, которые я вам задам. Интервью займет примерно 20 мин. После интервью мы взвесим и измерим вашего ребенка, а также возьмем у него небольшой анализ крови, из которого мы узнаем есть ли у вашего ребенка анемия, что будет очень полезно для вас. При этом, риски очень небольшие, состоящие только из непродолжительного и несильного дискомфорта при протыкании пальца ребенка при заборе крови. Все ваши ответы будут строго конфиденциальны, никто не узнает, что это ваши данные, только результаты измерения уровня гемоглобина будут предоставлены в вашу поликлинику. Вы можете согласиться или отказаться от участия в интервью, никаких негативных последствий в случае вашего отказа не будет. В любой момент интервью, измерения веса и роста или анализа крови мы можете отказаться от дальнейшего участия. Если вы согласны участвовать, пожалуйста подпишите эту форму. Мы можем начинать?															
<input style="width: 100px; height: 20px;" type="text"/>															
Пожалуйста подпишите?															
Если разрешение получено, попросите респондента подписать здесь и начинайте интервью.		Вы согласны участвовать? <input style="width: 50px;" type="text"/>													
		да <input type="checkbox"/> 1 нет <input type="checkbox"/> 0													
(Подпись) _____ Разрешение на участие ребенка		(Имя) _____													
Подпись интервьюера, подтверждающая, что респондент дал согласие на участие и подписал данную форму:															
<input style="width: 100%; height: 20px;" type="text"/>		<input style="width: 100%; height: 20px;" type="text"/>													
Подпись		Дата													

НН15. Ваше имя? (фамилия, имя, отчество) НН16. Имя ребенка? (фамилия, имя, отчество) НН17. Пол ребенка 1= мужской 2= женский НН18. Сколько у ребенка братьев/сестер? (поставьте 88 если неизвестно/невозможно узнать) НН19. Дата рождения матери? Д Д М М Г Г Г Г (поставьте 88 если неизвестно/невозможно узнать)	НН20. Кем вы приходите ребенку (имя ребенка): мать 1 <b>1→GU1</b> бабушка 2 тетя 3 другое _____ 6 (укажите) НН21. Почему сегодня мать ребенка (имя ребенка) не принимает участие в интервью? работает/учится 1 больна 2 не захотела участвовать 3 запрет семьи 4 занята домашней работой 5 другое _____ 7 (укажите) не знаю 8
<b>Модуль Гулязык</b>	
GU1. Получали ли вы такой набор Гулязык для вашего ребенка (имя ребенка)? (покажите пакетик Гулязык) 1=да 0 = нет 8= не знаю <b>0→C3</b>	GU6. Как часто (имя ребенка) употребляет Гулязык? Каждый день в течение одного месяца 1 1 пакетик в день в течение 15 дней, после перерыва в 15 дней 2 Через день 3 Другое расписание _____ 4 (укажите) Никогда не употреблял 5 Не знаю 8
GU2. Употребляет ли (имя ребенка) Гулязык в настоящее время? 1=да 0 = нет 8= не знаю <b>1→GU5</b>	GU7. Сколько пакетиков Гулязыка ребенок (имя ребенка) употребил в течение 2 месяцев? 88 если не знает/не помнит 00 если ни одного
GU3. Почему (имя ребенка) не употребляет Гулязык в настоящее время? <i>Не читать вслух! Обвести кружком ответ.</i> А. Тяжело помнить о том, что нужно дать Гулязык В. У ребенка побочные явления/диарея R. У ребенка побочные явления/кожные/аллергия S. У ребенка побочные явления/другое (укажите) _____ C. Ребенку не нравится еда с Гулязыком D. Гулязык портит вкус еды E. Гулязык изменяет цвет еды F. Матери/лицу, ухаживающему за ребенком не нравится Гулязык G. Медработник сказал не употреблять Гулязык H. Кончился Гулязык I. Дети должны получать натуральные витамины из продуктов L. Нас не было какое-то время дома N. Болезнь ребенка O. Не знаю P. Считает Гулязык вредным для здоровья ребенка Q. Слышала по ТВ или радио о вреде Гулязыка T. Ребенку больше 24 месяцев U. Другое _____ (укажите)	GU8. Заметили ли вы изменения цвета еды, в которую добавлялся Гулязык? 1=да 0 = нет <b>0→GU9</b>
GU4. Во сколько месяцев ребенок (имя ребенка) прекратил употреблять Гулязык? 88 если не знает/не помнит 77 никогда не употреблял	GU9. Заметили ли вы изменения вкуса еды, в которую добавлялся Гулязык? 1=да 0 = нет <b>0→GU10</b>
GU5. Где вы получаете Гулязык для (имя ребенка)? Хожу в мед. учреждение 1 Медработник приносит Гулязык на дом 2 Оба варианта 3 Другое _____ 8 (укажите)	GU10. Съедает ли (имя ребенка) обычно всю порцию еды, в которую добавлен Гулязык или он употребляет меньшую порцию еды? Съедает всю порцию 1 Съедает меньше 2 Не знаю 8
GU11. Когда у вас кончился Гулязык, вы планируете продолжить его употребление? 1=да 0 = нет 2= ребенку больше 24 мес 8= не знаю <b>1 или 2→E1</b>	GU12. Каковы причины возможного отказа от Гулязык? <i>Не читать вслух, обвести кружком ответ</i> А. Тяжело помнить о том, что нужно дать Гулязык B. У ребенка побочные явления C. Ребенку не нравится еда с Гулязыком D. Отказ члена семьи использовать Гулязык E. Медработник сказал не употреблять Гулязык F. Дети должны получать натуральные витамины из продуктов H. Считает Гулязык вредным для здоровья ребенка I. Слышала по ТВ или радио о вреде Гулязыка J. Другое (укажите) _____ K. Не знаю
<b>Модуль Влияние Гулязыка</b>	
E1. Заметили ли вы улучшения у вашего ребенка благодаря Гулязыку с тех пор, как он начал его употреблять? 1=да 0 = нет <b>0→E3</b>	E3. Рекомендовали ли вы Гулязык другим семьям? 1=да 0 = нет 8= не знаю
E2. Какие улучшения вы заметили у вашего ребенка? <i>Не читать вслух, обвести кружком ответ</i> A. Больше энергии B. Лучший рост C. Более любознательный/сообразительный D. Улучшилось зрение E. Меньше болеет F. Улучшение аппетита G. Вообще выглядит лучше/здоровее I. Другое _____ (укажите)	

Модуль Влияние Гулязыка		E																																																				
E1. Заметили ли вы улучшения у вашего ребенка благодаря Гулязыку с тех пор, как он начал его употреблять? 1=да 0=нет <input type="text"/> <b>0→E3</b>		E3. Рекомендовали ли вы Гулязык другим семьям? 1=да 0=нет 8= не знаю <input type="text"/>																																																				
E2. Какие улучшения вы заметили у вашего ребенка? <b>Не читать вслух, обвести кружком ответ</b> <table border="1"> <tr><td>A</td><td>Больше энергии</td></tr> <tr><td>B</td><td>Лучший рост</td></tr> <tr><td>C</td><td>Более любознательный/сообразительный</td></tr> <tr><td>D</td><td>Улучшилось зрение</td></tr> <tr><td>E</td><td>Меньше болеет</td></tr> <tr><td>F</td><td>Улучшение аппетита</td></tr> <tr><td>G</td><td>Вообщем выглядит лучше/здоровее</td></tr> <tr><td>I</td><td>Другое _____ (укажите)</td></tr> </table>		A	Больше энергии	B	Лучший рост	C	Более любознательный/сообразительный	D	Улучшилось зрение	E	Меньше болеет	F	Улучшение аппетита	G	Вообщем выглядит лучше/здоровее	I	Другое _____ (укажите)																																					
A	Больше энергии																																																					
B	Лучший рост																																																					
C	Более любознательный/сообразительный																																																					
D	Улучшилось зрение																																																					
E	Меньше болеет																																																					
F	Улучшение аппетита																																																					
G	Вообщем выглядит лучше/здоровее																																																					
I	Другое _____ (укажите)																																																					
Модуль Информация		C																																																				
C1. Получали ли вы эту брошюру про Гулязык? (покажите брошюру) <input type="text"/> 1=да 0=нет 8= не знаю		C4. Слышали ли вы о Гулязыке по радио? 1=да 0=нет 8= не знаю <input type="text"/>																																																				
C2. Читали ли вы эту брошюру? 1=да 0=нет 8= не знаю <input type="text"/>		C5. Есть ли у вас телевизор? 1=да 0=нет 8= не знаю <input type="text"/> 0:если мать→WM1 если НЕ мать→BF1																																																				
C3. Есть ли у вас радио? 1=да 0=нет 8= не знаю <input type="text"/> <b>0→C5</b>		C6. Слышали ли вы о Гулязыке по телевидению? 1=да 0=нет 8= не знаю <input type="text"/>																																																				
Если респондент мать→WM1 Если респондент НЕ мать: перейти на BF1																																																						
Модуль Женщина		WM																																																				
WM1. Ваша национальность? <table border="1"> <tr><td>Кыргызка</td><td>1</td><td><input type="text"/></td></tr> <tr><td>Русская</td><td>2</td><td></td></tr> <tr><td>Казашка</td><td>3</td><td></td></tr> <tr><td>Узбечка</td><td>4</td><td></td></tr> <tr><td>Таджичка</td><td>5</td><td></td></tr> <tr><td>Уйгурка</td><td>6</td><td></td></tr> <tr><td>Другое (укажите) _____</td><td>7</td><td></td></tr> <tr><td>Не знаю</td><td>8</td><td></td></tr> </table>		Кыргызка	1	<input type="text"/>	Русская	2		Казашка	3		Узбечка	4		Таджичка	5		Уйгурка	6		Другое (укажите) _____	7		Не знаю	8		WM5. Работаете ли вы или учитесь за пределами места проживания? 1= да 0=нет <input type="text"/> <b>0→BF1</b>																												
Кыргызка	1	<input type="text"/>																																																				
Русская	2																																																					
Казашка	3																																																					
Узбечка	4																																																					
Таджичка	5																																																					
Уйгурка	6																																																					
Другое (укажите) _____	7																																																					
Не знаю	8																																																					
WM2. Какой наивысший уровень вашего образования? <table border="1"> <tr><td>Никогда не училась</td><td>0</td><td><input type="text"/></td></tr> <tr><td>Начальная школа (1-4 классы)</td><td>1</td><td></td></tr> <tr><td>Незаконченное среднее (5-9)</td><td>2</td><td></td></tr> <tr><td>Законченное среднее (11 классов)</td><td>3</td><td></td></tr> <tr><td>Училище</td><td>4</td><td></td></tr> <tr><td>ВУЗ</td><td>5</td><td></td></tr> <tr><td>Религиозное образование</td><td>6</td><td></td></tr> <tr><td>Не знаю</td><td>8</td><td></td></tr> </table>		Никогда не училась	0	<input type="text"/>	Начальная школа (1-4 классы)	1		Незаконченное среднее (5-9)	2		Законченное среднее (11 классов)	3		Училище	4		ВУЗ	5		Религиозное образование	6		Не знаю	8		WM6. Кем вы работаете( или учитесь)? <table border="1"> <tr><td>рабочий</td><td>1</td><td><input type="text"/></td></tr> <tr><td>продавец продуктов, домашних изделий и т.д.</td><td>2</td><td></td></tr> <tr><td>наемный служащий в частном бизнесе</td><td>3</td><td></td></tr> <tr><td>владелец бизнеса</td><td>4</td><td></td></tr> <tr><td>специалист (медсестра, доктор, учитель, фармацевт и т.д.)</td><td>5</td><td></td></tr> <tr><td>студент</td><td>6</td><td></td></tr> <tr><td>другое (укажите) _____</td><td>7</td><td></td></tr> <tr><td>не знаю</td><td>8</td><td></td></tr> </table>		рабочий	1	<input type="text"/>	продавец продуктов, домашних изделий и т.д.	2		наемный служащий в частном бизнесе	3		владелец бизнеса	4		специалист (медсестра, доктор, учитель, фармацевт и т.д.)	5		студент	6		другое (укажите) _____	7		не знаю	8				
Никогда не училась	0	<input type="text"/>																																																				
Начальная школа (1-4 классы)	1																																																					
Незаконченное среднее (5-9)	2																																																					
Законченное среднее (11 классов)	3																																																					
Училище	4																																																					
ВУЗ	5																																																					
Религиозное образование	6																																																					
Не знаю	8																																																					
рабочий	1	<input type="text"/>																																																				
продавец продуктов, домашних изделий и т.д.	2																																																					
наемный служащий в частном бизнесе	3																																																					
владелец бизнеса	4																																																					
специалист (медсестра, доктор, учитель, фармацевт и т.д.)	5																																																					
студент	6																																																					
другое (укажите) _____	7																																																					
не знаю	8																																																					
WM3. Вы замужем? 1= да 0=нет <input type="text"/> <b>0→WM5</b>		WM7. Сколько часов в день вы заняты на работе или учебе? (поставьте 88 если не знает) <input type="text"/>																																																				
WM4. Какой наивысший уровень образования вашего супруга? <table border="1"> <tr><td>Никогда не учился</td><td>0</td><td><input type="text"/></td></tr> <tr><td>Начальная школа (1-4 классы)</td><td>1</td><td></td></tr> <tr><td>Незаконченное среднее (5-9)</td><td>2</td><td></td></tr> <tr><td>Законченное среднее (11 классов)</td><td>3</td><td></td></tr> <tr><td>Училище</td><td>4</td><td></td></tr> <tr><td>ВУЗ</td><td>5</td><td></td></tr> <tr><td>Религиозное образование</td><td>6</td><td></td></tr> <tr><td>Не знаю</td><td>8</td><td></td></tr> </table>		Никогда не учился	0	<input type="text"/>	Начальная школа (1-4 классы)	1		Незаконченное среднее (5-9)	2		Законченное среднее (11 классов)	3		Училище	4		ВУЗ	5		Религиозное образование	6		Не знаю	8		WM8. Кто обычно кормит (имя ребенка) когда вас нет дома? <table border="1"> <tr><td>Мать (берет ребенка с собой)</td><td>1</td><td><input type="text"/></td></tr> <tr><td>Бабушка</td><td>2</td><td></td></tr> <tr><td>Сестры/братья ребенка</td><td>3</td><td></td></tr> <tr><td>Отец ребенка</td><td>4</td><td></td></tr> <tr><td>Другой член семьи</td><td>5</td><td></td></tr> <tr><td>Наня</td><td>6</td><td></td></tr> <tr><td>В дневном центре/детском саду</td><td>7</td><td></td></tr> <tr><td>Другое (укажите) _____</td><td>8</td><td></td></tr> <tr><td>Не знаю</td><td>9</td><td></td></tr> </table>		Мать (берет ребенка с собой)	1	<input type="text"/>	Бабушка	2		Сестры/братья ребенка	3		Отец ребенка	4		Другой член семьи	5		Наня	6		В дневном центре/детском саду	7		Другое (укажите) _____	8		Не знаю	9	
Никогда не учился	0	<input type="text"/>																																																				
Начальная школа (1-4 классы)	1																																																					
Незаконченное среднее (5-9)	2																																																					
Законченное среднее (11 классов)	3																																																					
Училище	4																																																					
ВУЗ	5																																																					
Религиозное образование	6																																																					
Не знаю	8																																																					
Мать (берет ребенка с собой)	1	<input type="text"/>																																																				
Бабушка	2																																																					
Сестры/братья ребенка	3																																																					
Отец ребенка	4																																																					
Другой член семьи	5																																																					
Наня	6																																																					
В дневном центре/детском саду	7																																																					
Другое (укажите) _____	8																																																					
Не знаю	9																																																					

Модуль Грудное вскармливание и питание детей		BF
Теперь я хочу задать несколько вопросов о грудном вскармливании и питании (имя ребенка)		
BF1. Находился когда-либо (имя ребенка) на грудном вскармливании? 1=да 0=нет 8=Не знаю <b>0 или 8→BF3</b>	Теперь скажите что ел или пил (имя ребенка) со вчерашнего дня Не забудьте закуски, еду или питье в ночное время, которые могли давать ребенку другие члены семьи.	
BF2. Как скоро (имя ребенка) после рождения впервые был приложен к груди? Немедленно (через 1 час после рождения) 0 В течение первых 24 часов 1 Между 24 - 48 часами 2 После 48 часов 3 Не знаю/не помню 8	BF6. Ел ли (имя ребенка) со вчерашнего дня что-то из этих продуктов? <i>(зачитайте вслух и запишите каждый ответ отдельно)</i> 1=да 0=нет 8= не знаю a грудное молоко a b животное молоко, йогурт, кефир, сыр и т.д. b c детскую смесь или порошковое молоко c <i>(узнайте: название?)</i> Торговая марка? _____ d фасоль, горох или орехи d e каша, картофель, лапша, свекла e f мясо, рыба, птица, печень/внутренности f g яйца g h морковь, тыква, помидоры h i другие фрукты и овощи (шпинат, урюк, огурцы и т.д.) i j хлеб или печенье j k детские хлопья/питание которое было куплено k Торговая марка? _____	
Следующие вопросы о том, когда (имя ребенка) впервые покормили не грудным молоком.		
BF3. Сколько месяцев было (имя ребенка), когда он впервые получил животное молоко, сухое молоко или молочную смесь? ____ м ____ м <i>(укажите полных месяцев)</i> <i>(поставьте 00, если менее 1 мес.; 99-еще не получал; 88- не знаю)</i>	BF7. Со вчерашнего дня сколько раз кормили (имя ребенка) <i>(если больше 7 поставьте 7. Если не знаю поставьте 8)</i> ("кормили" означает любую пищу или закуску, кроме незначительной пищи) _____ <i>(раз)</i> a твердая, полутвердая или мягкая пища типа каши, крупы, мяса, овощей, печенья, фруктов и т.д. a b грудное молоко b c животное молоко, сухое молоко или смеси c d что-то из бутылочки d	
BF4. Следующий вопрос про жидкости. Пож-та обозначьте все жидкости такие, как животное молоко, сухое молоко, смеси, сок, вода, сладкая или фруктовая вода, чай или другие, которые (имя ребенка) возможно получает. Во сколько месяцев (имя ребенка) начал получать любую жидкость, даже чай, кроме грудного молока? ____ м ____ м <i>(укажите полных месяцев)</i> <i>(поставьте 00, если менее 1 мес.; 99-еще не получал; 88- не знаю)</i>	BF8. (имя ребенка) уже перестали кормить грудью? 1=да 0=нет <b>0→AB1</b>	
BF5. Следующий вопрос о твердой или полутвердой пище. Пож-та обозначьте всю пищу такую, как каша, рис, крупа, булочки или что-то еще, что получает (имя ребенка). Во сколько месяцев (имя ребенка) начал получать любую твердую пищу? ____ м ____ м <i>(укажите полных месяцев)</i> <i>(поставьте 00, если менее 1 мес.; 99-еще не получал; 88= не знаю)</i>	BF9. Во сколько месяцев (имя ребенка) прекратили кормить грудью? <i>(поставьте 88 если не знаю/не помню)</i> ____ м ____ м	
Модуль - Отношение, Поведение		AB
Нам интересно что думают матери о грудном вскармливании и питании их детей. Мне хотелось бы спросить вас, что вы думаете о грудном вскармливании и питании вашего ребенка. Помните нет правильных или неправильных ответов на эти вопросы. Мы просто хотим знать ваше мнение по этим вопросам.		
AB1. Используя эту шкалу, как бы вы оценили важность грудного вскармливания для здоровья и питания ребенка? <i>(покажите шкалу и запишите номер соответствующий ответу)</i> _____ 1=да 0=нет 8= не знаю <b>0 или 8→AB4</b>	AB4. По вашему, во сколько месяцев ребенку нужно начинать давать жидкости типа чая, воды, молока и т.д.? <i>(00 если &lt; 1мес, 88 - не знаю)</i> ____ м ____ м	
AB2. По вашему мнению, ребенка нужно кормить грудью? 1=да 0=нет 8= не знаю <b>0 или 8→AB4</b>	AB5. По вашему, во сколько месяцев ребенка нужно начинать кормить пищей в виде каши, крупы и булочки и т.д.? <i>(00 если &lt; 1 мес.; 88 - не знаю)</i> ____ м ____ м	
AB3. По вашему мнению, как долго ребенок должен находиться на грудном вскармливании? <i>(поставьте 00 если &lt; 1мес; 88 если не знаю)</i> ____ м ____ м	AB6. Некоторые считают, что грудное вскармливание полезно, некоторые - что нет, по вашему мнению грудное вскармливание полезно? 1=да 0=нет 8= не знаю	
Модуль Консультирование по пищевому рациону		DA
Когда в семье рождается ребенок, многие дают советы относительно грудного вскармливания и питания ребенка. Я хочу спросить про это, независимо следовали ли вы им или нет. Мне интересно что именно вам советовали и от кого вы получали эти советы.		
DA1. Как давно(месяцев) доктор, медсестра, акушер или фельдшер консультировал по поводу грудного вскармливания? 1=да 0=нет 8= не знаю <b>0 или 8→VS1</b>	DA3. Как давно(месяцев) доктор, медсестра, акушер или фельдшер советовал прекратить грудное вскармливание?  <i>00 если &lt; 1 мес. 88 не знаю/не помню 99 если они не консультировали по поводу роста или не измеряли точный рост ребенка</i> ____ м ____ м	
DA2. Как давно(месяцев) доктор, медсестра, акушер или фельдшер советовал грудное вскармливание без использования других жидкостей или твердой пищи? <i>(поставьте 00 если &lt; 1 мес; 88 - не знаю/не помню; 99 если не консультировали по поводу роста или не измеряли точный рост ребенка)</i> ____ м ____ м		

Модуль Витамины/Добавки		VS								
Я задам несколько вопросов про витамины и добавки, которые вы возможно даете вашему ребенку. Некоторые употребляют такие добавки, некоторые - нет, это нормально.										
VS1. Получал ли (имя ребенка) когда-либо такой Витамин А в капсулах? Покажите 100,000IU для 6-11-месячных детей Покажите 200,000IU для 12-29-месячных детей 1=да 0=нет 8=не знаю <b>0 или 8→VS3</b>	VS6. Как давно (месяцев) (имя ребенка) получал такой железосодержащий сироп или капли? (поставьте 00 если <1 мес; 88-если не знаю/не помню) <table border="1" style="float: right;"> <tr><td> </td><td> </td></tr> <tr><td>М</td><td>М</td></tr> </table>			М	М					
М	М									
VS2. Как давно (месяцев) (имя ребенка) получал Витамин А в капсулах? (поставьте 00 если < 1 мес; 88 - не знаю/не помню) <table border="1" style="float: right;"> <tr><td> </td><td> </td></tr> <tr><td>м</td><td>м</td></tr> </table>			м	м	VS7. Как долго (месяцев) (имя ребенка) употреблял или употребляет железосодержащий сироп или капли? (поставьте 00 если <1 мес; 88 -если не знаю/не помню) <table border="1" style="float: right;"> <tr><td> </td><td> </td></tr> <tr><td>М</td><td>М</td></tr> </table>			М	М	
м	м									
М	М									
VS3. Говорил ли вам доктор или медсестра, что у (имя ребенка) анемия? 1=нет 0=нет 8=не знаю	VS8. Давали ли вы или кто-то еще (имя ребенка) какие-либо из этих витаминов и минеральных добавок? (прочитайте вслух список и отметьте каждый ответ) а Витамин Д <table border="1" style="float: right;"><tr><td> </td></tr></table> б Рыбий жир <table border="1" style="float: right;"><tr><td> </td></tr></table> с Мультивитамины <table border="1" style="float: right;"><tr><td> </td></tr></table> д Цинк <table border="1" style="float: right;"><tr><td> </td></tr></table> е Другое _____ <table border="1" style="float: right;"><tr><td> </td></tr></table> (укажите) <table border="1" style="float: right; margin-top: 10px;"> <tr><td>1=да</td><td>0=нет</td><td>8=не знаю</td></tr> </table>							1=да	0=нет	8=не знаю
1=да	0=нет	8=не знаю								
VS4. Получал ли (имя ребенка) такой железосодержащий сироп или капли? (покажите сироп и капли) 1=да 0=нет 8=не знаю <b>0 или 8→VS8</b>										
VS5. Получает ли (имя ребенка) в настоящее время железосодержащий сироп или капли? 1=да 0=нет 8=не знаю <b>1→VS7</b>										
Модуль Характеристики домохозяйства		HC								
Я задам еще один вопрос о вашем доме и ваших домочадцах.										
HC1. Получает ли ваша семья универсальное ежемесячное пособие? 1=да 0=нет 8=не знаю										
Антропометрия		AN								
Сейчас я собираюсь измерить рост и вес вашего ребенка.										
AN1. Были ли сделаны антропометрические замеры ребенка? 1=да 0=нет <b>1→AN3</b>	AN4. Рост ребенка (см) <table border="1" style="float: right;"> <tr><td> </td><td> </td><td> </td><td> </td></tr> </table>									
AN2. Если нет, почему? 1 = не хотел (плакал, пинался и т.д.) 3=не пришли 2 = отказ матери/опекуна 4 другое (укажите) <b>1→BS1</b>	AN5. В каком положении был измерен рост ребенка -лежа или стоя? 1 = Лежа (дети до 24 месяцев) 2 = Стоя (дети 24-29 месяцев)									
AN3. Вес ребенка (кг) <table border="1" style="float: right;"> <tr><td> </td><td> </td><td> </td><td> </td></tr> </table>										
Модуль Анализ крови		BS								
В конце мы возьмем небольшой образец крови из пальца вашего ребенка. Это может доставить небольшой дискомфорт, но зато вы можете узнать есть ли у вашего ребенка анемия.										
BS1. Был ли получен анализ крови у ребенка? 1=да 0=нет <b>1→BS3</b>	BS4. Концентрация гемоглобина в Гемокью (поставьте 88.8 если не было измерено/не знаю) <table border="1" style="float: right;"> <tr><td> </td><td> </td><td> </td><td> </td></tr> </table> g/dL									
BS2. Если нет, почему? 1 = не хотел (плакал, пинался и т.д.) 4 = технические проблемы 2 = отказ матери/опекуна 5 = другое (укажите) 3 = не пришли <b>перейти на BS5</b>	BS5. Идентификационная бирка РЕБЕНКА Прикрепите здесь бирку ребенка									
BS3. Сколько примерно микролитров крови было собрано в микротейнер? <table border="1" style="float: right;"> <tr><td> </td><td> </td><td> </td><td> </td></tr> </table>										
Подпись координатора, подтверждающего, что ребенок был направлен в медучреждение для лечения, если уровень гемоглобина <8.0 g/dL _____ Дата _____										
Не забудьте отдать матери результаты измерения уровня гемоглобина ее ребенка. Попросите мать подписать здесь для подтверждения получения результатов измерения уровня гемоглобина и медицинского направления (если необходимо) _____ Дата _____										
Подпись координатора подтверждающего, что вопросник был проверен и заполнен: _____ Дата _____										
Комментарии интервьюера: _____ _____										



## APPENDIX II LIST OF CHILD HEMOGLOBIN RESULTS: FOR VILLAGE MEDICAL ATTENDANT

### List of Child Hemoglobin Results: For Village Medical Attendant

*Team supervisor: record hemoglobin results for all children in each cluster and indicating anyone with an Hgb co less than 8.0 g/dL; gave this form to the medical attendant at the end of each day.*

Team #:

Cluster #:

Date          
d d m m y y

No.	Child name	Mother	Hb level (g/dLiter)	Hb < 8.0 g/dLiter? (Yes/No)	Commen
01					
02					
03					
04					
05					
06					
07					
08					
09					
10					
11					
12					
13					
14					
15					
16					
17					
18					
19					
20					
21					
22					
23					
24					

### APPENDIX III CONFIDENCE INTERVALS, DESIGN EFFECTS, AND INTRA-CLASS CORRELATION COEFFICIENTS FOR MAJOR INDICATORS, CHILDREN 6–29 MONTHS — KYRGYZSTAN NATIONAL SURVEY, 2013

Indicator	Sample Size	Prevalence (%)	95% Confidence Interval	DEFF <sup>a</sup>	Number of Clusters	Average Cluster Size	ICC <sup>b</sup>
Anemia	2156	26	(22.2 - 29.8)	4.2	80	26.95	0.123
Hb<11.0 g/dL							
Serum Ferritin (<12µg/L)	2156	34.2	(31.3 - 37.2)	2.1	80	26.95	0.042
Serum Transferrin Receptor (sTfR) (>8.3 mg/L)	2156	39.3	(35.6 - 43.1)	3.2	80	26.95	0.085
Iron Deficiency Anemia (IDA)	2156	19.7	(16.5 - 22.9)	3.5	80	26.95	0.096
Hb<11.0 g/dL and sf<12 µg/L or elevated sTfR							
Serum Retinol Binding Protein (RBP) <0.71 µmol/L	2148	15.6	(13 - 18.1)	2.7	80	26.95	0.066
Alpha-1-glycoprotein (AGP) >1.0 g/L	2156	31.3	(27.9 - 34.7)	2.9	80	26.95	0.073
C-Reactive Protein (CRP) >5.0 mg/L	2156	13	(11.4 - 14.6)	1.3	80	26.95	0.012
Stunting	2149	11.7	(9.3 - 14.1)	3	80	26.95	0.077
Height-for-age z-score (HAZ) <-2 SD							
Underweight	2156	4.8	(3.7 - 5.9)	1.5	80	26.95	0.019
Weight-for-age z-score (WAZ) <-2 SD							
Wasting	2154	2.0	(1.1 - 2.8)	2.1	80	26.95	0.042
Weight-for-height z-score (WHZ) <-2 SD							

Note: CI=confidence interval; DEFF=design effect; ICC=intracluster correlation coefficient; Hb=hemoglobin; percent estimates weighted for non-response and 95% Confidence intervals adjusted for cluster survey design; average cluster size=sample size/ number of clusters.

<sup>a</sup>The design effect or DEFF is the ratio of the actual variance to the variance computed under the assumption of simple random sampling, thus calculating the loss of effectiveness by the use of cluster sampling, instead of simple random sampling; the larger the DEFF, the greater the variance.

<sup>b</sup> ICC=(DEFF-1)/(average cluster size - 1).

# APPENDIX IV

## QUESTIONNAIRES FOR 2011 AND 2012 LOT QUALITY ASSURANCE SAMPLING SURVEYS

Fill the following information before beginning the interview.

HH1. Cluster number:	<input type="text"/> <input type="text"/>	HH4. Supervisor's code	<input type="text"/> <input type="text"/>
HH2. Interviewer code	<input type="text"/> <input type="text"/>	HH5. Data entry operator code	<input type="text"/>
HH3. Day/Month/Year of interview:	___ / ___ /11	HH6. Oblast	_____
HH8. Result of interview	<input type="checkbox"/>	HH7. Rayon	_____
		ID1. Child number	<input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>

Completed in person -1; Completed by phone -2; Refused in person - 3; Refused by phone - 4; Not available to interview - 5

If data is not collected on a child. Find the following information from the clinic, the medical worker, the VHC volunteer or by visiting the home of the mother/child.

<p>HH9. Reason for not completing interview?</p> <p>Family moved from village 1</p> <p>Mother or family refused 2</p> <p>Mother/caregiver is temporarily unavailable 3</p> <p>Other (specify) _____ 7</p> <p>Can't find out 8</p> <p>If 3, 7 or 8 try to identify family's phone number</p> <p>0-996- _____ - _____</p>	<p>HH10. What is the child's birthdate?</p> <p>___ / ___ / 20__</p> <p>Write 8/8/2088 if DK</p>
---	---

For all children in the survey: Please abstract the information below from the Green Journal

<p>G1. Has (Child's name) ever received Gulazyk? <input type="checkbox"/></p> <p>1 = yes 0=no</p> <p>If not listed in green Journal put 0</p>	<p>G2. Has the child received Gulazyk within 3 months of or to today's date? <input type="checkbox"/></p> <p>1 = yes 0=no</p> <p>If not listed in green Journal put 0. <b>Proceed to the interview!</b></p>
<p>G3. What is the last date that the child received Gulazyk according to the green journal?</p> <p>___ / ___ / 20__</p> <p>Write 8/8/2088 if DK</p>	

Hello, my name is \_\_\_ and I am working with the Ministry of Health and the National Statistics Committee. I am working on a joint project concerned with nutrition of children 6 to 24 months of age. I would like to talk to you about this and record your answers to some questions that I have about your child (NAME of child). The interview will take approximately 5 minutes.

All of the information we obtain will remain strictly confidential and nobody will know that the information is yours. Participation is completely voluntary. If we come to any point at which you do not want to answer, let me know and I will go on to the next question, or the interview can be stopped at any time. If you agree to participate, I would like you to sign this form. May we begin? Would you please sign?

If permission is given, ask respondent to sign here and begin the interview.

(Signature) \_\_\_\_\_ Name: \_\_\_\_\_

<p>HH11. What is the name of the child?</p> <p>Name: _____</p>	<p>HH12. In what month and year was (child's name) born? - Probe: What is his/her birthday?</p> <p>___ / ___ / 20__</p> <p>Write 8/8/2088 if DK</p>
--	---

Check the birthday with the calendar to be sure the child is between 6 and 24 months of age. If the mother is unsure of the birthday, check the birthdate with the registry at the clinic. If the child is not 6 to 24 months he cannot participate. Explain this to the mother and thank her for coming.

Receipt and Use of Gulazyk Module																															
<p>GU1. Now I would like to ask you some questions about Gulazyk. Gulazyk is a vitamin and mineral powder that can be used to make a young child's food more nutritious. We would like to speak to a person in the household who knows what the child eats. May I speak to this person now?            1 = yes 0=no 8=don't know <input type="checkbox"/></p> <p><i>If there is no person in the household who can answer questions about the child, ask when that person could be available and arrange a time to return if possible.</i></p>																															
<p>GU2. Have you ever received a package of Gulazyk like this for (child's name)? - <i>Show Gulazyk sachets</i> <input type="checkbox"/>            1 = yes 0=no 8=don't know</p> <p><i>If no, encourage her to visit the health clinic to receive Gulazyk. End interview.</i></p>	<p>GU7. How many sachets of Gulazyk has (child's name) consumed within 2 months <input type="text"/> <input type="text"/>            (put 88 if don't know/remember; put 00 if none)</p>																														
<p>GU3. Is (child's name) currently taking Gulazyk? <input type="checkbox"/>            1 = yes 0=no 8=don't know  <b>If 1→GU5</b></p>	<p>GU8. How many full sachets of Gulazyk for (child's name) do you have remaining right now? - <i>Ask to see the remaining Gulazyk for this child and count the number of full sachets. Put 88 if cannot locate remaining sachets. Put 99 if more than 99</i></p>																														
<p>GU4. Why (child's name) is not currently taking Gulazyk?  <i>Don't read out! Circle each answer mentioned.</i></p> <table style="width: 100%; border-collapse: collapse;"> <tr><td style="width: 80%;">Too difficult to remember to give Gulazyk</td><td style="width: 20%; text-align: center; border: 1px solid black;">A</td></tr> <tr><td>Child experienced side effects/diarrhea</td><td style="text-align: center; border: 1px solid black;">B</td></tr> <tr><td>Child experienced side effects/skin/allergy</td><td style="text-align: center; border: 1px solid black;">R</td></tr> <tr><td>Child experienced side effects/other (specify) _____</td><td style="text-align: center; border: 1px solid black;">S</td></tr> <tr><td>Child does not like food when Gulazyk is added</td><td style="text-align: center; border: 1px solid black;">C</td></tr> <tr><td>Gulazyk makes the food taste bad</td><td style="text-align: center; border: 1px solid black;">D</td></tr> <tr><td>Gulazyk changes the color of the food</td><td style="text-align: center; border: 1px solid black;">E</td></tr> <tr><td>Mother/caretaker doesn't like Gulazyk</td><td style="text-align: center; border: 1px solid black;">F</td></tr> <tr><td>Health care worker told me to stop using Gulazyk</td><td style="text-align: center; border: 1px solid black;">G</td></tr> <tr><td>Ran out of Gulazyk</td><td style="text-align: center; border: 1px solid black;">H</td></tr> <tr><td>Children should receive natural vitamins from food</td><td style="text-align: center; border: 1px solid black;">I</td></tr> <tr><td>We have been away from home</td><td style="text-align: center; border: 1px solid black;">L</td></tr> <tr><td>Child has been sick</td><td style="text-align: center; border: 1px solid black;">N</td></tr> <tr><td>Don't know</td><td style="text-align: center; border: 1px solid black;">O</td></tr> <tr><td>Other _____</td><td style="text-align: center; border: 1px solid black;">P</td></tr> </table> <p>(specify)</p>	Too difficult to remember to give Gulazyk	A	Child experienced side effects/diarrhea	B	Child experienced side effects/skin/allergy	R	Child experienced side effects/other (specify) _____	S	Child does not like food when Gulazyk is added	C	Gulazyk makes the food taste bad	D	Gulazyk changes the color of the food	E	Mother/caretaker doesn't like Gulazyk	F	Health care worker told me to stop using Gulazyk	G	Ran out of Gulazyk	H	Children should receive natural vitamins from food	I	We have been away from home	L	Child has been sick	N	Don't know	O	Other _____	P	<p>GU9. Have you noticed any changes in the color or taste of the food to which Gulazyk is added? <input type="text"/> <input type="text"/>            Yes 1 <input type="checkbox"/>            No 0 <input type="checkbox"/>            If 0 =&gt; GU10</p> <p>GU9a. Is this a concern for you or your child? <input type="checkbox"/>            Yes 1            No 0</p>
Too difficult to remember to give Gulazyk	A																														
Child experienced side effects/diarrhea	B																														
Child experienced side effects/skin/allergy	R																														
Child experienced side effects/other (specify) _____	S																														
Child does not like food when Gulazyk is added	C																														
Gulazyk makes the food taste bad	D																														
Gulazyk changes the color of the food	E																														
Mother/caretaker doesn't like Gulazyk	F																														
Health care worker told me to stop using Gulazyk	G																														
Ran out of Gulazyk	H																														
Children should receive natural vitamins from food	I																														
We have been away from home	L																														
Child has been sick	N																														
Don't know	O																														
Other _____	P																														
<p>GU5. How did you obtain Gulazyk for (child's name)? <input type="checkbox"/></p> <table style="width: 100%; border-collapse: collapse;"> <tr><td style="width: 80%;">I go to the health clinic</td><td style="width: 20%; text-align: center;">1</td></tr> <tr><td>The medical worker brings Gulazyk to my home</td><td style="text-align: center;">2</td></tr> <tr><td>Both (of above answers)</td><td style="text-align: center;">3</td></tr> <tr><td>Other (specify) _____</td><td style="text-align: center;">8</td></tr> </table>	I go to the health clinic	1	The medical worker brings Gulazyk to my home	2	Both (of above answers)	3	Other (specify) _____	8	<p>GU10. When you run out of Gulazyk sachets do you plan to continue? <input type="checkbox"/>            1 = yes 0=no 8=don't know            If 0 or 8 =&gt; GU11, otherwise <b>End Interview</b></p>																						
I go to the health clinic	1																														
The medical worker brings Gulazyk to my home	2																														
Both (of above answers)	3																														
Other (specify) _____	8																														
<p>GU6. How often does/did (child's name) consume Gulazyk? <input type="checkbox"/></p> <p>1 - Every day for one month            2 - A sachet per day for 15 days, followed by a break for 15 days            3 - Every other day            4 - Other schedule (specify) _____            5 - Never yet received =&gt;GU10            8 - Don't know</p>	<p>GU11. What are the reasons that you will probably not continue to give Gulazyk? - <i>Do NOT read each item, circle all answers mentioned</i></p> <p>A - Too difficult to remember to give Gulazyk            B - Child experienced side effects            C - Child does not like food when Gulazyk is added            D - Family member doesn't want to use            E - Health care worker told me to stop using Gulazyk            F - Children should receive natural vitamins from diverse food            H - By the time I run out of Gulazyk my child will be &gt;24 months            I - Other (specify) _____            J - Don't know</p> <p><b>End Interview</b></p>																														

## ОПРОСНИК LQAS ПО ПРОГРАММЕ ГУЛАЗЫК

*Перед началом интервью укажите следующую информацию.*

НН1. Номер кластера:    
 НН4. Код супервайзера

НН2. Код интервьюера  
 НН5. Код оператора по воду данных

НН3. дд/мм/гг проведения интервью: \_\_\_/\_\_\_/12
 НН6. Область \_\_\_\_\_

НН8. Результат интервью 
 НН7. Район \_\_\_\_\_

ID. Номер ребенка

*Заполнено при встрече -1; Заполнено по телефону -2; Отказ участвовать в интервью при встрече - 3; Отказ участвовать в интервью по телефону – 4; Интервью не состоялось по причине отсутствия опрашиваемого - 5*

*В случае если не собраны данные о ребенке, обратитесь в медицинское учреждение, к медицинскому работнику или волонтеру СКЗ за информацией или посетите мать/ребенка на дому.*

НН9. Причина по которой интервью не состоялось? Семья съехала с деревни 1 Отказ матери или семьи 2 Мать/родитель временно не доступны 3 Другое (укажите) _____ 7 Не смогли найти 8 <i>В случае 3, 7 или 8 постарайтесь найти номер телефона семьи 0-996-_____ - _____</i>	НН10. Дата рождения ребенка ___/___/20___  <i>Напишите 8/8/2088 если не знает</i>
---	--

*Для всех детей, охваченных опросом: Пожалуйста, при ответе на нижеуказанные вопросы приведите информацию из зеленого журнала*

G1. Получал ли (имя ребенка) когда-нибудь Гулазык? 1 = да 0=нет <input type="checkbox"/> <i>Если нет записи в зеленом журнале, укажите 0</i>	G2. Получал ли ребенок Гулазык в течение последних 3 месяцев или в настоящее время? 1 = да 0=нет <input type="checkbox"/> <i>Если нет записи в зеленом журнале или со дня последней записи прошло более 3-х месяцев, укажите 0</i>
--	--

G3. Согласно зеленому журналу укажите последнюю дату когда ребенок получал Гулазык?  
 \_\_\_/\_\_\_/20\_\_\_  
*Напишите 8/8/2088 если не знает*

Здравствуйте, меня зовут \_\_\_\_\_. Я работаю с Министерством здравоохранения и Национальным статистическим комитетом по программе Гулазык. Я хотел(а) бы поговорить с Вами на эту тему и записать ваши ответы на некоторые из наших вопросов относительно вашего ребенка. Это интервью займет приблизительно 5 минут.

Вся полученная нами информация будет носить строго конфиденциальный характер, и никто не узнает о том, что полученная информация была предоставлена Вами. Ваше участие является добровольным. Если вы не хотите отвечать на какой либо вопрос, мы можем перейти на следующий или прервать интервью. Если вы согласны принять участие в нашем опросе, то, пожалуйста, подпишите данную форму. Можно начинать? Подпишите пожалуйста.

*Если разрешение получено, обратитесь с просьбой к респонденту поставить здесь подпись и начинайте интервью.*

(Подпись) \_\_\_\_\_ Фамилия \_\_\_\_\_

НН11. Назовите имя ребенка? Имя: _____	НН12. Укажите дату рождения (имя ребенка) ? ___/___/20___ <i>Запишите 8/8/2088 если не знает</i>
---	--

*Убедитесь по календарю, что ребенок находится в возрасте от 6 до 24 месяцев. Если мама не помнит точную дату рождения, уточните эту дату в медпункте. Если ребенок не находится в возрасте от 6 до 24 месяцев, то он не может участвовать. Объясните ситуацию матери и поблагодарите ее за то, что она согласилась на интервью.*

Как получают и используют Гулазык																																	
<p>GU1. Теперь я хотел бы задать вам несколько вопросов в отношении Гулазыка. Гулазык - это порошок, содержащий витамины и минералы, который можно использовать для того чтобы сделать еду вашего ребенка более питательной. Мы хотели бы поговорить с тем членом семьи, который знает чем ребенок питается. Можно поговорить с этим человеком сейчас?</p> <p>1 = да 0=нет 8=не знаю <input type="checkbox"/></p> <p><i>Если дома не оказалось того человека, который мог бы ответить на вопросы о ребенке, спросите когда можно будет с ним поговорить и назначьте время для повторной встречи, если возможно.</i></p>																																	
<p>GU2. Получали ли вы когда-нибудь вот такие пакетики Гулазыка для (имя ребенка)? - <i>Покажите пакетики Гулазыка</i></p> <p>1 = да 0=нет 8=не знаю <input type="checkbox"/></p> <p><i>Если нет, посоветуйте обратиться в медицинское учреждение для получения Гулазыка. <b>Конец интервью.</b></i></p>	<p>GU7. Сколько пакетиков Гулазыка (имя ребенка) употребил за последние 2 месяца <input type="text"/></p> <p><i>(поставьте 88 в случае ответа не знаю/не помню; поставьте 00 в случае отсутствия ответа)</i></p>																																
<p>GU3. Принимает ли (имя ребенка) Гулазык в настоящее время?</p> <p>1 = да 0=нет 8=не знаю <input type="checkbox"/></p> <p><b>If 1=&gt;GU5</b></p>	<p>GU8. Сколько полных пакетиков Гулазыка осталось у (имя ребенка) в данный момент? <input type="text"/></p> <p><i>Спросите можно ли посмотреть на оставшийся для ребенка запас Гулазыка и посчитайте количество полных пакетиков. Поставьте 88 в случае если не могут найти оставшиеся пакетики. Поставьте 99, если пакетиков 99 и более.</i></p>																																
<p>GU4. Почему (имя ребенка) в настоящее время не принимает Гулазык?</p> <p><i>Не зачитывайте варианты ответов.</i></p> <table border="0"> <tr><td>Очень трудно вспомнить о том, что надо дать Г.</td><td><input type="checkbox"/></td></tr> <tr><td>Наблюдались побочные эффекты, /диарея</td><td><input type="checkbox"/></td></tr> <tr><td>Наблюдались побочные эффекты, сыпь/аллергия</td><td><input type="checkbox"/></td></tr> <tr><td>Наблюдались побочные эффекты/прочие (укажите) _____</td><td><input type="checkbox"/></td></tr> <tr><td>Ребенку не нравится пища, когда в нее добавлен Г,</td><td><input type="checkbox"/></td></tr> <tr><td>Гулазык портит вкус пищи</td><td><input type="checkbox"/></td></tr> <tr><td>Гулазык меняет цвет пищи</td><td><input type="checkbox"/></td></tr> <tr><td>Матери/воспитателю не нравится Гулазык</td><td><input type="checkbox"/></td></tr> <tr><td>Медработник сказал мне прекратить прием Г.</td><td><input type="checkbox"/></td></tr> <tr><td>Закончился запас Гулазыка</td><td><input type="checkbox"/></td></tr> <tr><td>Дети должны получать натуральные витамины посредством разнообразного питания</td><td><input type="checkbox"/></td></tr> <tr><td>Прием Гулазыка может привести к смерти ребенка</td><td><input type="checkbox"/></td></tr> <tr><td>Находились вдали от дома</td><td><input type="checkbox"/></td></tr> <tr><td>Ребенок болел</td><td><input type="checkbox"/></td></tr> <tr><td>Не знаю</td><td><input type="checkbox"/></td></tr> <tr><td>Другое (укажите) _____</td><td><input type="checkbox"/></td></tr> </table>	Очень трудно вспомнить о том, что надо дать Г.	<input type="checkbox"/>	Наблюдались побочные эффекты, /диарея	<input type="checkbox"/>	Наблюдались побочные эффекты, сыпь/аллергия	<input type="checkbox"/>	Наблюдались побочные эффекты/прочие (укажите) _____	<input type="checkbox"/>	Ребенку не нравится пища, когда в нее добавлен Г,	<input type="checkbox"/>	Гулазык портит вкус пищи	<input type="checkbox"/>	Гулазык меняет цвет пищи	<input type="checkbox"/>	Матери/воспитателю не нравится Гулазык	<input type="checkbox"/>	Медработник сказал мне прекратить прием Г.	<input type="checkbox"/>	Закончился запас Гулазыка	<input type="checkbox"/>	Дети должны получать натуральные витамины посредством разнообразного питания	<input type="checkbox"/>	Прием Гулазыка может привести к смерти ребенка	<input type="checkbox"/>	Находились вдали от дома	<input type="checkbox"/>	Ребенок болел	<input type="checkbox"/>	Не знаю	<input type="checkbox"/>	Другое (укажите) _____	<input type="checkbox"/>	<p>GU9. Заметили ли вы какие-либо изменения в цвете или вкусе пищи, в которую добавлен Гулазык? <input type="checkbox"/></p> <p>Да 1</p> <p>Нет 0</p> <p><b>If 0 =&gt; GU10</b></p>
Очень трудно вспомнить о том, что надо дать Г.	<input type="checkbox"/>																																
Наблюдались побочные эффекты, /диарея	<input type="checkbox"/>																																
Наблюдались побочные эффекты, сыпь/аллергия	<input type="checkbox"/>																																
Наблюдались побочные эффекты/прочие (укажите) _____	<input type="checkbox"/>																																
Ребенку не нравится пища, когда в нее добавлен Г,	<input type="checkbox"/>																																
Гулазык портит вкус пищи	<input type="checkbox"/>																																
Гулазык меняет цвет пищи	<input type="checkbox"/>																																
Матери/воспитателю не нравится Гулазык	<input type="checkbox"/>																																
Медработник сказал мне прекратить прием Г.	<input type="checkbox"/>																																
Закончился запас Гулазыка	<input type="checkbox"/>																																
Дети должны получать натуральные витамины посредством разнообразного питания	<input type="checkbox"/>																																
Прием Гулазыка может привести к смерти ребенка	<input type="checkbox"/>																																
Находились вдали от дома	<input type="checkbox"/>																																
Ребенок болел	<input type="checkbox"/>																																
Не знаю	<input type="checkbox"/>																																
Другое (укажите) _____	<input type="checkbox"/>																																
	<p>GU9a. Если да, это беспокоит Вас или Вашего ребенка?</p> <p>Да 1 <input type="checkbox"/></p> <p>Нет 0</p>																																
	<p>GU10. Когда у вас закончатся пакетики Гулазыка. Вы планируете продолжить прием Гулазыка?</p> <p>1 = да 0=нет 8=не знаю <input type="checkbox"/></p> <p>Если 1 =&gt; GU12</p>																																
<p>GU5. Каким образом вы получаете Гулазык для (имя ребенка)? <input type="checkbox"/></p> <p>Я иду к врачу 1</p> <p>Медработник приносит Гулазык мне домой 2</p> <p>Оба варианта (вышеуказанные варианты ответа) 3</p> <p>Другое (укажите) _____ 8</p>	<p>GU11. По каким причинам вы не будете продолжать прием Гулазыка? – <i>НЕ зачитывайте каждый вариант ответа, обведите все полученные ответы</i></p> <table border="0"> <tr><td>Очень трудно помнить о том, что надо дать Гулазык</td><td><input type="checkbox"/></td></tr> <tr><td>У ребенка наблюдались побочные эффекты</td><td><input type="checkbox"/></td></tr> <tr><td>Ребенку не нравится пища, когда в нее добавляют Г.</td><td><input type="checkbox"/></td></tr> <tr><td>Члены семьи не хотят, чтобы ребенок потреблял Г.</td><td><input type="checkbox"/></td></tr> <tr><td>Медработник сказал мне прекратить прием Гулазыка</td><td><input type="checkbox"/></td></tr> <tr><td>Дети должны получать натуральные витамины посредством разнообразного питания</td><td><input type="checkbox"/></td></tr> </table>	Очень трудно помнить о том, что надо дать Гулазык	<input type="checkbox"/>	У ребенка наблюдались побочные эффекты	<input type="checkbox"/>	Ребенку не нравится пища, когда в нее добавляют Г.	<input type="checkbox"/>	Члены семьи не хотят, чтобы ребенок потреблял Г.	<input type="checkbox"/>	Медработник сказал мне прекратить прием Гулазыка	<input type="checkbox"/>	Дети должны получать натуральные витамины посредством разнообразного питания	<input type="checkbox"/>																				
Очень трудно помнить о том, что надо дать Гулазык	<input type="checkbox"/>																																
У ребенка наблюдались побочные эффекты	<input type="checkbox"/>																																
Ребенку не нравится пища, когда в нее добавляют Г.	<input type="checkbox"/>																																
Члены семьи не хотят, чтобы ребенок потреблял Г.	<input type="checkbox"/>																																
Медработник сказал мне прекратить прием Гулазыка	<input type="checkbox"/>																																
Дети должны получать натуральные витамины посредством разнообразного питания	<input type="checkbox"/>																																
<p>GU6. Как часто (имя ребенка) принимает Гулазык? <input type="checkbox"/></p> <p>1 – Каждый день в течение одного месяца</p> <p>2 – Пакетик в день в течение 15 дней с перерывом в 15 дней</p> <p>3 – Каждый второй день</p> <p>4 – Другой график приема</p> <p>5 - Никогда не принимал =&gt; <b>GU10</b></p> <p>8 - Не знаю</p>	<table border="0"> <tr><td>1) Прием Гулазыка может привести к смерти ребенка</td><td><input type="checkbox"/></td></tr> <tr><td>К тому времени, когда закончится у меня запас</td><td><input type="checkbox"/></td></tr> <tr><td>2) Гулазыка мой ребенок будет старше 24 месяцев</td><td><input type="checkbox"/></td></tr> <tr><td>Другое (укажите) _____</td><td><input type="checkbox"/></td></tr> <tr><td>3) Не знаю</td><td><input type="checkbox"/></td></tr> <tr><td>8</td><td><input type="checkbox"/></td></tr> </table>	1) Прием Гулазыка может привести к смерти ребенка	<input type="checkbox"/>	К тому времени, когда закончится у меня запас	<input type="checkbox"/>	2) Гулазыка мой ребенок будет старше 24 месяцев	<input type="checkbox"/>	Другое (укажите) _____	<input type="checkbox"/>	3) Не знаю	<input type="checkbox"/>	8	<input type="checkbox"/>																				
1) Прием Гулазыка может привести к смерти ребенка	<input type="checkbox"/>																																
К тому времени, когда закончится у меня запас	<input type="checkbox"/>																																
2) Гулазыка мой ребенок будет старше 24 месяцев	<input type="checkbox"/>																																
Другое (укажите) _____	<input type="checkbox"/>																																
3) Не знаю	<input type="checkbox"/>																																
8	<input type="checkbox"/>																																
	<p>GU12. Получали ли вы когда-нибудь брошюру «Гулазык»</p> <p>1 = да 0=нет 8=не знаю <input type="checkbox"/></p>																																

**Завершите интервью!**

---

## APPENDIX V

# ALGORITHMS FOR CALCULATING INDICATORS OF INFANT AND YOUNG CHILD FEEDING PRACTICES

The algorithm is patterned after the Indicators for Assessing Young Child Feeding Practices Part 2 (WHO, 2008); however the survey instruments used to assess infant feeding practices are not identical and therefore algorithm has been modified accordingly. Note that children with missing values or values out of acceptable range are excluded from all calculations

### 1. Early initiation of breastfeeding

**Definition:** Proportion of children 6 to 24 months who were put to the breast within one hour of birth.

$$\frac{\text{Children 6 to <24 months who were put to the breast within one hour of birth}}{\text{Children 6 to <24 months}}$$

**Calculation (ages 6 to <24 months only):**

$$\frac{\text{BF2=0}}{\text{All children 6 to <24 months}} \times 100$$

**Note:** Children who never initiated breast feeding (BF1=0) are included in the denominator.

### 2. Exclusive breastfeeding until 6 months of age

**Definition:** Proportion of infants fed exclusively with breast milk until 6 months of age.

$$\frac{\text{Children 6 to <24 months who were exclusively breast feed until 6 months of age}}{\text{Children 6 to <24 months}}$$

**Calculation (ages 6 to <24 months only):**

$$\frac{(\text{BF3} \geq 6 \text{ or } 99) \text{ AND } (\text{BF4} \geq 6 \text{ or } 99) \text{ AND } (\text{BF4} \geq 6 \text{ or } 99)}{\text{All children 6 to <24 months}} \times 100$$

### 3. Continued breastfeeding at 1 year

**Definition:** Proportion of children 12–15 months of age who are fed breast milk.

$$\frac{\text{Children 12–15 months of age who received breast milk during the previous day}}{\text{Children 12–15 months of age}}$$

**Calculation (ages 12 to 15 months only):**

$$\frac{(\text{BF6a} = 1)}{\text{Children 12 to 15 months of age}} \times 100$$

### 4. Introduction of solid, semi-solid or soft foods

**Definition:** Proportion of infants 6–8 months of age who receive solid, semi-solid or soft foods.

$$\frac{\text{Infants 6–8 months of age who received solid, semi-solid or soft foods during the previous day}}{\text{Infants 6–8 months of age}}$$

---

**Calculation (ages 6 to 8 months only):**

$$\frac{\text{BF7a} = 1 \text{ or } 2 \text{ or } 3 \text{ or } 4 \text{ or } 5 \text{ or } 6 \text{ or } 7}{\text{Children 6 to 8 months of age}} \times 100$$

**5. Minimum dietary diversity**

**Definition:** Proportion of children 6–23 months of age who receive foods from 4 or more food groups.

$$\frac{\text{Children 6–23 months of age who received foods from } \geq 4 \text{ food groups during the previous day}}{\text{Children 6–23 months of age}}$$

**Calculation (6 to <24 months only):**

To calculate a value for this indicator, a 7 food group score variable needs to be created. The instructions below show how to calculate the 7 food group score. This is followed by instructions to calculate the Minimum dietary diversity indicator.

**Calculation of 7 food group score:**

The 7 foods groups used for calculation of this indicator are:

1. grains, roots and tubers
2. legumes and nuts
3. flesh foods (meat, fish, poultry and liver/organ meats)
4. eggs
5. vitamin-A rich fruits and vegetables
6. other fruits and vegetables
7. dairy products (milk, yogurt, cheese)

Construct the 7 food group score as follows:

Begin with a score of 0.

For each of the 7 food groups, add a point if any food in the group was consumed.

Food group 1 Add 1 point if: Bf6e=1 or Bf6j=1 or Bf6k=1

Food group 2 Add 1 point if: Bf6d=1

Food group 3 Add 1 point if: Bf6f=1

Food group 4 Add 1 point if: Bf6g=1

Food group 5 Add 1 point if: Bf6h=1

Food group 6 Add 1 point if: Bf6i=1

Food group 7 Add 1 point if: Bf6b=1 or Bf6c=1

**Calculation of Minimum dietary diversity indicator:**

$$\frac{7 \text{ food group score } \geq 4}{\text{All children 6 to } < 24 \text{ months}} \times 100$$

**6. Minimum meal frequency**

**Definition:** Proportion of breastfed and non-breastfed children 6–23 months of age who receive solid, semi-solid, or soft foods (but also including milk feeds for non-breastfed children) the minimum number of times or more.

Breastfed children 6–23 months of age who received solid, semi-solid or soft foods the minimum number of times or more during the previous day \_\_\_\_\_ X100

Breastfed children 6–23 months of age

**and**



---

Non-breastfed children 6–23 months of age who received solid, semi-solid or soft foods or milk feeds the minimum number of times or more during the previous day \_\_\_\_\_X100\_

Non-breastfed children 6–23 months of age

**Calculation:**

To calculate a value for the overall indicator of Minimum Meal Frequency, combine the two numerators shown above, and the two denominators.

$[(Bf6a=1) \text{ AND } (age=6 \text{ to } 8 \text{ months}) \text{ and } BF7 \geq 2] \text{ OR } [(Bf6a=1) \text{ AND } (age=9 \text{ to } <24 \text{ months}) \text{ AND } (Bf6a \geq 3)] \text{ OR } [(Bf6a=2) \text{ AND } (age=6 \text{ to } 24 \text{ months}) \text{ AND } (Bf7a + Bf7c) \geq 4]$  \_\_\_\_\_X100

---

All children 6 to <24 months

**Notes:**

- For breastfed children, the minimum number of times varies with age (2 times if 6–8 months and 3 times if 9–23 months). For non-breastfed children the minimum number of times does not vary by age (4 times for all children 6–23 months)
- Bf6b and Bf6c ask about whether the child consumed infant formula, milk, etc. yesterday (each yes answer is counted in the numerator for non-breastfed children only).

**Minimum acceptable diet**

**Definition:** Proportion of children 6–23 months of age who receive a minimum acceptable diet (apart from breast milk).

Breastfed children 6–23 months of age who had at least the minimum dietary diversity and the minimum meal frequency during the previous day \_\_\_\_\_

Breastfed children 6–23 months of age

**and**

Non-breastfed children 6–23 months of age who received at least 2 milk feedings and had at least the minimum dietary diversity not including milk feeds and the minimum meal frequency during the previous day

Non-breastfed children 6–23 months of age

**Calculation of Minimum acceptable diet indicator:**

For non-breastfed children, the dietary diversity component of this indicator is different than for the Minimum dietary diversity indicator. A 6 food group score (instead of a 7 food group score) that excludes dairy products is used for non-breastfed children for this indicator. In addition, this indicator requires that non-breastfed children receive a minimum number of milk feeds.

The instructions below show how to calculate the 6 food group score for non-breastfed children. This is followed by instructions to calculate the Minimum acceptable diet indicator.

**Calculation of 6 food group score:**

The 6 foods groups used for calculation of the dietary diversity component of the indicator for non-breastfed children are:

1. grains, roots and tubers
2. legumes and nuts
3. flesh foods (meat, fish, poultry and liver/organ meats)
4. eggs
5. vitamin-A rich fruits and vegetables
6. other fruits and vegetables

---

Construct the 6 food group score as follows:

Begin with a score of 0.

For each of the 7 food groups, add a point if any food in the group was consumed.

Food group 1 Add 1 point if: Bf6e=1 or Bf6j=1 or Bf6k=1

Food group 2 Add 1 point if: Bf6d=1

Food group 3 Add 1 point if: Bf6f=1

Food group 4 Add 1 point if: Bf6g=1

Food group 5 Add 1 point if: Bf6h=1

Food group 6 Add 1 point if: Bf6i=1

(The 6 food group score is calculated by adding the 6 food groups the same as it was for minimum dietary diversity except for Food Group # 7 [Dairy products] which is not included in the calculation.)

**Calculation:**

[(Bf6a=1) AND (age 6 to 8 months) AND (7 food group score  $\geq$ 4) AND (BF7  $\geq$ 2)] OR [(Bf6a=1) AND (age 9 to <24 months) AND (7 food group score  $\geq$ 4) AND (BF7  $\geq$ 3)] OR ((Bf6a=0) AND (age 6 to <24 months) AND (BF7c  $\geq$ 2) AND (6 food group frequency score  $\geq$ 4) and AND (Bfa + Bf7c  $\geq$ 4))] X100

All children 6 to <24 months

**Children ever breastfed**

**Definition:** Proportion of children 6 to 24 months who were ever breastfed.

Children 6-24 months of age who were ever breastfed X100

Children born in the last 24 months

**Calculation (age 6 -24 months):**

$$\frac{BF1=1}{\text{Children 6 -24 months of age}} \times 100$$

**Continued breastfeeding at 2 years**

**Definition:** Proportion of children 20–23 months of age who are fed breast milk.

Children 20 to <24 months of age who received breast milk during the previous day

Children 20 to <24 months of age

**Calculation (children 20 to <24 months of age):**

$$\frac{(Bf6a=1) \text{ и } (Bf7a \geq 1)}{\text{Children 20 to <24 months of age}} \times 100$$

**Age-appropriate breastfeeding**

**Definition:** Proportion of children 6 to <24 months of age who are appropriately breastfed.

Children 6–23 months who received breast milk, as well as solid, semi-solid or soft foods, the previous day X100

Children 6–23 months of age

---

**Calculation (Children 6 to <24 months):**

$$\frac{(\text{Bf6a}=1) \text{ and } (\text{Bf7a} \geq 1)}{\text{All children 6 to } <24 \text{ months}} \times 100$$

**Milk feeding frequency for non-breastfed children**

**Definition:** Proportion of non-breastfed children 6–23 months of age who receive at least 2 milk feedings.

Non-breastfed children 6–23 months of age who received at least 2 milk feedings during the previous day \_\_\_\_ X100

Non-breastfed children 6–23 months of age

**Calculation (children 6 to <24 months of age):**

(Bf6a= 0) AND(Bf7c ≥2)

Bf6a=0)

## APPENDIX VI NATIONAL SURVEY OF NUTRITIONAL STATUS, KYRGYZSTAN 2013: RESULTS FOR THE WHO GLOBAL DATABASE ON CHILD GROWTH AND MALNUTRITION

GLOBAL DATABASE ON CHILD GROWTH AND MALNUTRITION

Ref. No.:

Country:

Author:

Reference:

Administrative level:

Month and year survey:

AGE GROUPS	N	WEIGHT//AGE (%)			HEIGHT (LENGTH)//AGE (%)			WEIGHT//HEIGHT (LENGTH) (%)			BMI//AGE (%)			NOTES										
		<-3 CO	<-2 CO <sup>1</sup>	Mean z-score	SD z-score	<-3 CO	<-2 CO <sup>1</sup>	Mean z-score	SD z-score	<-3 CO	<-2 CO <sup>1</sup>	Mean z-score	SD z-score											
(Months)																								
TOTAL (6-29)	2 162	0.4	4.8	0.2	1.1	3.3	11.7	0.0	1.3	0.6	2.0	19.7	3.2	0.3	0.3	1.1	0.6	2.1	22.7	4.4	0.7	0.3	1.2	
6-11	520	0.7	3.4	-0.2	1.0	1.4	5.0	-0.5	1.2	1.3	2.7	27.4	4.1	0.2	0.1	1.0	1.4	3.7	24.3	3.9	0.2	0.2	1.1	
12-17	569	0.7	3.7	-0.3	1.0	2.6	8.6	-0.9	1.2	0.7	2.5	17.1	3.7	0.6	0.1	1.0	0.7	2.5	19.9	4.8	0.7	0.3	1.0	
18-23	539	0.2	6.3	-0.4	0.9	4.1	17.1	-1.2	1.0	0.5	2.3	13.2	2.2	0.1	0.3	1.0	0.5	1.8	19.6	3.8	0.7	0.5	1.0	
24-29	534	0.2	5.6	0.2	1.1	4.8	15.5	0.0	1.3	0.0	0.5	21.9	2.8	0.3	0.3	1.1	0.0	0.5	27.1	4.9	1.3	0.3	1.2	

AGE GROUPS	N	WEIGHT/AGE (%)			HEIGHT (LENGTH)//AGE (%)			WEIGHT/HEIGHT (LENGTH)// (%)			BMI/AGE (%)			NOTES									
		<-3 CO	<-2 CO <sup>1</sup>	Mean z-score	SD z-score	<-3 CO	<-2 CO <sup>1</sup>	Mean z-score	SD z-score	<-3 CO	<-2 CO <sup>1</sup>	Mean z-score	SD z-score										
															>+1 CO <sup>1</sup>	>+2 CO <sup>1</sup>	>+3 CO	>+1 CO <sup>1</sup>	>+2 CO <sup>1</sup>	>+3 CO			
(Months)																							
Male (6-29)	1 102	0.8	5.3	0.2	1.2	3.8	13.1	-0.1	1.4	1.1	3.0	18.3	2.6	0.5	0.3	1.2	1.1	3.1	21.4	3.4	1.0	0.3	1.2
6-11	268	1.4	6.2	-0.3	1.1	2.4	8.9	-0.6	1.2	2.5	4.5	26.6	3.7	0.3	0.0	1.1	2.5	5.0	24.2	3.4	0.3	0.1	1.1
12-17	283	1.1	6.0	-0.3	1.0	3.8	10.0	-0.9	1.2	1.2	4.7	14.1	3.9	0.9	0.1	1.0	1.4	4.7	16.8	4.0	1.1	0.3	1.0
18-23	284	0.3	4.9	-0.4	1.0	3.9	16.4	-1.2	1.1	0.9	2.5	13.6	1.5	0.1	0.2	1.0	0.9	2.5	18.9	2.8	1.2	0.4	1.0
24-29	267	0.4	4.1	0.2	1.2	4.8	16.3	-0.1	1.4	0.0	0.4	19.9	1.7	0.5	0.3	1.2	0.0	0.4	25.9	3.3	1.4	0.3	1.2

GLOBAL DATABASE ON CHILD GROWTH AND MALNUTRITION

Ref. No.:

AGE GROUPS	N	WEIGHT/AGE (%)			HEIGHT(LENGTH)//AGE (%)			WEIGHT/HEIGHT(LENGTH)// (%)			BMI/AGE (%)			NOTES									
		<-3 CO	<-2 CO <sup>1</sup>	Mean z-score	SD z-score	<-3 CO	<-2 CO <sup>1</sup>	Mean z-score	SD z-score	<-3 CO	<-2 CO <sup>1</sup>	Mean z-score	SD z-score										
															>+1 CO <sup>1</sup>	>+2 CO <sup>1</sup>	>+3 CO	>+1 CO <sup>1</sup>	>+2 CO <sup>1</sup>	>+3 CO			
(Months)																							
Female (6-29)	1 060	0.1	4.3	0.2	1.1	2.7	10.3	0.0	1.3	0.1	1.0	21.1	3.7	0.1	0.3	1.1	0.1	1.0	23.9	5.4	0.4	0.3	1.1
6-11	252	0.0	0.6	-0.1	1.0	0.4	1.0	-0.5	1.1	0.2	0.9	28.2	4.5	0.0	0.2	1.0	0.4	2.3	24.3	4.5	0.0	0.2	1.0
12-17	286	0.3	1.6	-0.3	1.0	1.5	7.2	-0.9	1.2	0.1	0.4	20.0	3.6	0.3	0.1	1.0	0.0	0.4	22.8	5.5	0.3	0.3	1.0
18-23	255	0.1	7.9	-0.4	0.9	4.3	17.9	-1.1	1.0	0.1	2.2	12.9	3.0	0.1	0.3	0.9	0.0	1.1	20.3	4.9	0.1	0.5	1.0
24-29	267	0.0	7.1	0.2	1.1	4.8	14.7	0.0	1.3	0.0	0.5	24.0	3.9	0.2	0.3	1.1	0.0	0.5	28.5	6.7	1.1	0.3	1.1

	N	WEIGHT/AGE (%)		HEIGHT(LENGTH)//AGE (%)		WEIGHT/HEIGHT(LENGTH)/ (%)		BMI/AGE (%)		Mean	SD	NOTES											
		<-3 CO	<-2 CO <sup>1</sup>	Mean z-score	SD z-score	<-3 CO	<-2 CO <sup>1</sup>	>+1 CO <sup>1</sup>	>+2 CO <sup>1</sup>				>+3 CO	Mean z-score	SD z-score								
RESIDENCE																							
Urban	552	0.3	5.0	-0.2	1.1	3.6	11.3	-0.6	1.3	0.7	2.0	19.1	3.1	0.2	0.1	1.0	0.7	2.0	21.3	4.3	0.7	0.2	1.1
Rural	1 610	0.7	4.3	-0.2	1.0	2.5	12.6	-0.7	1.2	0.4	2.0	21.0	3.5	0.6	0.2	1.0	0.4	2.3	25.7	4.4	0.7	0.3	1.1
REGIONS																							
NOTES																							

1

% <-2SD includes %<-3SD; %>+2SD includes %>+3SD; %>+1SD includes %>+2SD and %>+3SD.

---

## **APPENDIX VII**

### **BRIEF SUMMARY QUALITY ASSURANCE FOR MICRONUTRIENT MEASUREMENTS FROM THE 2009 AND 2013 KYRGYZSTAN SURVEYS**

#### **2009 SURVEY**

##### **External Quality Assurance**

The VitMin Lab (Willstaett, Germany) has participated in CDC's external quality assurance program since 2006. The laboratory measures ferritin, soluble transferrin receptor (sTfR), C-reactive protein (CRP), and retinol binding protein (RBP) concentrations in plasma using an enzyme-linked immunosorbent assay (ELISA) technique. The precision and bias were Optimal or Desirable for all of the above indicators (>90% precision of the VITAL-EQA results, with <0.5% bias). This quality assurance analysis is based on exercises immediately preceding and during the survey (Rounds 12-13).

##### **Internal Quality Control**

The VitMin Lab (Willstaett, Germany) analyzed the survey samples for ferritin, sTfR, CRP, RBP, and AGP using an ELISA technique. The lab routinely tested a single QC pool in 16 different wells randomly distributed in each 384-well plate. The inter-assay coefficients variation (CV) for these analytes were 3.3% for ferritin, 3.7% for TfR, 8.9% for CRP, 3.0% for RBP and 3.5% for AGP. A CV of about 10% provides acceptable precision using an ELISA technique. These data indicate that the lab's performance was good to excellent while analyzing the survey samples.

#### **2013 SURVEY**

##### **External Quality Assurance**

The VitMin Lab (Willstaett, Germany) has participated in CDC's external quality assurance program, VITAL-EQA, since 2006. The laboratory measures ferritin, sTfR, CRP, and RBP concentrations in plasma using an enzyme-linked immunosorbent assay (ELISA) technique. The precision and bias were Optimal or Desirable for ferritin, sTfR, and CRP (>90% precision of the VITAL-EQA results, with <0.5% bias). For the 2013 survey, the precision and bias were Minimal or Unacceptable for RBP (>80-85% precision of the VITAL-EQA results, with 15-20% bias). This quality assurance analysis is based on exercises immediately preceding and during the survey (Rounds 20-21).

##### **Internal Quality Control**

The VitMin Lab analyzed the survey samples for ferritin, sTfR, CRP, RBP, and AGP using an ELISA technique. The lab routinely tested a single QC pool in 10 different wells randomly distributed in each 384-well plate. The inter-assay coefficients variation (CV) for these analytes were 3.8% for RBP, 3.2% for ferritin, 5.1% for AGP, 3.0% for TfR, and 5.2% for CRP. A CV of about 10% provides acceptable precision using an ELISA technique (Erhardt, 2004; Haynes, 2008). These data indicate that the lab's performance exceeded the acceptable performance expectations while analyzing the survey samples.





HH19. What is your name?  (last, first, middle initial)	HH23. What is your relationship to (child's name): mother 1 grandmother 2 aunt 3 other(specify) 6	<input type="text"/>	1-->HC1												
HH20. What is the name of the child?  (last, first, middle initial)	HH24. Why is (child's name)'s mother not here today? working/studying 1 sick 2 did not want to attend 3 family did not allow 4 busy at home/with housework 5 other (specify) 7 don't know 8	<input type="text"/>	<b>**IF MOTHER IS NOT PRESENT, RECORD REASON AND SKIP TO AN1**</b>												
HH21. In what day, month, and year was (child's name) born? <i>Probe: What is his/her birthday?</i> <table border="1" style="display: inline-table; vertical-align: middle;"> <tr> <td><input type="text"/></td> <td><input type="text"/></td> <td><input type="text"/></td> <td><input type="text"/></td> <td><input type="text"/></td> <td><input type="text"/></td> </tr> <tr> <td>d</td> <td>d</td> <td>m</td> <td>m</td> <td>y</td> <td>y</td> </tr> </table> <i>(put 88 for day, month, or year if don't know/remember)</i>	<input type="text"/>	<input type="text"/>		<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	d	d	m	m	y	y		
<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>										
d	d	m	m	y	y										
<i>Check to make sure the baby's birthday is between June 1, 2004 and December 31, 2008. If the birthday does not fall between these dates, the child cannot participate. Explain this to the mother and thank her for coming. If the mother is unsure of the birthday, check the birthdate with the registry at the clinic.</i>															
HH22. child's sex?		<input type="text"/>													
	1=male 2=female														
<b>Household Characteristics Module</b>			<b>HC</b>												
I would now like to ask you a few questions about your home and those who live in it.															
HC1. Does your family currently receive the universal monthly benefit?	HC2. Including (child's name) how many children age 6 - 59 months live in your home?	<input type="text"/>	<input type="text"/>												
<input type="text"/>	1=yes 0=no 8= don't know														

Woman's Module		WM
WM1. What is your native language? Kyrgyz 1 Russian 2 Kazakh 3 Uzbek 4 Other ( <i>specify</i> ) _____ 6 Don't know 8 <input type="text"/>		WM8. What type of work or study do you do? <input type="text"/> laborer (in the fields) 1 vender of food, fruit, homemade goods or other 2 employee in a business 3 business owner 4 professional (nurse, doctor, teacher, pharmacists, etc) 5 student 6 other (specific) _____ 7 Don't know 8
WM2. What is your date of birth? <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> d d m m y y (put 88 for day, month, or year if don't know/remember)		WM9. How many hours a day do you USUALLY work or study outside of the home? (put 88 if don't know) <input type="text"/> <input type="text"/>
WM3. How many live children do you have? <input type="text"/> <input type="text"/>		WM10. Who USUALLY takes care of (child's name) while you are outside of the home? <input type="text"/> The mother ( <i>takes the child with her</i> ) 1 Baby's grandmother 2 Baby's sisters/brothers 3 Baby's father 4 Other family member 5 Baby sitter 6 Day care / children's garden 7 Other (specify) _____ 8 Don't know 9
WM4. What is the highest level of school you completed? <input type="text"/> Never attended 0 Primary (1-4 grades) 1 Incomplete secondary (5-9) 2 Complete secondary 3 Technical school 4 Higher 5 Religious curriculum 6 Don't know 8		WM11. Who USUALLY feeds (child's name) while you are outside of the home? <input type="text"/> The mother ( <i>takes the child with her</i> ) 1 Baby's grandmother 2 Baby's sisters/brothers 3 Baby's father 4 Other family member 5 Baby sitter 6 Day care / children's garden 7 Other (specify) _____ 8 Don't know 9
WM5. Are you currently married? <input type="text"/> 1= yes 0= no <b>0→WM7</b>		WM12. How often does someone other than the mother feed (child's name) meals? <input type="text"/> never 0 < 1 time / day 1 1 time / day 2 2 times / day 3 3 times / day 4 > 3 times / day 5 Don't know 8
WM6. What is the highest level of school your spouse completed? <input type="text"/> Never attended 0 Primary (1-4 grades) 1 Incomplete secondary (5-9) 2 Complete secondary 3 Technical school 4 Higher 5 Religious curriculum 6 Don't know 8		Sometimes if mothers have to leave their child with a friend or family member while they are out of the house, they may not know everything the baby eats because someone else feeds them meals or snacks.....
WM7. Do you currently work or study outside the home (for example, as an employee, business owner, laborer in fields, etc.)? <input type="text"/> 1 = yes 0 = no <b>0→WM12</b>		WM13. Using the scale, can you estimate how much you know about what (child's name) usually eats? <input type="text"/> (show scale and note number that corresponds to the answer)

Breastfeeding and Infant Feeding		BF
Now I would like to ask you some questions about the breastfeeding and feeding of (child's name)		
BF1. Was (child's name) ever breastfed? 1=yes 0=no	<input type="text"/>	0→BF3
BF2. Approximately, how long after birth was (child's name) first put to the breast? Immediately (< 1 hour after birth) 0 During first 24 hours 1 Between 24 - 48 hours 2 > 48 hours 3 Don't know/remember 8	<input type="text"/>	
The next few questions are about the first time (child's name) was fed something other than breastmilk.		
BF3. How old was [child's name] in months when (he/she) was first fed animal milk, powdered milk or formula? <i>(if less than 1 month put 00, if NEVER fed milk, powdered milk or formula put 99, if don't know put 88)</i> <input type="text"/> <input type="text"/> (round down to nearest whole month) m m		
BF4. The next question is about liquids. Please include all liquids such as animal milk, powdered milk, formula, juice, water, sugar or fruit water, tea, or anything else that (child's name) might have been given. How old was (child's name) in months when he/she was first given any liquid, even tea, other than breastmilk? <i>(if less than 1 month put 00, if NEVER fed anything other than breastmilk put 99 if don't know put 88)</i> <input type="text"/> <input type="text"/> (round down to nearest whole month) m m		
BF5. The next question is about solid or semi-solid foods. Please include all solids such as porridge, rice, cereal, bulymak or anything else that (child's name) might have been given. How old was (child's name) in months when he/she was first fed any solid food? <i>(if less than 1 month put 00, if NEVER fed anything other than breastmilk put 99 if don't know put 88)</i> <input type="text"/> <input type="text"/> (round down to nearest whole month) m m		
Now think about everything (child's name) has drunk or eaten since this time yesterday. Don't forget snacks and eating or drinking during the night or things (child's name) ate with someone other than yourself.		
BF6. Since this time yesterday, was (child's name) fed any of the following items? <i>(read each item aloud and record response before proceeding to the next item)</i> 1=yes 0=no 8= don't know		
a Breastmilk	a	<input type="text"/>
b Animal milk, yogurt, kefir, cheese, etc	b	<input type="text"/>
c infant formula or powdered milk <i>(probe: what was the name?)</i> Brand name?	c	<input type="text"/>
d haricot, pea or nuts	d	<input type="text"/>
e kasha, potatoes, noodles, beet	e	<input type="text"/>
f meat, fish, poultry, liver/organ meat	f	<input type="text"/>
g eggs	g	<input type="text"/>
h carrots, pumpkin, tomatoes	h	<input type="text"/>
i other fruit or vegetable (spinach, dried apricots cucumbers)	i	<input type="text"/>
j bread or biscuit	j	<input type="text"/>
k baby cereal/food which was purchased Brand name?	k	<input type="text"/>
l any food with Sprinkles added (show packet)	l	<input type="text"/>
BF7. Since this time yesterday, how many times was (child's name) fed: <i>(if more than 7 put 7. If don't know put 8)</i> <i>("fed" means any meal or snack, excluding trivial amounts)</i>		
a any solid, semisolid, or soft food such as porridge, cereal, meat, vegetables, cookies, fruit, etc.	a	<input type="text"/>
b Breastmilk	b	<input type="text"/>
c animal milk, powdered milk or formula	c	<input type="text"/>
d anything from a bottle	d	<input type="text"/>
BF8. Has (child's name) stopped breastfeeding? 1=yes 0=no 0→AB1		
BF9. At what age in months did you stop breastfeeding (child's name)? <input type="text"/> <input type="text"/> <i>(put 88 if don't know/can't remember)</i> m m		

Attitude, Behavior Module		AB
<p>We are interested in knowing what mothers think about breastfeeding and feeding of their babies. I would like to ask you what you think about breastfeeding and feeding of your baby. Remember there are no right or wrong answers to any of these questions. We just want to know what you think about these topics.</p>		
<p>AB1. Using this scale, how would you describe the importance of breastfeeding for a baby's health and nutrition?</p> <p style="text-align: right;">[ ]</p> <p><i>(show the scale and note the number that corresponds to the answer)</i></p>	<p>AB8. In your opinion, what are some advantages to breastfeeding? <i>(don't read, mark all mentioned with 1)</i></p> <p>a healthy for baby and/or mother a [ ]</p> <p>b breastmilk is rich with vitamins/nutrients b [ ]</p> <p>c saves money c [ ]</p> <p>d saves time d [ ]</p> <p>e protects baby from infections e [ ]</p> <p>f safer than feeding from a bottle f [ ]</p> <p>g Other <i>(specify)</i> g [ ]</p>	
<p>AB2. Using this scale, how would you describe the importance of feeding other types of milk or formula for a baby's health and nutrition?</p> <p style="text-align: right;">[ ]</p> <p><i>(show the scale and note the number that corresponds to the answer)</i></p>	<p>AB9. Some people think there are disadvantages to breastfeeding while some people do not. In your opinion, are there disadvantages to breastfeeding?</p> <p style="text-align: right;">[ ]</p> <p>1=yes 0=no 8= don't know 0→DA1 8→DA1</p>	
<p>AB3. In your opinion, should a baby be breastfed?</p> <p style="text-align: center;">1=yes 0=no [ ]</p> <p style="text-align: right;">0→AB5</p>	<p>AB10. In your opinion, what are some disadvantages to breastfeeding? The things that make it more difficult.</p> <p><i>(don't read, mark all mentioned with 1)</i></p> <p>a mother cannot leave baby for very long (ie, to work or be outside the home) a [ ]</p> <p>b mother must be very careful about her diet b [ ]</p> <p>c causes sore nipples c [ ]</p> <p>d concerned they are not producing enough milk d [ ]</p> <p>e concerned mother's milk does not contain enough nutrients e [ ]</p> <p>f Other <i>(specify)</i> f [ ]</p>	
<p>AB4. In your opinion, how long in months should a baby be breastfed?</p> <p style="text-align: center;">[ ] [ ]</p> <p><i>(note 00 if &lt; 1 m; 88 if don't know)</i> m m</p>	<p>AB7. Some people think there are advantages to breastfeeding while some people do not. In your opinion, are there advantages to breastfeeding?</p> <p style="text-align: center;">[ ]</p> <p>1=yes 0=no 8= don't know 0→AB9 8→AB9</p>	
<p>AB5. In your opinion, at what age in months should a baby start drinking other liquids like tea, water, milk, etc?</p> <p style="text-align: center;">[ ] [ ]</p> <p><i>(note 00 if &lt; 1 m; 88 if don't know)</i> m m</p>		
<p>AB6. In your opinion, at what age in months should a baby start eating foods like porridge, cereal, bulymak, etc?</p> <p style="text-align: center;">[ ] [ ]</p> <p><i>(note 00 if &lt; 1 m; 88 if don't know)</i> m m</p>		

Dietary Advice Module		DA
<p>When a woman is pregnant and after she has a baby, many people give advice on her diet, breastfeeding and feeding the baby. I want to ask just about the advice you have received, it doesn't matter if it is advice you followed or not. I am just interested in what people have told you and who you have heard it from.</p>		
<p>DA1. Did you ever receive advice on your diet or nutrition when you were pregnant?</p> <p>1=yes 0=no 8= don't know</p>	<input type="text"/> <p>0→DA4 8→DA4</p>	<p>DA7. Using the scale, how important is the advice we get on breastfeeding from a doctor, nurse, midwife or feldsher?</p> <input type="text"/>
<p>DA2. Did a doctor, nurse, midwife or feldsher give you advice on your diet?</p> <p>1= yes 0= no</p>	<input type="text"/>	<p>DA8. Did family, friends or neighbors give you advice on breastfeeding?</p> <input type="text"/>
<p>DA3. Did a family member, friend or neighbor give you advice on your diet?</p> <p>1= yes 0= no</p>	<input type="text"/>	<p>1=yes 0= no 0→VS1</p>
<p>DA4. Did a doctor, nurse, midwife or feldsher give you advice on breastfeeding?</p> <p>1=yes 0= no 8= don't know</p>	<input type="text"/> <p>0→DA8 8→DA8</p>	<p>DA9. For how long (in months) did family, friends or neighbors advise you to breastfeed without giving other liquids or solids?</p> <input type="text"/> <input type="text"/> <small>(Put 00 if &lt; 1 m; 88 if don't know/remember; 99 if they did not give advice on length or did not specify exact length)</small>
<p>DA5. For how long (in months) did a doctor, nurse, midwife or feldsher advise you to breastfeed without giving other liquids or solids?</p> <input type="text"/> <input type="text"/> <small>(Put 00 if &lt; 1 m; 88 if don't know/remember; 99 if they did not give advice on length or did not specify exact length)</small>	<input type="text"/> <input type="text"/> <small>m m</small>	<p>DA10. At what age (in months) did family, friends, or neighbors advise you to stop breastfeeding?</p> <input type="text"/> <input type="text"/> <small>(Put 00 if &lt; 1 m; 88 if don't know/remember; 99 if they did not give advice on length or did not specify exact length)</small>
<p>DA6. At what age (in months) did a doctor, nurse, midwife or feldsher advise you to stop breastfeeding?</p> <input type="text"/> <input type="text"/> <small>(Put 00 if &lt; 1 m; 88 if don't know/remember; 99 if they did not give advice on length or did not specify exact length)</small>	<input type="text"/> <input type="text"/> <small>m m</small>	<p>DA11. Using the scale, how important is the advice we get on breastfeeding from family, friends or neighbors?</p> <input type="text"/> <small>(show the scale and note the number that corresponds to the answer)</small>
<p><b>Ask the Q. is at least on of answers of DA4, DA8=1</b></p>		
<p>DA12. Using the scale, rate the extent to which this advice would influence your own decisions regarding breastfeeding.</p> <input type="text"/> <small>(show the scale and note the number that corresponds to the answer)</small>		

Vitamins/Supplements Module		VS
I am now going to ask some questions about vitamins and supplements you and your baby might have taken. Some people take these supplements and some don't and that is okay. I will start with the supplements you might have taken.		
VS1. During your most recent pregnancy, did you take a folic acid supplement like this? (Show dispenser) <input type="text"/> 1=yes 0 = no 8= don't know	VS7. How long ago (in months) did (child's name) take the most recent vitamin A capsule? (note 00 if < 1 m; put 88 if don't know/remember) <input type="text"/> m <input type="text"/> m	
VS2. During your most recent pregnancy, did you take an iron supplement like this? (Show dispenser) <input type="text"/> 1=yes 0 = no 8= don't know	VS8. Have you ever been told by a doctor or nurse that (child's name) had anemia? 1=yes 0 = no 8= don't know	<input type="text"/> 0→VS10 8→VS10
VS3. In the first two months after the birth of your youngest child, did you take a Vitamin A dose like this? (Show Vitamin A capsule) <input type="text"/> 1=yes 0 = no 8= don't know	VS9. Did (child's name) take iron syrup or tablets to improve his/her anemia status? 1=yes 0 = no 8= don't know	<input type="text"/>
VS4. Have you ever been told by a doctor or nurse that you have anemia? <input type="text"/> 1=yes 0 = no 8= don't know 0→VS6 8→VS6	VS10. Have you, or someone else, ever given (child's name) any of these other vitamin or mineral supplements? (read the list and mark each answer)	a <input type="text"/> b <input type="text"/> c <input type="text"/> d <input type="text"/> 1=yes 0 = no 8= don't know
VS5. Did you take iron capsules or iron syrup to improve your anemia status? <input type="text"/> 1=yes 0 = no 8= don't know		
Now I'd like to ask a few questions about vitamins, minerals and supplements that (child's name) might have received. It is okay if (child's name) hasn't received these supplements.	VS11. Have you ever seen a Sprinkles package like this? Show Sprinkles sachet <input type="text"/> 1=yes 0 = no 8= don't know	<input type="text"/> 0→FF1 8→FF1
VS6. Has (child's name) ever taken a Vitamin A capsule like this one? Show 100,000IU for 6-11 month old <input type="text"/> 0→VS8 Show 200,000IU for 12-59 month old <input type="text"/> 8→VS8 1=yes 0 = no 8= don't know	VS12. Have you ever received a Sprinkles package like this? Show Sprinkles sachet <input type="text"/> 1=yes 0 = no 8= don't know	<input type="text"/> 0→FF1 8→FF1
	VS13. Has (child's name) ever consumed Sprinkles? <input type="text"/> 1=yes 0 = no 8= don't know	<input type="text"/>

Fortified Flour Module		FF
Now I would like to ask you about the flour you use for baking bread or cakes or any other food in your house.		
FF1. How much flour does your family consume in one month (in kilos)	<input type="text"/> <input type="text"/> . <input type="text"/>	FF8. In your opinion what are the benefits of fortified flour? (Don't read. Probe and mark all mentioned with a 1. If not mentioned, mark with a 0)
If the family does not consume/use flour skip to FF6		a Flour is better quality <input type="text"/>
FF2. What grade of flour do you usually use for cooking?		b Flour tastes better <input type="text"/>
Extra	1 <input type="text"/>	c Kids like the flour more <input type="text"/>
First grade	2 <input type="text"/>	d Makes children grow better <input type="text"/>
Second grade	3 <input type="text"/>	e Contains vitamins/minerals <input type="text"/>
Milled flour from own grain	4 <input type="text"/>	f It makes kids smarter <input type="text"/>
	1→FF4	g It makes kids stronger <input type="text"/>
	2→FF4	h It makes kids healthier <input type="text"/>
	3→FF4	i It makes women/men healthier <input type="text"/>
	4→FF3	j Prevents anemia <input type="text"/>
FF3. With which type of flour do you mix your milled flour from your own grain?		k Prevents illness <input type="text"/>
	<input type="text"/>	m It has adverse/negative effects <input type="text"/>
	0→FF6	n other (specify) <input type="text"/>
Do not mix milled flour with other flour	0	FF9. If you were given the choice of two loaves of bread of the same size and cost, but one had added iron and vitamins and the other did not, which would you prefer?
Extra	1	Loaf with added iron or vitamins <input type="text"/>
First grade	2	Loaf without added iron or vitamins <input type="text"/>
Second grade	3	Don't care <input type="text"/>
Don't know	8	Don't know <input type="text"/>
FF4 Where do you usually buy the flour for baking?		
Grocery store	1 <input type="text"/>	
Market	2	
Local mill	3	
other (specify) _____	4	
Don't know	8	
FF5. When you buy flour, what type do you most often purchase?		
Kazakh	1 <input type="text"/>	
National (Kyrgyz)	2	
Local (from your region)	3	
Don't know	8	
FF6. Have you ever heard about fortified flour?		
	<input type="text"/> 0→VHC1	
1=yes 0 = no		
FF7. Do you think there are any benefits to using fortified flour?		
	<input type="text"/>	
1=yes 0 = no 8=don't know		
	0→VHC1	
	0→VHC1	

VHC Contact Module		VHC	
<p>In some villages around Talas Oblast, there are Village Health Committees. I would like to ask you what you have heard of about these Village Health Committees, if anything. Remember no answers are right or wrong we just want to know what you have heard and your personal experience.</p>			
<p>VHC1. Have you ever heard of the Village Health Committee (VHC)?</p> <p style="text-align: right;"><input type="text"/></p> <p>1=yes 0 = no 8= don't know <b>0→P1</b></p>		<p>VHC4. Have you talked to a VHC member about your diet during pregnancy, or about breastfeeding or feeding your baby?</p> <p style="text-align: right;"><input type="text"/></p> <p>1=yes 0 = no 8= don't know <b>0→P1</b> <b>8→P1</b></p>	
<p>VHC2. Have you ever talked to someone from the VHC about health issues?</p> <p style="text-align: right;"><input type="text"/></p> <p>1=yes 0 = no 8= don't know <b>0→P1</b> <b>8→P1</b></p>		<p>VHC5. Using the scale, how helpful do you think the visit(s) with the VHC member was (the visit(s) on diet, breastfeeding and feeding a baby)?</p> <p style="text-align: right;"><input type="text"/></p> <p><i>(show the scale and note the number that corresponds to the answer)</i></p>	
<p>VHC3. How long ago was the last time you talked with a VHC member about health issues?</p> <p>&gt; 1 year ago                    0 <input type="text"/></p> <p>6-12 months ago                1</p> <p>3 - 6 months ago                2</p> <p>1- 3 months ago                 3</p> <p>&lt; 1 month ago                    4</p>		<p>VHC6. Are you interested in receiving more advice from the VHC?</p> <p style="text-align: right;"><input type="text"/></p> <p>1=yes 0 = no 8= don't know <b>0→P1</b> <b>8→P1</b></p>	
		<p>VHC7. What topics would you be interested in receiving advice about?</p> <p><i>(Write in all topics mentioned)</i></p> <p>_____</p> <p>_____</p>	



Pregnancy	
Before we continue, I need to know if you are pregnant. Even if you think you may be pregnant, but do not know for sure, we would still like to know that.	
P1. Are you pregnant right now? 1=yes 0= no 7=may be, but not sure 8= don't know	P2. How many weeks pregnant are you right now? (put 88 if don't know)
0→AN1 7→AN1 8→AN1	<input type="text"/> <input type="text"/>
If YES (1) Do not take blood or anthropometric measures from the mother. Take measurments only from the child.	
<b>Anthropometry AN</b>	
Now I am going to measure your height and weight and the length and weight of your baby.	
AN1. Were anthropometrics taken from mother? 1 = yes 0= no	AN5. Were anthropometrics taken from the child? 1 = yes 0= no
1→AN3	1→AN7
AN2. Why not? 1 = refused 3= not present 2 = pregnant 4 other (specify)	AN6. Why not? 1 = refused (cried, kicked, etc) 3=not present 2 = mother/guardian refused 4 other (specify)
AN3. Mother's height (cms) <input type="text"/> <input type="text"/> <input type="text"/> . <input type="text"/>	AN7. Child's weight (kg) <input type="text"/> <input type="text"/> . <input type="text"/>
AN4. Mother's weight (kg) <input type="text"/> <input type="text"/> <input type="text"/> . <input type="text"/>	AN8. Child's length/height (cm) <input type="text"/> <input type="text"/> <input type="text"/> . <input type="text"/>
<b>Blood Sample Module BS</b>	
The last thing we will do today is take a small sample of blood from your finger and the finger of your baby. This might cause a little discomfort from the stick but we will be able to tell you if your or your baby has anemia.	
BS1. What time did you eat for the last time? <input type="text"/> <input type="text"/> : <input type="text"/> <input type="text"/> h h m m	BS8. At what time did (child's name) eat for the last time? <input type="text"/> <input type="text"/> : <input type="text"/> <input type="text"/> h h m m
BS2. Was a capillary sample obtained from the mother? 1=yes 0= no	BS9. Was a capillary sample obtained from the child? 1=yes 0= no
1→BS4	1→BS11
BS3. Why not? 1 = refused 3=Not present →BS7 2 = pregnant 4=Technical difficulties 5=other (specify)	BS10. Why not? 1 = refused (cried, kicked, etc) 3=not present →BS14 2 = mother/guardian refused 4=technical difficulties 5=other (specify)
BS4. Approximately how many microliters of blood were collected in the microtainer? <input type="text"/> <input type="text"/> <input type="text"/>	BS11. Approximately how many microliters of blood were collected in the microtainer? <input type="text"/> <input type="text"/> <input type="text"/>
BS5. At what time was the sample obtained? <input type="text"/> <input type="text"/> : <input type="text"/> <input type="text"/> h h m m	BS12. At what time was the sample obtained? <input type="text"/> <input type="text"/> : <input type="text"/> <input type="text"/> h h m m
BS6. Hemoglobin concentration from Hemocue <input type="text"/> <input type="text"/> . <input type="text"/> g/dL (put 88.8 if not measured/don't know)	BS13. Hemoglobin concentration from Hemocue <input type="text"/> <input type="text"/> . <input type="text"/> g/dL (put 88.8 if not measured/don't know)
BS7. ID Label - MOTHER Affix mother label for blood here	BS14. ID Label - CHILD Affix child label for blood here

---

Signature of site supervisor confirming woman or child was referred to primary health care provider for treatment if hemoglobin <7.0 g/dL _____ Signature Date
Don't forget to provide the mother with the Hb measurement results for herself and her baby. Ask mother to sign here to confirm receipt of Hb measurement results and referral (if appropriate). _____ Signature Date
Signature of site supervisor confirming they have checked the questionnaire and it is complete: _____ Signature Date
Interviewer Comments: _____ _____

ИССЛЕДОВАНИЯ ПО ПИТАНИЮ НА НАЦИОНАЛЬНОМ УРОВНЕ					
Заполните следующую информацию до проведения интервью.					
НН1. Номер кластера	<input type="text"/>	НН4. Код руководителя	<input type="text"/>	<input type="text"/>	<input type="text"/>
НН2. Код интервьюера	<input type="text"/>	НН5. Код оператора ввода данных	<input type="text"/>	<input type="text"/>	<input type="text"/>
НН3. День/Месяц/Год интервью:	<input type="text"/>	НН6. Код области	<input type="text"/>	<input type="text"/>	<input type="text"/>
	<input type="text"/>	02 Иссык-Кульская	05 Баткенская	08 Чуйская	
Д	Д	М	М	Г	Г
			0		9
		03 Джалал-Абадская	06 Ошская	11 г.Бишкек	
		04 Нарынская	07 Таласская	21 г.Ош	
НН7. Результат интервью	<input type="text"/>	НН8. Результат антропометрии	<input type="text"/>	НН9. Результат забора крови	<input type="text"/>
Заполнено	1	Заполнено на мать/ребенка	1	Заполнено на мать/ребенка	1
Отказано	2	Заполнено только на мать	2	Заполнено только на мать	2
Заполнено частично	3	Заполнено только на ребенка	3	Заполнено только на ребенка	3
Нет на интервью	4	На обоих не заполнено	4	На обоих не заполнено	4
Наклейка с ИН - РЕБЕНОК Вклейте ее сюда			L1. На каком языке было проведено интервью?		
			На кыргызском языке 1		
			На русском языке 2		
			На другом (укажите) 3		
Если нет данных, собранных на маму и ребенка. Найдите информацию в клинике, у медицинского работника, члена СКЗ или посетив дом матери/ребенка.					
НН11. Сколько лет (в годах) матери?			НН15. Где живет семья?		
<input type="text"/>			<input type="text"/>		
<i>(поставьте 88, если не знаете/не можете выяснить)</i>			Около или в центре села /города 1		
			На окраине села/города 2		
НН12. Каков возраст (в месяцах) ребенка?			Не в селе/городе 3		
<input type="text"/>			Прочее (укажите) 4		
<i>(поставьте 88, если не знаете/не можете выяснить)</i>			Не знаю 8		
НН13. Каков пол ребенка?			НН16. Сколько братьев/сестер есть у ребенка?		
<input type="text"/>			<input type="text"/>		
1=мужской 2=женский			<i>(поставьте 88, если не знаете/не можете выяснить)</i>		
НН14. Какова этническая группа матери?			НН17. Работает/учится мать вне дома?		
Кыргызы 1 <input type="text"/>			<input type="text"/>		
Русские 2			1= да 0 = нет 8 = не знаю/не могу выяснить		
Казахи 3					
Узбеки 4			НН18. Причина неявки на интервью?		
Таджики 5			<input type="text"/>		
Уйгуры 6			семья переехала из села/города 1		
Прочее (укажите) 7			мать отказалась 2		
Не знаю 8			семья (муж, свекровь, и т.д.) отказалась 3		
			мать больна 4		
			ребенок болен 5		
			матери нужно работать 6		
			не была приглашена в поликлинику 7		
			Прочее (укажите) 8		
			Не знаю/не могу выяснить 9		

Мы из Министерства Здравоохранения и работаем по совместному проекту по вопросам питания и здоровья матери и ребенка. Я бы хотел(а) поговорить с вами об этом и записать ваши ответы на некоторые вопросы. На это интервью потребуется приблизительно 20 минут. После интервью мы взвесим и сделаем замеры вас и вашего ребенка, и возьмем небольшую пробу крови с вашего пальца и пальца вашего ребенка. На основе этой пробы мы сможем проинформировать вас, имеется ли у вас или вашего ребенка анемия. Единственной прямой выгодой для вас является знание статуса анемии у вас и вашего ребенка. Риск для вас маленький и состоит из возможного дискомфорта, вызванного прокалыванием пальца для получения пробы крови. Дискомфорт будет только временным и не будет очень большим. Вся полученная нами информация останется строго конфиденциальной и никто не будет знать, что это ваша информация, однако результаты гемоглобина будут переданы медицинскому работнику ФАПа/ГСВ. Вы вправе выбрать, участвовать в интервью, или нет, и если вы решите не участвовать, то это не будет иметь для вас никаких последствий. Если вы согласны участвовать, я бы хотел(а), чтобы вы подписали эту форму.

Мы можем начинать? Пожалуйста, подпишите.

Если вы получили разрешение, попросите респондента подписаться здесь, и начинайте интервью.

(Подпись) \_\_\_\_\_ (Ф.И.О.) \_\_\_\_\_

Согласие на участие женщины

(Подпись) \_\_\_\_\_ (Ф.И.О.) \_\_\_\_\_

Согласие на участие ребенка

НН19. Ваши Ф.И.О.? _____ (фамилия, имя, инициал отчества)	НН23. Кем вы являетесь (имя ребенка): <table border="1"> <tr> <td>мать</td> <td>1</td> <td rowspan="4" style="background-color: #cccccc; text-align: center;">1→НС1</td> </tr> <tr> <td>бабушка</td> <td>2</td> </tr> <tr> <td>тетя</td> <td>3</td> </tr> <tr> <td>прочее (укажите)</td> <td>6</td> </tr> </table>	мать	1	1→НС1	бабушка	2	тетя	3	прочее (укажите)	6													
мать	1	1→НС1																					
бабушка	2																						
тетя	3																						
прочее (укажите)	6																						
НН20. Имя ребенка? _____ (фамилия, имя, инициал отчества)	НН24. Почему (имя ребенка) мать не пришла сегодня? <table border="1"> <tr> <td>на работе/на учебе</td> <td>1</td> <td><input type="text"/></td> <td rowspan="8" style="background-color: #cccccc; text-align: center;">**ЕСЛИ МАТЬ ОТСУТСТВУЕТ, УКАЗАТЬ ПРИЧИНУ И ПЕРЕЙТИ К АН1**</td> </tr> <tr> <td>болеет</td> <td>2</td> <td></td> </tr> <tr> <td>не захотела прийти</td> <td>3</td> <td></td> </tr> <tr> <td>семья не разрешила</td> <td>4</td> <td></td> </tr> <tr> <td>занята дома/домашней работой</td> <td>5</td> <td></td> </tr> <tr> <td>прочее (укажите)</td> <td>7</td> <td></td> </tr> <tr> <td>не знаю</td> <td>8</td> <td></td> </tr> </table>	на работе/на учебе	1	<input type="text"/>	**ЕСЛИ МАТЬ ОТСУТСТВУЕТ, УКАЗАТЬ ПРИЧИНУ И ПЕРЕЙТИ К АН1**	болеет	2		не захотела прийти	3		семья не разрешила	4		занята дома/домашней работой	5		прочее (укажите)	7		не знаю	8	
на работе/на учебе		1	<input type="text"/>	**ЕСЛИ МАТЬ ОТСУТСТВУЕТ, УКАЗАТЬ ПРИЧИНУ И ПЕРЕЙТИ К АН1**																			
болеет	2																						
не захотела прийти	3																						
семья не разрешила	4																						
занята дома/домашней работой	5																						
прочее (укажите)	7																						
не знаю	8																						
НН21. День, месяц и год рождения (имя ребенка)? Спросите день его/ее рождения. <table border="1" style="width: 100%;"> <tr> <td style="width: 20px; text-align: center;">д</td> <td style="width: 20px; text-align: center;">д</td> <td style="width: 20px; text-align: center;">м</td> <td style="width: 20px; text-align: center;">м</td> <td style="width: 20px; text-align: center;">г</td> <td style="width: 20px; text-align: center;">г</td> </tr> </table> (напишите 88 вместо день, месяц, или год, в случае, если не знаете/ не помните)	д	д	м		м	г	г	Убедитесь, что ребенок родился в период между 1 июня 2004 г. и 31 декабря 2008 г. Если дата рождения не соответствует этому периоду, ребенок не может участвовать. Объясните это матери и поблагодарите ее за то, что пришла. Если мать не уверена, проверьте дату рождения по журналу регистрации в поликлинике.															
д	д	м	м	г	г																		
НН22. Каков пол ребенка? <input type="text"/> 1=мужской 2=женский																							
<b>Модуль характеристики домохозяйств</b> <span style="float: right;">НС</span>																							
Я теперь хотела бы задать вам несколько вопросов о вашем доме и о тех, кто живет в нем.																							
НС1. Ваша семья получает сейчас единое ежемесячное пособие малообеспеченным семьям (ЕЕП)?	НС2. Сколько детей включая (имя ребенка) в возрасте 6-59 месяцев живут в вашем доме?																						
<input type="text"/> <span style="background-color: #cccccc; padding: 2px;">1=да</span> <span style="padding: 2px;">0=нет</span> <span style="padding: 2px;">8= не знаю</span>	<input type="text"/>																						

Модуль для женщины		WM
WM1. Какой у вас [у матери] родной язык?		WM8. Какой вид работы или учебы у вас [у матери]?
Кыргызский 1		<input type="text"/>
Русский 2		работник (на полях) 1
Казахский 3		продавец продуктов, фруктов, товаров домашнего приг. и др. 2
Узбекский 4		сотрудник предприятия 3
Другой (укажите) _____ 6		владелец бизнеса 4
Не знаю 8 <input type="text"/>		специалист (медсестра, врач, учитель, фармацевт, т.д.) 5
WM2. Дата вашего [матери] рождения?		студент 6
<input type="text"/>		другое (укажите) _____ 7
Д Д М М Г Г		не знаю 8
<i>(напишите 88 вместо день, месяца, или год, если не знаете/не помните)</i>		WM9. Сколько часов в день вы [мать] ОБЫЧНО
WM3. Сколько живых детей вы [мать] имеете?	<input type="text"/>	находитесь на работе или учебе вне дома?
		<i>(напишите 88, если не знаете)</i>
WM4. Какой самый высокий уровень образования вы закончили?	<input type="text"/>	WM10. Кто ОБЫЧНО заботится о ребенке (имя
Никогда не посещала 0		ребенка) пока вы [мать] вне дома? <input type="text"/>
Начальное (1-4 класс) 1		Мать ( <i>берет ребенка с собой</i> ) 1
Неполное среднее (5-9) 2		Бабушка ребенка 2
Полное среднее 3		Сестры/братья ребенка 3
Средне-техническое образование 4		Отец ребенка 4
Высшее 5		Другой член семьи 5
Религиозная учебная программа 6		Няня 6
Не знаю 8		Детский сад 7
		Прочие (укажите) _____ 8
		Не знаю 9
WM5. Вы замужем в настоящее время?	<input type="text"/>	WM11. Кто ОБЫЧНО кормит ребенка (имя ребенка)
1= да 0= нет	<b>0→WM7</b>	пока вы [мать] вне дома? <input type="text"/>
WM6. Какой самый высокий уровень образования получил ваш		Мать ( <i>берет ребенка с собой</i> ) 1
супруг? <input type="text"/>		Бабушка ребенка 2
Никогда не посещал 0		Сестры/братья ребенка 3
Начальное (1-4 класс) 1		Отец ребенка 4
Неполное среднее (5-9) 2		Другой член семьи 5
Полное среднее 3		Няня 6
Средне-техническое образование 4		Детский сад 7
Высшее 5		Прочие (укажите) _____ 8
Религиозная учебная программа 6		Не знаю 9
Не знаю 8		WM12. Как часто кто-нибудь помимо матери кормит
WM7. В настоящее время вы [мать] работаете, либо учитесь		(имя ребенка)? <input type="text"/>
вне дома (например, как сотрудник, владелец бизнеса,		никогда 0
работник на поле, и т.д.)? <input type="text"/>		Менее 1 раза в день 1
1 = да 0 = нет	<b>0→WM12</b>	1 раз в день 2
		2 раза в день 3
		3 раза в день 4
		Более 3 раз в день 5
		Не знаю 8
		Иногда мамы оставляют своих детей с друзьями или членами семьи,
		чтобы куда-то уйти, поэтому они могут не знать всего, что ребенок ест в
		это время, потому что кто-то другой кормит ребенка.....
		WM13. Используя шкалу, оцените, как много вы знаете о том,
		что обычно ест (имя ребенка)? <input type="text"/>
		<i>(покажите шкалу и отметьте цифры, соответствующую ответу)</i>

Грудное вскармливание и питание младенца		BF
Теперь бы я хотела задать вам несколько вопросов о кормлении грудью и питании (имя ребенка)		
BF1. (имя ребенка) когда-либо кормили грудью? 1=да 0 = нет	<input type="text"/>	0→BF3
BF2. Приблизительно в течение какого времени после рождения (имя ребенка) был впервые приложен к груди? Сразу же (менее 1 часа после рождения) 0 В течение первых 24 часов 1 Между 24 - 48 часами 2 Более 48 часов 3 Не знаю/не помню 8	<input type="text"/>	
Следующие несколько вопросов касаются того, когда впервые (имя ребенка) дали что-то еще помимо грудного молока. BF3. Сколько было [имя ребенка] в месяцах, когда (ему/ей) дали впервые молоко животных, сухое молоко или смесь? (если младше 1 месяца, поставьте 00; если НИКОГДА не давали молоко, сухое молоко или смесь, поставьте 99; если не знаете, поставьте 88) <input type="text"/> <input type="text"/> (округлите в меньшую сторону до ближайшего целого месяца) М М		
BF4. Следующий вопрос будет о жидкостях. Пожалуйста, включите все жидкости, такие как молоко животных, сухое молоко, смесь, сок, воду, сладкую воду, компот, чай или другое, что могли давать (имя ребенка). Сколько было (имя ребенка) в месяцах, когда ему/ей впервые дали жидкость, даже чай, помимо грудного молока? (если младше 1 месяца, поставьте 00, если НИКОГДА не давали ничего, кроме грудного молока, поставьте 99, если не знаете, поставьте 88) <input type="text"/> <input type="text"/> (округлите в меньшую сторону до ближайшего целого месяца) М М		
BF5. Следующий вопрос будет о густой или твердой пище. Пожалуйста, включите всю твердую пищу, такую как каши, рис, злаковые, булмык или другое, что могли давать (имя ребенка). Сколько было (имя ребенка) в месяцах, когда ему/ей впервые дали твердую пищу? (если младше 1 месяца, поставьте 00, если НИКОГДА не давали ничего, кроме грудного молока, поставьте 99, если не знаете, поставьте 88) <input type="text"/> <input type="text"/> (округлите в меньшую сторону до ближайшего целого месяца) М М		
		Теперь вспомните обо всем, что (имя ребенка) выпил или съел с этого часа со вчерашнего дня. Не забудьте про легкий прием пищи и еду, или жидкости ночью, или что (имя ребенка) ел с кем-то, помимо вас.
		BF6. С этого времени вчера давалось ли (имя ребенка) что-нибудь из следующего: (прочитайте вслух каждое название и сделайте запись ответа, прежде чем приступить к следующему наименованию) 1=да 0 = нет 8= не знаю
		a Грудное молоко a <input type="text"/>
		b Молоко животных, йогурт, кефир, сыр и т.д. b <input type="text"/>
		c Молочная смесь, сухое молоко c <input type="text"/>
		(образец: как называется?) Название торговой марки?
		d фасоль, горох или орехи d <input type="text"/>
		e каша, картофель, лапша, свекла e <input type="text"/>
		f мясо, рыба, птица, печень/внутренности f <input type="text"/>
		g яйца g <input type="text"/>
		h морковь, тыква, помидоры h <input type="text"/>
		i другие фрукты или овощи (шпинат, сушеный урюк, огурцы) i <input type="text"/>
		j хлеб или печенье j <input type="text"/>
		k купленная детская крупа/питание k <input type="text"/>
		Название торговой марки? l любое питание с добавками Спринклз (показать пакет) l <input type="text"/>
		BF7. С этого часа вчера сколько раз кормили (имя ребенка): (если больше 7, поставьте 7. Если не знаете, поставьте 8) ("кормили" означает любая еда или перекус, не включая маленькие объемы)
		a любая твердая, густая или мягкая пища, типа каши, злаковых, мяса, овощей, печенья, фруктов и т.д. a <input type="text"/>
		b Грудное молоко b <input type="text"/>
		c Молоко животных, сухое молоко, смесь c <input type="text"/>
		d что-нибудь из бутылочки d <input type="text"/>
		BF8. Вы перестали кормить грудью (имя ребенка)? 1=да 0 = нет 0→AB1
		BF9. В каком возрасте в месяцах вы перестали кормить грудью (имя ребенка)? (поставьте 88, если не знаете/не помните) М М

Модуль отношения, поведения		AB
Мы заинтересованы в том, чтобы знать, что матери думают о грудном вскармливании и питании своих детей. Я хочу спросить вас, что вы думаете о грудном вскармливании и питании своего ребенка. Помните, что здесь нет ни правильных, ни неверных ответов на любой из этих вопросов. Мы просто хотим знать, что вы думаете по этому поводу.		
AB1. Используя эту шкалу, как бы вы описали важность грудного вскармливания для здоровья и питания ребенка? <input type="text"/> <i>(покажите шкалу и отметьте цифру, которая соответствует ответу)</i>	AB8. По вашему мнению, каковы преимущества грудного вскармливания? <i>(не читайте, отметьте упомянутое цифрой 1)</i> a полезно для малыша и/или матери a <input type="text"/> б грудное молоко богато витаминами/полезными веществами b <input type="text"/> в экономия денег c <input type="text"/> d экономия времени d <input type="text"/> e защита ребенка от инфекций e <input type="text"/> f безопаснее, чем кормление ч/з бутылочку f <input type="text"/> г Другое <i>(укажите)</i> g <input type="text"/>	
AB2. Используя эту шкалу, как бы вы описали важность кормления другими видами молока или смесью для здоровья и питания ребенка? <input type="text"/> <i>(покажите шкалу и отметьте цифру, которая соответствует ответу)</i>	AB9. Некоторые считают, что грудное вскармливание имеет недостатки, а некоторые так не считают. По-вашему, есть ли недостатки при грудном вскармливании? 1=да 0=нет 8= не знаю <input type="text"/>	
AB3. По вашему мнению, следует ли вскармливать ребенка грудью? 1=да 0=нет <input type="text"/> 0→AB5	AB10. По вашему мнению, каковы недостатки при грудном вскармливании? Причины, которые делают его более трудным. <i>(не читайте, отметьте упомянутое цифрой 1)</i> а мама не может оставить ребенка надолго (т.е., работа, гости) a <input type="text"/> б мама должна следить за своим питанием b <input type="text"/> в болезненность грудных сосков c <input type="text"/> г беспокойство о нехватке молока d <input type="text"/> е беспокойство о том, что молоко матери не содержит достаточно питательных веществ e <input type="text"/> ф Другое <i>(укажите)</i> f <input type="text"/>	
AB4. По вашему мнению, сколько месяцев следует вскармливать ребенка грудью? <i>(поставьте 00, если &lt; 1 мес; 88 - если не знаете)</i> <input type="text"/> М <input type="text"/> М	1=да 0=нет 8= не знаю <input type="text"/>	
AB5. По вашему мнению, в каком возрасте в месяцах ребенок должен начинать пить другие жидкости (чай, вода, молоко и т.д.)? <i>(поставьте 00, если &lt; 1 мес; 88 - если не знаете)</i> <input type="text"/> М <input type="text"/> М		
AB6. По вашему мнению, в каком возрасте в месяцах ребенок должен начинать есть пищу типа каши, злаковых, булмык, и т.д.? <i>(поставьте 00, если &lt; 1 мес; 88 - если не знаете)</i> <input type="text"/> М <input type="text"/> М		
AB7. Некоторые считают, что грудное вскармливание имеет преимущества, а некоторые так не считают. По-вашему, есть ли преимущества при грудном вскармливании? 1=да 0=нет 8= не знаю <input type="text"/> 0→AB9 8→AB9		
Модуль советов о питании		DA
Когда женщина беременна и после рождения ребенка, многие люди дают советы относительно ее питания, кормления грудью и питания ребенка. Некоторым советам мы следуем, а некоторым нет. Я хочу спросить вас именно о совете, полученном вами. не имеет значения следовали ли вы этому совету или нет. Мне просто интересно, какой совет вам дали и кто.		
DA1. Получали ли вы советы по питанию в период вашей беременности? 1=да 0=нет 8= не знаю <input type="text"/> 0→DA4 8→DA4	DA7. Используя шкалу, оцените насколько важны советы по грудному вскармливанию, которые мы получаем от врача, медсестры, акушера или фельдшера? <input type="text"/> <i>(покажите шкалу и отметьте цифру, соответствующую ответу)</i>	
DA2. Давали ли вам врач, медсестра, акушер или фельдшер советы по питанию? 1= да 0 = нет <input type="text"/>	DA8. Давали ли вам советы по грудному вскармливанию члены семьи, друзья или соседи? 1=да 0=нет <input type="text"/>	
DA3. Давал ли вам член семьи, друг или сосед советы по питанию? 1= да 0 = нет <input type="text"/>	DA9. Как долго (в месяцах) советовали вам члены семьи, друзья или соседи кормить грудью без кормления другими жидкостями или сухим питанием? <input type="text"/> М <input type="text"/> М <i>(Поставьте 00, если &lt; 1 мес; 88 -если не знаете/ не помните; 99 - если не давали советы по длительности или не указывали длительность)</i>	
DA4. Давали ли вам врач, медсестра, акушер или фельдшер советы по грудному вскармливанию? 1=да 0 = нет 8= не знаю <input type="text"/> 0→DA8 8→DA8	DA10. В каком возрасте (в месяцах) советовали вам члены семьи, друзья, соседи прекратить грудное вскармливание? <input type="text"/> М <input type="text"/> М <i>(Поставьте 00, если &lt; 1 мес; 88 -если не знаете/ не помните; 99 - если не давали советы по длительности или не указывали длительность)</i>	
DA5. Как долго (в месяцах) советовали вам врач, медсестра, акушер или фельдшер кормить грудью, не давая другие жидкости или густую пищу? <input type="text"/> М <input type="text"/> М <i>(Поставьте 00, если &lt; 1 мес; 88 - если не знаете/не помните; 99 - если не давали советы по длительности или не указывали длительность)</i>	DA11. Используя шкалу, оцените насколько важны советы по грудному вскармливанию, которые мы получаем от семьи, друзей или соседей? <input type="text"/> <i>(покажите шкалу и отметьте цифру, соответствующую ответу)</i>	
DA6. В каком возрасте (в месяцах) советовали вам врач, медсестра, акушер или фельдшер прекратить грудное вскармливание? <input type="text"/> М <input type="text"/> М <i>(Поставьте 00 если &lt;1мес; 88 - если не знает помните; 99 - если не давали советы по длительности или не указывали длительность)</i>	<b>Интервьюер! Задайте вопрос, если хотя бы один из ответов DA4, DA8=1</b>	
	DA12. Используя шкалу, оцените, насколько полученные советы повлияли на ваши собственные решения в отношении	

	грудного вскармливания. <input type="text"/> (покажите шкалу и отметьте цифру, соответствующую ответу)								
<b>Модуль витаминов/добавок</b>	<b>VS</b>								
Я теперь задам вам вопросы о витаминах и добавках, которые могли принимать вы и ваш ребенок. Некоторые люди принимают эти добавки, а некоторые не принимают их, и это нормально. Я начну с добавок, которые вы, возможно, могли принимать.									
<b>VS1.</b> Во время вашей последней беременности принимали ли вы подобную добавку фолиевой кислоты? (Покажите диспенсер) <input type="text"/> 1=да 0 = нет 8= не знаю	<b>VS7.</b> Как давно (в месяцах) (имя ребенка) в последний раз принимал капсулу с витамином А? <input type="text"/> (поставьте 00, если < 1 мес; 88 - если не знаете/не помните) М М								
<b>VS2.</b> Во время вашей последней беременности принимали ли вы подобную добавку железа? (Покажите диспенсер) <input type="text"/> 1=да 0 = нет 8= не знаю	<b>VS8.</b> Говорил ли вам врач или медсестра, что у (имя ребенка) анемия? 1=да 0 = нет 8= не знаю <b>0→VS10</b> <b>8→VS10</b>								
<b>VS3.</b> Принимали ли вы подобную дозу витамина А в первые два месяца после рождения самого младшего ребенка? (Покажите капсулу витамина А) <input type="text"/> 1=да 0 = нет 8= не знаю	<b>VS9.</b> (имя ребенка) принимал железосодержащий сироп или таблетки для лечения анемии? <input type="text"/> 1=да 0 = нет 8= не знаю								
<b>VS4.</b> Говорил ли вам врач или медсестра о том, что у вас анемия? <input type="text"/> 1=да 0 = нет 8= не знаю <b>0→VS6</b> <b>8→VS6</b>	<b>VS10.</b> Давали ли вы, или кто-то другой, когда-либо (имя ребенка) что-нибудь из этих витаминно-минеральных добавок? (прочитайте список и отметьте каждый ответ) <table border="1"> <tr> <td>a Витамин Д</td> <td>a <input type="text"/></td> </tr> <tr> <td>b Рыбий жир</td> <td>b <input type="text"/></td> </tr> <tr> <td>c Мультивитамины</td> <td>c <input type="text"/></td> </tr> <tr> <td>d Другое (укажите)</td> <td>d <input type="text"/></td> </tr> </table> 1=да 0 = нет 8= не знаю	a Витамин Д	a <input type="text"/>	b Рыбий жир	b <input type="text"/>	c Мультивитамины	c <input type="text"/>	d Другое (укажите)	d <input type="text"/>
a Витамин Д	a <input type="text"/>								
b Рыбий жир	b <input type="text"/>								
c Мультивитамины	c <input type="text"/>								
d Другое (укажите)	d <input type="text"/>								
<b>VS5.</b> Принимали ли вы железосодержащие капсулы или сироп для лечения анемии у вас? <input type="text"/> 1=да 0 = нет 8= не знаю	<b>VS11.</b> Вы когда-нибудь видели подобную упаковку Спринклз? Покажите пакет Спринклз <input type="text"/> 1=да 0 = нет 8= не знаю <b>0→FF1</b> <b>8→FF1</b>								
Теперь я хочу задать несколько вопросов насчет витаминов, минералов и добавок, которые, возможно, мог принимать (имя ребенка). Это нормально, если (имя ребенка) не принимал этих добавок.	<b>VS12.</b> Вы когда-нибудь получали подобную упаковку Спринклз? Покажите пакет Спринклз <input type="text"/> 1=да 0 = нет 8= не знаю <b>0→FF1</b> <b>8→FF1</b>								
<b>VS6.</b> Принимал ли (имя ребенка) подобную капсулу витамина А? <input type="text"/> Покажите 100000МЕ для детей 6-11 месяцев <b>0→VS8</b> Покажите 200000МЕ для детей 12-59 мес. <b>8→VS8</b> 1=да 0 = нет 8= не знаю	<b>VS13.</b> (имя ребенка) когда-нибудь употреблял Спринклз? <input type="text"/> 1=да 0 = нет 8= не знаю								



Модуль обогащенной муки		FF																																																					
Сейчас я задам вам вопрос о муке, которую вы используете для выпечки хлеба или тортов, либо других блюд в вашем доме.																																																							
FF1. Сколько муки потребляет ваша семья за один месяц (в кг)?  <div style="display: flex; justify-content: space-around; width: 100px;"> <input style="width: 20px; height: 20px; border: 1px solid black;" type="text"/> <input style="width: 20px; height: 20px; border: 1px solid black;" type="text"/> <input style="width: 20px; height: 20px; border: 1px solid black;" type="text"/> </div> <p style="text-align: center; font-size: small;">Если семья не потребляет муку, перейдите к FF6</p>	FF6. Вы слышали когда-нибудь о муке, обогащенной витаминно-минеральными добавками?  <div style="display: flex; justify-content: space-between; width: 100px;"> <input style="width: 20px; height: 20px; border: 1px solid black;" type="text"/> <span>0→VHC1</span> </div> <p style="text-align: center; font-size: small;">1=да 0=нет</p>																																																						
FF2. Какой сорт муки вы обычно используете в приготовлении пищи? <table style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 60%;">Высшего сорта</td> <td style="width: 10%; text-align: center;">1</td> <td style="width: 30%; border: 1px solid black;">1→FF4</td> </tr> <tr> <td>Первого сорта</td> <td style="text-align: center;">2</td> <td style="border: 1px solid black;">2→FF4</td> </tr> <tr> <td>Второго сорта</td> <td style="text-align: center;">3</td> <td style="border: 1px solid black;">3→FF4</td> </tr> <tr> <td>Муку, перемолотую из собственного зерна</td> <td style="text-align: center;">4</td> <td style="border: 1px solid black;">4→FF3</td> </tr> </table>	Высшего сорта	1	1→FF4	Первого сорта	2	2→FF4	Второго сорта	3	3→FF4	Муку, перемолотую из собственного зерна	4	4→FF3	FF7. Есть ли, по-вашему, преимущества использования обогащенной муки?  <div style="display: flex; justify-content: space-between; width: 100px;"> <span>1=да 0=нет 8=не знаю</span> <input style="width: 20px; height: 20px; border: 1px solid black;" type="text"/> </div> <div style="display: flex; justify-content: space-between; width: 100px; font-size: small;"> <span>0→VHC1</span> <span>8→VHC1</span> </div>																																										
Высшего сорта	1	1→FF4																																																					
Первого сорта	2	2→FF4																																																					
Второго сорта	3	3→FF4																																																					
Муку, перемолотую из собственного зерна	4	4→FF3																																																					
FF3. С какой мукой вы смешиваете муку, перемолотую из собственного зерна?  <table style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 60%;">Не смешиваю перемолотую муку с другой мукой</td> <td style="width: 10%; text-align: center;">0</td> <td style="width: 30%; border: 1px solid black;">0→FF6</td> </tr> <tr> <td>Высшего сорта</td> <td style="text-align: center;">1</td> <td></td> </tr> <tr> <td>Первого сорта</td> <td style="text-align: center;">2</td> <td></td> </tr> <tr> <td>Второго сорта</td> <td style="text-align: center;">3</td> <td></td> </tr> <tr> <td>Не знаю</td> <td style="text-align: center;">8</td> <td></td> </tr> </table>	Не смешиваю перемолотую муку с другой мукой	0	0→FF6	Высшего сорта	1		Первого сорта	2		Второго сорта	3		Не знаю	8		FF8. По вашему мнению, каковы преимущества обогащенной муки? <i>(Не читайте. Отметьте упомянутые пункты цифрой 1, неупомянутые - 0)</i> <table style="width: 100%; border-collapse: collapse;"> <tr><td>a Мука имеет более высокое качество</td><td style="width: 20px;"></td><td>a</td></tr> <tr><td>b Вкус муки лучше</td><td></td><td>b</td></tr> <tr><td>c Дети больше любят муку</td><td></td><td>c</td></tr> <tr><td>d Дети растут лучше</td><td></td><td>d</td></tr> <tr><td>e Содержит витамины/минералы</td><td></td><td>e</td></tr> <tr><td>f Дети становятся умнее</td><td></td><td>f</td></tr> <tr><td>g Дети становятся крепче</td><td></td><td>g</td></tr> <tr><td>h Оздоровляет детей</td><td></td><td>h</td></tr> <tr><td>i Оздоровляет мужчин/женщин</td><td></td><td>i</td></tr> <tr><td>j Предупреждает анемию</td><td></td><td>j</td></tr> <tr><td>k Предупреждает болезни</td><td></td><td>k</td></tr> <tr><td>l Влияет отрицательно</td><td></td><td>l</td></tr> <tr><td>m другое (укажите)</td><td></td><td>m</td></tr> </table>	a Мука имеет более высокое качество		a	b Вкус муки лучше		b	c Дети больше любят муку		c	d Дети растут лучше		d	e Содержит витамины/минералы		e	f Дети становятся умнее		f	g Дети становятся крепче		g	h Оздоровляет детей		h	i Оздоровляет мужчин/женщин		i	j Предупреждает анемию		j	k Предупреждает болезни		k	l Влияет отрицательно		l	m другое (укажите)		m
Не смешиваю перемолотую муку с другой мукой	0	0→FF6																																																					
Высшего сорта	1																																																						
Первого сорта	2																																																						
Второго сорта	3																																																						
Не знаю	8																																																						
a Мука имеет более высокое качество		a																																																					
b Вкус муки лучше		b																																																					
c Дети больше любят муку		c																																																					
d Дети растут лучше		d																																																					
e Содержит витамины/минералы		e																																																					
f Дети становятся умнее		f																																																					
g Дети становятся крепче		g																																																					
h Оздоровляет детей		h																																																					
i Оздоровляет мужчин/женщин		i																																																					
j Предупреждает анемию		j																																																					
k Предупреждает болезни		k																																																					
l Влияет отрицательно		l																																																					
m другое (укажите)		m																																																					
FF4. Где вы обычно покупаете муку для выпечки? <table style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 60%;">Продовольственный магазин</td> <td style="width: 10%; text-align: center;">1</td> <td style="width: 30%; border: 1px solid black;"><input style="width: 20px; height: 20px;" type="text"/></td> </tr> <tr> <td>Рынок</td> <td style="text-align: center;">2</td> <td></td> </tr> <tr> <td>Мельница</td> <td style="text-align: center;">3</td> <td></td> </tr> <tr> <td>Другое (укажите)</td> <td style="text-align: center;">4</td> <td></td> </tr> <tr> <td>Не знаю</td> <td style="text-align: center;">8</td> <td></td> </tr> </table>	Продовольственный магазин	1	<input style="width: 20px; height: 20px;" type="text"/>	Рынок	2		Мельница	3		Другое (укажите)	4		Не знаю	8		FF9. Если бы у вас было право выбора из двух буханок хлеба одинакового веса и цены, но одна бы содержала витаминно-минеральные добавки, а другая - нет; какую буханку вы бы купили? <table style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 60%;">Буханка с витаминно-минеральными добавками</td> <td style="width: 10%; text-align: center;">1</td> <td style="width: 30%; border: 1px solid black;"><input style="width: 20px; height: 20px;" type="text"/></td> </tr> <tr> <td>Буханка без витаминно-минеральных добавок</td> <td style="text-align: center;">2</td> <td></td> </tr> <tr> <td>Мне все равно</td> <td style="text-align: center;">3</td> <td></td> </tr> <tr> <td>Не знаю</td> <td style="text-align: center;">8</td> <td></td> </tr> </table>	Буханка с витаминно-минеральными добавками	1	<input style="width: 20px; height: 20px;" type="text"/>	Буханка без витаминно-минеральных добавок	2		Мне все равно	3		Не знаю	8																												
Продовольственный магазин	1	<input style="width: 20px; height: 20px;" type="text"/>																																																					
Рынок	2																																																						
Мельница	3																																																						
Другое (укажите)	4																																																						
Не знаю	8																																																						
Буханка с витаминно-минеральными добавками	1	<input style="width: 20px; height: 20px;" type="text"/>																																																					
Буханка без витаминно-минеральных добавок	2																																																						
Мне все равно	3																																																						
Не знаю	8																																																						
<b>Модуль контактов с СКЗ</b>		<b>VHC</b>																																																					
В некоторых селах Кыргызской Республики существуют Сельские комитеты здоровья. Я хотела вас спросить, что вы слышали об СКЗ. Помните, нет правильных ответов или неправильных ответов, мы просто хотим знать что вы слышали об этом и ваш личный опыт.																																																							
VHC1. Вы когда-нибудь слышали о Сельском комитете здоровья (СКЗ)?  <div style="display: flex; justify-content: space-between; width: 100px;"> <input style="width: 20px; height: 20px; border: 1px solid black;" type="text"/> <span>0→P1</span> </div> <p style="text-align: center; font-size: small;">1=да 0=нет 8=не знаю</p>	VHC4. Разговаривали ли вы с членом СКЗ о вопросах вашего питания во время беременности, или о грудном вскармливании или питании ребенка?  <div style="display: flex; justify-content: space-between; width: 100px;"> <input style="width: 20px; height: 20px; border: 1px solid black;" type="text"/> <span>0→P1</span> </div> <div style="display: flex; justify-content: space-between; width: 100px; font-size: small;"> <span>1=да 0=нет 8=не знаю</span> <span>8→P1</span> </div>																																																						
VHC2. Вы когда-нибудь разговаривали с кем-нибудь из СКЗ о вопросах здоровья?  <div style="display: flex; justify-content: space-between; width: 100px;"> <input style="width: 20px; height: 20px; border: 1px solid black;" type="text"/> <span>0→P1</span> </div> <div style="display: flex; justify-content: space-between; width: 100px; font-size: small;"> <span>1=да 0=нет 8=не знаю</span> <span>8→P1</span> </div>	VHC5. Используя шкалу, насколько полезными, по-вашему, были визиты к членам СКЗ (по вопросам питания, грудного вскармливания или питания ребенка)?  <div style="text-align: center; font-size: small;">(покажите шкалу и отметьте цифру, соответствующую ответу)</div>																																																						
VHC3. Сколько времени прошло с тех пор, как вы в последний раз разговаривали с членом СКЗ о вопросах здоровья? <table style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 60%;">&gt; 1 года назад</td> <td style="width: 10%; text-align: center;">0</td> <td style="width: 30%; border: 1px solid black;"><input style="width: 20px; height: 20px;" type="text"/></td> </tr> <tr> <td>6-12 месяцев назад</td> <td style="text-align: center;">1</td> <td></td> </tr> <tr> <td>3 - 6 месяцев назад</td> <td style="text-align: center;">2</td> <td></td> </tr> <tr> <td>1- 3 месяца назад</td> <td style="text-align: center;">3</td> <td></td> </tr> <tr> <td>&lt; 1 месяца назад</td> <td style="text-align: center;">4</td> <td></td> </tr> </table>	> 1 года назад	0	<input style="width: 20px; height: 20px;" type="text"/>	6-12 месяцев назад	1		3 - 6 месяцев назад	2		1- 3 месяца назад	3		< 1 месяца назад	4		VHC6. Заинтересованы ли вы в получении советов от СКЗ?  <div style="display: flex; justify-content: space-between; width: 100px;"> <input style="width: 20px; height: 20px; border: 1px solid black;" type="text"/> <span>0→P1</span> </div> <div style="display: flex; justify-content: space-between; width: 100px; font-size: small;"> <span>1=да 0=нет 8=не знаю</span> <span>8→P1</span> </div>																																							
> 1 года назад	0	<input style="width: 20px; height: 20px;" type="text"/>																																																					
6-12 месяцев назад	1																																																						
3 - 6 месяцев назад	2																																																						
1- 3 месяца назад	3																																																						
< 1 месяца назад	4																																																						
	VHC7. По каким вопросам вы бы хотели получить совет? <i>(Запишите все упомянутые темы)</i> <div style="border-bottom: 1px solid black; height: 20px; width: 100%;"></div>																																																						
<b>Беременность</b>																																																							
Прежде чем мы продолжим, мне нужно знать, беременны ли вы. Даже если вы не уверены, но думаете, что возможно вы беременны, но не знаете точно, нам бы хотелось знать это.																																																							
P1. Вы беременны сейчас?  <div style="display: flex; justify-content: space-between; width: 100px;"> <input style="width: 20px; height: 20px; border: 1px solid black;" type="text"/> <span>0→AN1</span> </div> <div style="display: flex; justify-content: space-between; width: 100px; font-size: small;"> <span>1=да 0=нет 7=может быть, но не уверена</span> <span>8=не знаю</span> <span>7→AN1</span> <span>8→AN1</span> </div>	P2. Сколько недель составляет ваша беременность на данный момент? <i>(поставьте 88 - если не знаете)</i> <div style="display: flex; justify-content: space-between; width: 100px;"> <input style="width: 20px; height: 20px; border: 1px solid black;" type="text"/> <input style="width: 20px; height: 20px; border: 1px solid black;" type="text"/> </div>																																																						
Если да (1), то не нужно брать кровь или производить антропометрические замеры у матери. Возьмите кровь и замеры только у ребенка																																																							

Антропометрия		AN	
Теперь, я измерю ваш рост и вес и рост и вес вашего ребенка.			
AN1. Были ли взяты антропометрические данные у матери? <input type="checkbox"/>	1 = да 0 = нет	1→AN3	
AN2. Почему нет? <input type="checkbox"/>	1 = отказалась 3 = не присутствовала 2 = беременна 4 прочее (укажите)		
AN3. Рост матери (см) <input type="text"/> . <input type="text"/>		AN5. Были ли взяты антропометрические данные у ребенка? <input type="checkbox"/>	1 = да 0 = нет 1→AN7
AN4. Вес матери (кг) <input type="text"/> . <input type="text"/>		AN6. Почему нет? <input type="checkbox"/>	1 =отказался (плакал, бил ножками и т.д.) 3=не присутствовал 2 = мать/сопровождающий отказались 4 прочее (укажите)
		AN7. Вес ребенка (кг) <input type="text"/> . <input type="text"/>	
		AN8. Рост ребенка (см) <input type="text"/> . <input type="text"/>	
Модуль пробы крови		BS	
Последнее, что мы сделаем сегодня, это возьмем небольшое количество крови с вашего пальца и пальца вашего ребенка. Это может причинить немного дискомфорта, но мы сможем сказать, есть ли у вас или вашего ребенка анемия.			
BS1. Когда вы ели в последний раз? <input type="text"/> : <input type="text"/>	ч ч : м м	BS8. Когда (имя ребенка) в последний раз принимал питание? <input type="text"/> : <input type="text"/>	ч ч : м м
BS2. Была ли получена капиллярная проба от матери? <input type="checkbox"/>	1=да 0 = нет 1→BS4	BS9. Была ли получена капиллярная проба у ребенка? <input type="checkbox"/>	1=да 0 = нет 1→BS11
BS3. Почему нет? <input type="checkbox"/>	1 = отказалась 3=не присутствовала →BS7 2 = беременна 4=технические трудности 5=прочее (укажите)	BS10. Почему нет? <input type="checkbox"/>	1 = отказался (плакал, бил ножками и т.д.) 3=не присутствовал →BS14 2 = мать/сопровожд. отказались 4=технические трудности 5=прочее (укажите)
BS4. Приблизительно сколько микролитров крови было собрано в микротейнере? <input type="text"/>		BS11. Приблизительно сколько микролитров крови было собрано в микротейнере? <input type="text"/>	
BS5. В какое время была взята проба? <input type="text"/> : <input type="text"/>	ч ч : м м	BS12. В какое время была взята проба? <input type="text"/> : <input type="text"/>	ч ч : м м
BS6. Концентрация гемоглобина из Гемокью <input type="text"/> . <input type="text"/> г/длитр <i>(поставьте 88.8 если не измерялось/не знаете)</i>		BS13. Концентрация гемоглобина из Гемокью <input type="text"/> . <input type="text"/> г/длитр <i>(поставьте 88.8 если не измерялось/не знаете)</i>	
BS7. Идентификационная бирка - <b>МАТЬ</b> Поставьте бирку крови матери здесь		BS14. Идентификационная бирка - <b>РЕБЕНОК</b> Поставьте бирку крови ребенка здесь	
<p>Подпись супервайзера по исследованию, подтверждающая, что женщина или ребенок были направлены в медицинское учреждение на лечение, в случае, если гемоглобин &lt;7,0 г/длитр</p> <p>_____</p> <p>Подпись _____ Дата _____</p>			
<p>Не забудьте предоставить матери результаты ее анализа гемоглобина и анализы гемоглобина ее ребенка. Попросите мать расписаться здесь в подтверждение того, что она получила результаты анализа гемоглобина и направление (если необходимо).</p> <p>_____</p> <p>Подпись _____ Дата _____</p>			
<p>Подпись супервайзера по исследованию, подтверждающая, что вопросник проверен и заполнен:</p> <p>_____</p> <p>Подпись _____ Дата _____</p>			
<p>Комментарии интервьюера:</p> <p>_____</p> <p>_____</p>			

## APPENDIX IX COMPARISON OF VARIOUS CHARACTERISTICS AMONG CHILDREN (6-24 MONTHS) — KYRGYZSTAN, 2009 AND 2013

Table IX.1: Anthropometric characteristics by age — Kyrgyzstan, 2009 and 2013

Characteristics	2009		2013		p- value <sup>a</sup>
	N	% (95% CI)	N	% (95% CI)	
<b>Wasted</b> (weight-for-length Z<-2.0)					
	400	2.0 (0.6 - 3.5)	1722	2.4 (1.4 - 3.5)	0.659
6-11 months	131	1.9 (0.0 - 4.1)	520	2.7 (0.8 - 4.5)	0.568
12-24 months	269	2.1 (0.2 - 4)	1202	2.3 (1.1 - 3.5)	0.861
<b>Stunted</b> (length-for-age Z<-2.0)					
	401	14.1 (10.4 - 17.9)	1717	10.6 (8.4 - 12.9)	0.116
6-11 months	131	6.9 (2.8 - 11.2)	515	5.0 (2.3 - 7.6)	0.451
12-24 months	270	17.7 (12.5 - 22.9)	1202	12.9(10.2 - 15.7)	0.109
<b>Underweight</b> (weight-for-age Z<-2.0)					
	403	3.1 (1.1 - 5.2)	1724	4.5 (3.1 - 6)	0.272
6-11 months	132	2.6 (0.0 - 5.4)	520	3.4 (0.9 - 6)	0.671
12-24 months	271	3.4 (0.7 - 6.1)	1204	5 (3.4 - 6.6)	0.315

NOTE: CI=confidence interval; 95% CI's adjusted for survey design.

Anthropometric values based on WHO growth reference curves (WHO, 2006).

<sup>a</sup>Bivariate tests of statistical significance were conducted using the Pearson chi-square test, accounting for the cluster survey design.

**TABLE IX.2  
PREVALENCE OF ANEMIA, IRON DEFICIENCY, AND VITAMIN A DEFICIENCY BY AGE (6-24 MONTHS) — KYRGYZSTAN, 2009 AND 2013**

Characteristics	All Participants				Participants without Inflammation <sup>a</sup>				
	2009		2013		2009		2013		p
	N	% (95% DI)	N	% (95% CI)	N	% (95% CI)	N	% (95% CI)	
Anemia <sup>a</sup>	450	40.1 (33.1 - 47.0)	1 720	34.5 (30.9 - 38.1)	342	38.6 (31.5 - 45.8)	1 147	31.4 (27.3 - 35.4)	<b>0.085</b>
6-11 months	137	40.7 (29.9 - 51.5)	519	35.6 (27.8 - 43.4)	112	35 (25 - 45.1)	354	33.6 (25.4 - 41.7)	<b>0.831</b>
12-24 months	313	39.8 (32.6 - 46.9)	1 201	34 (30.7 - 37.3)	230	40.4 (32.4 - 48.4)	793	30.4 (26.1 - 34.7)	<b>0.031</b>
Low ferritin (<12 µg/L)	450	50.3 (44.7 - 56)	1 720	33.6 (30.8 - 36.4)	342	56.7 (50.6 - 62.8)	1 147	39.8 (36.4 - 43.2)	<b>0.000</b>
6-11 months	137	44.9 (35.1 - 54.7)	519	29.9 (23.7 - 36.1)	112	48.7 (37.9 - 59.6)	354	34.2 (25.5 - 42.9)	<b>0.041</b>
12-24 months	313	52.8 (46.5 - 59.2)	1 201	35.1 (31.4 - 38.7)	230	60.6 (53.7 - 67.6)	793	42.2 (38.2 - 46.1)	<b>0.000</b>
High sTfR (>8.3 mg/L)	450	51.6 (45 - 58.1)	1 720	39.9 (36.1 - 43.7)	342	50.7 (43.6 - 57.8)	1 147	39.1 (34.6 - 43.5)	<b>0.007</b>
6-11 months	137	53.2 (44.3 - 62.1)	519	38.5 (33 - 44.1)	112	50.4 (41.2 - 59.7)	354	37.7 (30.8 - 44.6)	<b>0.031</b>
12-24 months	313	50.8 (43.6 - 58)	1 201	40.4 (35.6 - 45.2)	230	50.9 (42.5 - 59.3)	793	39.7 (34.1 - 45.2)	<b>0.030</b>
Iron deficiency anemia <sup>a</sup>	450	33.0 (27.2 - 38.9)	1 720	24.7 (21.1 - 28.3)	342	32.9 (26.5 - 39.3)	1 147	23.5 (19.7 - 27.4)	<b>0.014</b>
6-11 months	137	32.4 (22.8 - 41.9)	519	24.1 (16.9 - 31.4)	112	28.5 (19.4 - 37.5)	354	22.3 (14.3 - 30.3)	<b>0.312</b>
12-24 months	313	33.3 (27.3 - 39.3)	1 201	25.0 (21.6 - 28.3)	230	35.1 (27.8 - 42.4)	793	24.0 (19.7 - 28.4)	<b>0.011</b>
Vitamin A deficiency <sup>e</sup>	450	7.3 (5 - 9.6)	1 716	15.4 (12.9 - 17.9)	342	4 (2.1 - 6)	1 147	7.4 (5 - 9.7)	<b>0.029</b>
6-11 months	137	9.2 (4.8 - 13.6)	517	16.0 (11.1 - 20.9)	112	5.3 (1.3 - 9.2)	354	9.5 (5.3 - 13.8)	<b>0.155</b>
12-24 months	313	6.4 (3.8 - 9.1)	1 199	15.2 (12.3 - 18.1)	230	3.4 (1.1 - 5.7)	793	6.4 (3.8 - 9.1)	<b>0.093</b>

NOTE: CI=confidence interval; 95% CI's adjusted for survey design.

<sup>a</sup> Inflammation not present (low C-reactive protein [CRP≤5 mg/L] and low α1-glycoprotein acid [AGP≤1.0 g/L]).

<sup>b</sup> Anemia: hemoglobin < 11.0 g/dL adjusted for altitude.

<sup>c</sup> Iron deficiency anemia: Hemoglobin < 11.0 g/dL and low plasma ferritin (<12 µg/L) or high sTfR (>8.3 mg/L).

<sup>d</sup> Bivariate tests of statistical significance were conducted using the Pearson chi-square test, accounting for the cluster survey design.

<sup>e</sup> Vitamin A deficiency, defined as RBP < 0.71 µmol/L.