



Children and AIDS

Country fact sheets



Children and AIDS: Country Fact Sheets

*Companion to the
Second Stocktaking Report*

UNAIDS, the Joint United Nations Programme on HIV/AIDS, brings together the efforts and resources of ten UN system organizations to the global AIDS response. Cosponsors include UNHCR, UNICEF, WFP, UNDP, UNFPA, UNODC, ILO, UNESCO, WHO and the World Bank. Based in Geneva, the UNAIDS secretariat works on the ground in more than 75 countries worldwide.



CONTENTS

Page iii	Note on the estimates
-----------------	---------------------------------------

Page iv	Definitions and methodology
----------------	---

Page v	Notes on country data sources
---------------	---

Page 1	Country fact sheets
---------------	-------------------------------------

Page 1	Afghanistan
Page 2	Albania
Page 3	Algeria
Page 4	Angola
Page 5	Antigua and Barbuda
Page 6	Argentina
Page 7	Armenia
Page 8	Azerbaijan
Page 9	Bahamas
Page 10	Bahrain
Page 11	Bangladesh
Page 12	Barbados
Page 13	Belarus
Page 14	Belize
Page 15	Benin
Page 16	Bhutan
Page 17	Bolivia
Page 18	Bosnia and Herzegovina
Page 19	Botswana
Page 20	Brazil
Page 21	Brunei Darussalam
Page 22	Bulgaria
Page 23	Burkina Faso
Page 24	Burundi
Page 25	Cambodia
Page 26	Cameroon
Page 27	Cape Verde
Page 28	Central African Republic
Page 29	Chad
Page 30	Chile
Page 31	China
Page 32	Colombia
Page 33	Comoros
Page 34	Congo
Page 35	Congo, Democratic Republic of the
Page 36	Cook Islands
Page 37	Costa Rica
Page 38	Côte d'Ivoire
Page 39	Croatia
Page 40	Cuba
Page 41	Djibouti
Page 42	Dominica
Page 43	Dominican Republic
Page 44	Ecuador
Page 45	Egypt
Page 46	El Salvador

Page 47	Equatorial Guinea	Page 102	Niue
Page 48	Eritrea	Page 103	Occupied Palestinian Territory
Page 49	Ethiopia	Page 104	Oman
Page 50	Fiji	Page 105	Pakistan
Page 51	Gabon	Page 106	Palau
Page 52	Gambia	Page 107	Panama
Page 53	Georgia	Page 108	Papua New Guinea
Page 54	Ghana	Page 109	Paraguay
Page 55	Grenada	Page 110	Peru
Page 56	Guatemala	Page 111	Philippines
Page 57	Guinea	Page 112	Qatar
Page 58	Guinea-Bissau	Page 113	Romania
Page 59	Guyana	Page 114	Russian Federation
Page 60	Haiti	Page 115	Rwanda
Page 61	Honduras	Page 116	Saint Kitts and Nevis
Page 62	India	Page 117	Saint Lucia
Page 63	Indonesia	Page 118	Saint Vincent and the Grenadines
Page 64	Iran (Islamic Republic of)	Page 119	Samoa
Page 65	Iraq	Page 120	Sao Tome and Principe
Page 66	Jamaica	Page 121	Saudi Arabia
Page 67	Jordan	Page 122	Senegal
Page 68	Kazakhstan	Page 123	Serbia
Page 69	Kenya	Page 124	Seychelles
Page 70	Kiribati	Page 125	Sierra Leone
Page 71	Korea, Democratic People's Republic of	Page 126	Singapore
Page 72	Korea, Republic of	Page 127	Solomon Islands
Page 73	Kuwait	Page 128	Somalia
Page 74	Kyrgyzstan	Page 129	South Africa
Page 75	Lao People's Democratic Republic	Page 130	Sri Lanka
Page 76	Lebanon	Page 131	Sudan
Page 77	Lesotho	Page 132	Suriname
Page 78	Liberia	Page 133	Swaziland
Page 79	Libyan Arab Jamahiriya	Page 134	Syrian Arab Republic
Page 80	Madagascar	Page 135	Tajikistan
Page 81	Malawi	Page 136	Tanzania, United Republic of
Page 82	Malaysia	Page 137	Thailand
Page 83	Maldives	Page 138	The former Yugoslav Republic of Macedonia
Page 84	Mali	Page 139	Timor-Leste
Page 85	Marshall Islands	Page 140	Togo
Page 86	Mauritania	Page 141	Tonga
Page 87	Mauritius	Page 142	Trinidad and Tobago
Page 88	Mexico	Page 143	Tunisia
Page 89	Micronesia (Federated States of)	Page 144	Turkey
Page 90	Moldova	Page 145	Turkmenistan
Page 91	Mongolia	Page 146	Tuvalu
Page 92	Montenegro	Page 147	Uganda
Page 93	Morocco	Page 148	Ukraine
Page 94	Mozambique	Page 149	United Arab Emirates
Page 95	Myanmar	Page 150	Uruguay
Page 96	Namibia	Page 151	Uzbekistan
Page 97	Nauru	Page 152	Vanuatu
Page 98	Nepal	Page 153	Venezuela (Bolivarian Republic of)
Page 99	Nicaragua	Page 154	Viet Nam
Page 100	Niger	Page 155	Yemen
Page 101	Nigeria	Page 156	Zambia
		Page 157	Zimbabwe

CHILDREN AND AIDS: COUNTRY FACT SHEETS

The 'Children and AIDS Fact Sheets' in this publication reflect the situation of children affected by AIDS in low- and middle-income countries and territories. They capture the minimum data needed to monitor progress associated with *Unite for Children, Unite against AIDS* in relation to the 'Four Ps': Prevent mother-to-child transmission of HIV; Provide paediatric treatment; Prevent infection among adolescents and young people; and Protect and support children affected by HIV/AIDS. The fact sheets contain statistics currently available in UNICEF global databases on key indicators for children and AIDS. These databases contain only statistically sound and nationally representative data from household surveys, including Multiple Indicator Cluster Surveys (MICS) and Demographic and Health Surveys (DHS).

The statistics in the fact sheets include estimates and indicators:

- The *estimates* are generated from statistical models and are used to establish the magnitude of the problem. The estimates are included in the fact sheets for informational purposes.
- The *indicators* are collected at the country level through population-based surveys or programme data. They provide data regarding a country's response to a particular issue associated with the epidemic and are used to monitor changes over time. The indicators included in the fact sheets are the primary ones required for the United Nations General Assembly Special Session (UNGASS) on HIV and AIDS reporting or other global commitments.

Where trend data exist, graphs are included in the fact sheets to show changes in coverage or service delivery over time.

NOTE ON THE ESTIMATES

In 2007, UNAIDS and WHO released new global HIV and AIDS estimates, which were derived from a more refined estimation methodology and reflect the availability of more reliable data from population-based surveys and expanded national sentinel surveillance systems in a number of countries. As a result, estimates of the number of HIV-infected pregnant women needing interventions for the prevention of mother-to-child transmission (PMTCT) have also been refined, and coverage rates reported for 2004 and 2005 have thus been recalculated using the newly generated estimates. The new UNAIDS estimates for 2007 are lower than estimates used to report on PMTCT coverage in previous reports; differences between previous estimates and the newly generated estimates are not related to trends over time and therefore are not comparable. More details on the HIV estimates methodology can be found at <www.unaids.org>.

The following estimates are still under review and were not available at the time of publication of the country fact sheets:

- Estimated number of children living with HIV
- Percentage of infants born to HIV-positive women receiving cotrimoxazole prophylaxis within 2 months of birth
- Percentage of infants born to HIV-positive women receiving a virological test for HIV diagnosis within 2 months of birth
- Estimated number of children in need of antiretroviral treatment (ART)
- Percentage of children in need receiving ART
- Estimated number of children orphaned by all causes
- Estimated number of children orphaned by AIDS
- Estimated number of double orphans by all causes

DEFINITIONS AND METHODOLOGY

General

- **Estimated adult HIV prevalence rate:**
Percentage of adults aged 15–49 living with HIV as of 2005.

Prevent mother-to-child transmission of HIV

- **Antenatal care coverage:** Percentage of women aged 15–49 attended at least once during pregnancy by skilled health personnel, i.e., doctors, nurses or midwives. These data are collected from household surveys, including DHS and MICS.
- **Estimated number of births per year:** Estimated number of live births in 2006.
- **Estimated number of HIV+ pregnant women:** Estimated number of pregnant women living with HIV as of 2006.
- **Number and % of HIV+ pregnant women receiving ARVs for PMTCT:** Number of pregnant women who tested positive for HIV during visits to antenatal clinics and were provided with antiretrovirals (ARVs) to prevent mother-to-child transmission of HIV. The percentage of HIV-infected women who received ARVs for PMTCT is calculated by dividing the reported number of HIV-positive pregnant women who received ARVs for PMTCT by the estimated unrounded number of HIV-positive pregnant women in 2006. Ranges in estimates of coverage of ARVs for PMTCT are based on plausibility, or uncertainty, bounds in the denominator, i.e., low and high estimated numbers of HIV-positive pregnant women. No coverage has been calculated where the number of HIV-positive pregnant women is estimated to be less than 500. For these countries, plausibility bounds in the denominator are very wide, and coverage data are likely to be unreliable. Because the numerator and denominator are derived from different sources and developed utilizing different methods, care should be taken in interpreting changes over time with respect to this indicator. This is an UNGASS indicator.

Provide paediatric HIV treatment

- **Estimated number of children living with HIV:** Estimated number of children aged 0–14 living with HIV.
- **Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth:** These data are collected from national programme records that include public and private providers. The denominator of the percentage is the estimated number of HIV-positive pregnant women as a proxy for infants in need of cotrimoxazole, while the numerator reflects programme data. Because the numerator and denominator are derived from different sources and developed utilizing

different methods, care should be taken in interpreting changes over time with respect to this indicator.

- **Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth:** These data are collected from national programme records including public and private providers. The denominator of the percentage is the estimated number of children born to HIV-positive pregnant women. Assuming the ratio of one child to one mother, the figure of HIV-exposed children is assumed to be equal to the number of HIV-positive pregnant women. In addition, because the denominator is based on an estimate while the numerator is based on programme data, care should be taken in interpreting changes over time with respect to this indicator.
- **Estimated number of children in need of ART:** Estimated number of children aged 0–14 living with HIV in need of antiretroviral treatment (ART).
- **Number and % of children in need receiving ART:** Number of children aged 0–14 living with HIV receiving ART as of 2006. These data are collected from national programme records and should include data from public and private providers. The denominator of the percentage is the estimated number of children in need of ART, while the numerator reflects programme data. Because the numerator and denominator are derived from different sources and developed utilizing different methods, care should be taken in interpreting changes over time with respect to this indicator.

Prevent infection among adolescents and young people

- **Estimated HIV prevalence among young people aged 15–24:** Percentage of young men and women aged 15–24 living with HIV as of 2005. These estimates are generated from statistical models for most countries. More recently, an increasing number of countries that include HIV testing in population-based surveys are reporting from these sources.
- **HIV prevalence among pregnant women aged 15–24 in the capital city:** Percentage of blood samples taken from pregnant women aged 15–24 who test positive for HIV during ‘unlinked anonymous’ sentinel surveillance at selected antenatal clinics. These data are collected in routine HIV sentinel surveillance by the Ministry of Health or a designated coordinating authority.
- **HIV prevalence among most-at-risk populations:** Behavioural surveillance surveys (BSS) of groups with behaviours that put them at high risk of HIV infection collect these data. The national AIDS coordinating body should report these data where relevant. These data are also compiled in UNAIDS’ *Report on the Global AIDS Epidemic* (forthcoming in 2008). This is an UNGASS indicator.
- **Knowledge of HIV prevention among young people aged 15–24:** Percentage of young men and women aged 15–24 who correctly identify the two major ways of preventing the sexual transmission of HIV (using condoms and limiting sex to one faithful, uninfected partner), who reject the two most common local misconceptions about HIV transmission and who know that a healthy-looking person can transmit HIV. This is an UNGASS and MDG indicator and is collected routinely in household surveys.
- **Sex before age 15 among young people aged 15–19:** Percentage of young people aged 15–19 who say they had sex before age 15. This is an UNGASS indicator and is collected routinely in household surveys.
- **Condom use at last higher-risk sex among young people aged 15–24:** Percentage of young men and women aged 15–24 who say they used a condom the last time they had sex with a non-marital, non-cohabiting partner, of those who have had sex with such a partner in the past 12 months. This is an UNGASS and MDG indicator and is routinely collected in household surveys.

- **% of sex workers who used a condom with their most recent client:** Percentage of female and male sex workers reporting the use of a condom with their most recent client. Behavioural surveillance surveys should collect this indicator and the national AIDS coordinating body should report these data where relevant. This is an UNGASS indicator.
- **% of injecting drug users (IDUs) who used sterile equipment at last injection:** Percentage of IDUs reporting the use of sterile injecting equipment the last time they injected. Behavioural surveillance surveys should collect this indicator, and the national AIDS coordinating body should report these data where relevant. This is an UNGASS indicator.
- **% of men who used a condom the last time they had sex with a male partner:** Percentage of men reporting the use of a condom the last time they had sexual intercourse. Data for this indicator are collected from behavioural surveillance surveys (BSS). This is an UNGASS indicator.

Protect and support children affected by HIV and AIDS

- **Estimated number of children orphaned by all causes:** Estimated number of children aged 0–17 who have lost one or both parents to any cause.
- **Estimated number of children orphaned by AIDS:** Estimated number of children aged 0–17 who have lost one or both parents to AIDS.
- **Estimated number of double orphans due to all causes:** Estimated number of children aged 0–17 who have lost both their biological mother and father to any cause.
- **Orphan school attendance ratio:** Percentage of children aged 10–14 who have lost both biological parents and are currently attending school as a percentage of non-orphaned children of the same age, both of whose parents are alive, living with at least one parent and who are attending school. This is an UNGASS and MDG indicator and is collected routinely in household surveys.
- **% of orphans receiving external support:** Percentage of orphaned and vulnerable children whose households received free basic external support in caring for the child. Data are presented only for countries with HIV prevalence of 5 per cent or greater. These data are collected from household surveys.

NOTES ON COUNTRY DATA SOURCES

Number of HIV+ pregnant women receiving ARVs for PMTCT

- **Armenia:** World Health Organization, Sexually Transmitted Infections/HIV/AIDS Programme, *WHO/Europe Survey on HIV/AIDS and Antiretroviral Therapy: 31 December 2006*, WHO Regional Office for Europe, Copenhagen, 2007.
- **Bangladesh:** Data were provided by a non-governmental organization and are not national-level data.
- **Papua New Guinea:** Data were collected only from Port Moresby Hospital; hence, coverage was not calculated.
- **Thailand:** Country report indicates only 80 per cent of data were reported.

Number of children in need receiving ART

- **Albania:** WHO, Sexually Transmitted Infections/HIV/AIDS Programme, *WHO/Europe Survey on HIV/AIDS and Antiretroviral Therapy: 31 December 2006*, WHO Regional Office for Europe, Copenhagen, 2007.
- **Armenia:** WHO, Sexually Transmitted Infections/HIV/AIDS Programme, *WHO/Europe Survey on HIV/AIDS and Antiretroviral Therapy: 31 December 2006*, WHO Regional Office for Europe, Copenhagen, 2007.
- **Bulgaria:** WHO, Sexually Transmitted Infections/HIV/AIDS Programme, *WHO/Europe Survey on HIV/AIDS and Antiretroviral Therapy: 31 December 2006*, WHO Regional Office for Europe, Copenhagen, 2007.
- **Côte d'Ivoire:** Data reported are partial data from Elizabeth Glaser Pediatric AIDS Foundation and ACONDA-VS, an NGO, only and do not reflect national-level data.
- **Croatia:** WHO, Sexually Transmitted Infections/HIV/AIDS Programme, *WHO/Europe Survey on HIV/AIDS and Antiretroviral Therapy: 31 December 2006*, WHO Regional Office for Europe, Copenhagen, 2007.
- **Indonesia:** Expanded Inter-Agency Task Team (IATT) on Prevention of HIV Infection in Pregnant Women, Mothers and their Children, *A Report Card on Prevention of Mother-to-Child Transmission of HIV and Paediatric HIV Care and Treatment in Low- and Middle-Income Countries, Scaling Up Progress from 2004 to 2005*, UNICEF, New York, February 2007.
- **Iran:** World Health Organization, UNAIDS and UNICEF, *Towards Universal Access: Scaling up priority HIV/AIDS interventions in the health sector, Progress report, April 2007*, WHO, Geneva, 2007.
- **Lao People's Democratic Republic:** Expanded Inter-Agency Task Team (IATT) on Prevention of HIV Infection in Pregnant Women, Mothers and their Children, *A Report Card on Prevention of Mother-to-Child Transmission of HIV and Paediatric HIV Care and Treatment in Low- and Middle-Income Countries, Scaling Up Progress from 2004 to 2005*, UNICEF, New York, February 2007.
- **Oman:** WHO, UNAIDS and UNICEF, *Towards Universal Access: Scaling up priority HIV/AIDS interventions in the health sector, Progress report, April 2007*, WHO, Geneva, 2007.
- **Papua New Guinea:** Data reported are from Port Moresby Hospital only and do not reflect national-level data.
- **Russian Federation:** WHO, UNAIDS and UNICEF, *Towards Universal Access: Scaling up priority HIV/AIDS interventions in the health sector, Progress report, April 2007*, WHO, Geneva, 2007.
- **Thailand:** Data are from 66 out of 76 provinces.
- **Venezuela:** Data are reported from WHO Region of the Americas.

Condom use at last higher-risk sex among young people aged 15–19

- **Niger** (2006): Percentage is based on less than 25 unweighted cases.

Orphan school attendance ratio

- **Eritrea** (1995), **Ghana** (1998 and 2003), **Madagascar** (1997 and 2000), **Nicaragua** (2001), **Niger** (1992), **Nigeria** (2003), **Peru** (2000), **Senegal** (1992–1993 and 2000) and the **United Republic of Tanzania** (1999): Proportion of orphans aged 10–14 attending school is based on small denominators (typically 25–49 unweighted cases).

Children and AIDS fact sheet: Afghanistan

Estimated adult HIV prevalence rate <0.1%
 Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	16%
Source: MICS, 2003	
Estimated number of births per year	1,272,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	<100
Source: UNAIDS/WHO unpublished estimates, 2007	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	0 0.0%
Source: MOH, 2006	

PROVIDE PAEDIATRIC HIV TREATMENT






Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	0 -
Source: MOH, 2006	
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	0 -
Source: MOH, 2006	
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	0 -
Source: MOH, 2006	

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: -	F: -
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Albania

Estimated adult HIV prevalence rate -

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage <small>Source: MICS, 2005</small>	97%
Estimated number of births per year <small>Source: UN Population Division, 2006</small>	52,000
Estimated number of HIV+ pregnant women	-
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	- -

PROVIDE PAEDIATRIC HIV TREATMENT



Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	- -
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	- -
Estimated number of children in need of ART	-
Number and % of children in need receiving ART <small>Source: MOH-National AIDS Program, 2006</small>	5 -

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE


Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24 <small>Source: MICS, 2005</small>	M: -	F: 5.6%
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection <small>Source: BSS, 2005</small>	-	46.0%
% of men who used a condom the last time they had sex with a male partner <small>Source: BSS, 2005</small>	-	61.2%

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS


Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
* Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Algeria

Estimated adult HIV prevalence rate

0.1%

Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	89%
Source: MICS, 2006	
Estimated number of births per year	694,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	<500
Source: UNAIDS/WHO unpublished estimates, 2007	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	-
	[<500 - 783]*

PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	-


PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M:	-	F:	-
HIV prevalence among pregnant women aged 15-24 in capital city				-
HIV prevalence among most-at-risk populations	FSW:	-	IDU:	-
Knowledge of HIV prevention among young people aged 15-24	M:	-	F:	13.1%
Source: MICS, 2006				
Sex before age 15 among young people aged 15-19	M:	-	F:	-
Condom use at last higher-risk sex among young people aged 15-24	M:	-	F:	-
% of sex workers who used a condom with their most recent client	M:	-	F:	-
% of IDUs who used sterile equipment at last injection				-
% of men who used a condom the last time they had sex with a male partner				-


PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	0.99
Source: MICS, 2006	
% of orphans receiving external support	-

Number of HIV+ pregnant women receiving ARVs for PMTCT




Number of children in need receiving ART




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.

* Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Angola

Estimated adult HIV prevalence rate 3.7%
 Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	66%
Source: MICS, 2001	
Estimated number of births per year	792,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	14,094 [8,948 - 19,080]*
Source: UNAIDS/WHO unpublished estimates, 2007	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	1,923 13.6% [10.1% - 21.5%]*
Source: MOH, 2006	

PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	361
Source: MOH, 2006	


PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: 0.9%	F: 2.5%
Source: UNAIDS, 2006		
HIV prevalence among pregnant women aged 15-24 in capital city	2.8%	
Source: UNAIDS, 2006		
HIV prevalence among most-at-risk populations	FSW: 33.3%	IDU: -
Source: UNAIDS, 2006		
Knowledge of HIV prevention among young people aged 15-24	M: -	F: -
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	
% of men who used a condom the last time they had sex with a male partner	-	


PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	0.90
Source: MICS, 2000	
% of orphans receiving external support	-


Number of HIV+ pregnant women receiving ARVs for PMTCT




Number of children in need receiving ART




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Antigua and Barbuda

Estimated adult HIV prevalence rate -

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage		100%
<small>Source: PAHO/WHO, 2005</small>		
Estimated number of births per year		0
<small>Source: UN Population Division, 2006</small>		
Estimated number of HIV+ pregnant women		-
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	4	-
<small>Source: MOH, 2006</small>		

PROVIDE PAEDIATRIC HIV TREATMENT






Estimated number of children living with HIV		-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	4	-
<small>Source: MOH, 2006</small>		
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	0	-
<small>Source: MOH, 2006</small>		
Estimated number of children in need of ART		-
Number and % of children in need receiving ART	7	-
<small>Source: MOH, 2006</small>		

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M:	-	F:	-
HIV prevalence among pregnant women aged 15-24 in capital city				-
HIV prevalence among most-at-risk populations	FSW:	-	IDU:	-
Knowledge of HIV prevention among young people aged 15-24	M:	-	F:	-
Sex before age 15 among young people aged 15-19	M:	-	F:	-
Condom use at last higher-risk sex among young people aged 15-24	M:	-	F:	-
% of sex workers who used a condom with their most recent client	M:	-	F:	-
% of IDUs who used sterile equipment at last injection				-
% of men who used a condom the last time they had sex with a male partner				-

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes		-
Estimated number of children orphaned by AIDS		-
Estimated number of double orphans by all causes		-
Orphan school attendance ratio		-
% of orphans receiving external support		-



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Argentina

Estimated adult HIV prevalence rate

0.6%

Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	99%
Source: Encuesta Nacional de Nutrición y Salud, 2005	
Estimated number of births per year	690,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	1,654 [1,153 - 2,255]*
Source: UNAIDS/WHO unpublished estimates, 2007	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	2,120 >95.0% [94.0% - >95.0%]*
Source: MOH, 2006	

PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	2,148 -
Source: MOH, 2006	
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	3,760 -
Source: MOH, 2006	


PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	
HIV prevalence among most-at-risk populations	FSW: 3.4%	IDU: 23.0%
Source: UNAIDS, 2006		
Knowledge of HIV prevention among young people aged 15-24	M: -	F: -
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	
% of men who used a condom the last time they had sex with a male partner	-	


PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-


Number of HIV+ pregnant women receiving ARVs for PMTCT




Number of children in need receiving ART




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.

* Represents the uncertainty range around the estimates

Rev. 7 March 2008

Children and AIDS fact sheet: Armenia

Estimated adult HIV prevalence rate 0.1%
 Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage <small>Source: DHS, 2005</small>	93%
Estimated number of births per year <small>Source: UN Population Division, 2006</small>	36,000
Estimated number of HIV+ pregnant women <small>Source: UNAIDS/WHO unpublished estimates, 2007</small>	<100 [<100]*
Number and % of HIV+ pregnant women receiving ARVs for PMTCT <small>Source: WHO Regional Office for Europe, 2006</small>	5 -

PROVIDE PAEDIATRIC HIV TREATMENT



Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	- -
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	- -
Estimated number of children in need of ART	-
Number and % of children in need receiving ART <small>Source: MOH, 2006</small>	2 -

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE


Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations <small>Source: UNAIDS, 2006</small>	FSW: 0.4%	IDU: 9.3%
Knowledge of HIV prevention among young people aged 15-24 <small>Source: DHS, 2005</small>	M: 15.1%	F: 22.6%
Sex before age 15 among young people aged 15-19 <small>Source: DHS, 2005</small>	M: 2.6%	F: 0.3%
Condom use at last higher-risk sex among young people aged 15-24 <small>Source: DHS, 2005</small>	M: 86.4%	F: -
% of sex workers who used a condom with their most recent client <small>Source: UNAIDS, 2006</small>	M: 100.0%	F: 89.2%
% of IDUs who used sterile equipment at last injection <small>Source: UNAIDS, 2006</small>	-	46.0%
% of men who used a condom the last time they had sex with a male partner <small>Source: UNAIDS, 2006</small>	-	30.4%

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS


Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Azerbaijan

Estimated adult HIV prevalence rate

0.1%

Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	70%
Source: RHS, 2001	
Estimated number of births per year	129,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	<100
Source: UNAIDS/WHO unpublished estimates, 2007	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	1 -
Source: MOH, 2006	

PROVIDE PAEDIATRIC HIV TREATMENT



Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	0
Source: MOH, 2006	
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	0 -
Source: MOH, 2006	
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	0 -
Source: MOH, 2006	

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE


Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: -	F: 2.0%
Source: MICS, 2000		
Sex before age 15 among young people aged 15-19	M: -	F: 1.0%
Source: RCHS, 2001		
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS


Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.

* Represents the uncertainty range around the estimates

Rev. 7 March 2008

Children and AIDS fact sheet: Bahamas

Estimated adult HIV prevalence rate

3.3%

Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	-	-
Estimated number of births per year	6,000	
<small>Source: UN Population Division, 2006</small>		
Estimated number of HIV+ pregnant women	<100	[<100 - <200]*
<small>Source: UNAIDS/WHO unpublished estimates, 2007</small>		
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	-	-

PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-	-
Estimated number of children in need of ART	-	-
Number and % of children in need receiving ART	-	-

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: -	F: -
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-


PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-


Number of HIV+ pregnant women receiving ARVs for PMTCT




Number of children in need receiving ART




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.

* Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Bahrain

Estimated adult HIV prevalence rate -

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	-
Estimated number of births per year	13,000
<small>Source: UN Population Division, 2006</small>	
Estimated number of HIV+ pregnant women	-
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	-

PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	-


PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M:	-	F:	-
HIV prevalence among pregnant women aged 15-24 in capital city				-
HIV prevalence among most-at-risk populations	FSW:	-	IDU:	-
Knowledge of HIV prevention among young people aged 15-24	M:	-	F:	-
Sex before age 15 among young people aged 15-19	M:	-	F:	-
Condom use at last higher-risk sex among young people aged 15-24	M:	-	F:	-
% of sex workers who used a condom with their most recent client	M:	-	F:	-
% of IDUs who used sterile equipment at last injection				-
% of men who used a condom the last time they had sex with a male partner				-


PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-


Number of HIV+ pregnant women receiving ARVs for PMTCT




Number of children in need receiving ART




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Bangladesh

Estimated adult HIV prevalence rate

<0.1%

Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	48%
Source: MICS, 2006	
Estimated number of births per year	4,013,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	<200 [<100 - <500]*
Source: UNAIDS/WHO unpublished estimates, 2007	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	5 -
Source: MOH, 2006	

PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	5 -
Source: MOH, 2006	
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	0 -
Source: MOH, 2006	
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	1 -
Source: MOH, 2006	


PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	
HIV prevalence among most-at-risk populations	FSW: 0.2%	IDU: 4.9%
Source: UNAIDS, 2006		
Knowledge of HIV prevention among young people aged 15-24	M: -	F: 15.8%
Source: MICS, 2006		
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: 44.1%	F: 31.8%
Source: UNAIDS, 2006		
% of IDUs who used sterile equipment at last injection	8.3%	
Source: UNAIDS, 2006		
% of men who used a condom the last time they had sex with a male partner	49.2%	
Source: UNAIDS, 2006		


PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	0.84
Source: MICS, 2006	
% of orphans receiving external support	-

Number of HIV+ pregnant women receiving ARVs for PMTCT




Number of children in need receiving ART




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.

* Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Barbados

Estimated adult HIV prevalence rate 1.5%
 Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage <small>Source: PAHO/WHO, 2004</small>	100%
Estimated number of births per year <small>Source: UN Population Division, 2006</small>	3,000
Estimated number of HIV+ pregnant women <small>Source: UNAIDS/WHO unpublished estimates, 2007</small>	<100 [<100]*
Number and % of HIV+ pregnant women receiving ARVs for PMTCT <small>Source: MOH, 2006</small>	37 -

PROVIDE PAEDIATRIC HIV TREATMENT






Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	- -
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	- -
Estimated number of children in need of ART	-
Number and % of children in need receiving ART <small>Source: MOH, 2006</small>	17 -

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: -	F: -
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Belarus

Estimated adult HIV prevalence rate 0.3%
 Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage Source: MICS, 2005	99%
Estimated number of births per year Source: UN Population Division, 2006	91,000
Estimated number of HIV+ pregnant women Source: UNAIDS/WHO unpublished estimates, 2007	<100 [<100 - <200]*
Number and % of HIV+ pregnant women receiving ARVs for PMTCT Source: MOH, 2006	158 -

PROVIDE PAEDIATRIC HIV TREATMENT



Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	- -
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth Source: MOH, 2006	114 -
Estimated number of children in need of ART	-
Number and % of children in need receiving ART Source: MOH, 2006	57 -

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE


Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations Source: UNAIDS, 2006	FSW: -	IDU: 30.9%
Knowledge of HIV prevention among young people aged 15-24 Source: MICS, 2005	M: -	F: 33.5%
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client Source: UNAIDS, 2006	M: 100.0%	F: 77.3%
% of IDUs who used sterile equipment at last injection Source: UNAIDS, 2006	-	50.7%
% of men who used a condom the last time they had sex with a male partner Source: UNAIDS, 2006	-	61.6%

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS


Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio Source: MICS, 2005	1.01
% of orphans receiving external support	-




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Belize

Estimated adult HIV prevalence rate

2.5%

Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	-	-
Estimated number of births per year	7,000	
<small>Source: UN Population Division, 2006</small>		
Estimated number of HIV+ pregnant women	<200	[<100 - <500]*
<small>Source: UNAIDS/WHO unpublished estimates, 2007</small>		
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	57	-
<small>Source: MOH, 2006</small>		

PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-	-
Estimated number of children in need of ART	-	-
Number and % of children in need receiving ART	53	-
<small>Source: MOH, 2006</small>		


PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M:	-	F:	-
HIV prevalence among pregnant women aged 15-24 in capital city				-
HIV prevalence among most-at-risk populations	FSW:	-	IDU:	-
Knowledge of HIV prevention among young people aged 15-24	M:	-	F:	-
Sex before age 15 among young people aged 15-19	M:	-	F:	-
Condom use at last higher-risk sex among young people aged 15-24	M:	-	F:	-
% of sex workers who used a condom with their most recent client	M:	-	F:	-
% of IDUs who used sterile equipment at last injection				-
% of men who used a condom the last time they had sex with a male partner				-


PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-


Number of HIV+ pregnant women receiving ARVs for PMTCT




Number of children in need receiving ART




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.

* Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Benin

Estimated adult HIV prevalence rate 1.2%
 Source: DHS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	88%
Source: DHS, 2006	
Estimated number of births per year	358,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	4,497 [3,882 - 5,278]*
Source: UNAIDS/WHO unpublished estimates, 2007	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	2,378 52.9% [45.1% - 61.3%]*
Source: MOH, 2006	

PROVIDE PAEDIATRIC HIV TREATMENT






Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	1,111
Source: MOH, 2006	
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	330
Source: MOH, 2006	

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: 0.4%	F: 1.1%
Source: UNAIDS, 2006		
HIV prevalence among pregnant women aged 15-24 in capital city	1.7% - 2.1%	
Source: SOWC, 2003		
HIV prevalence among most-at-risk populations	FSW: 60.5%	IDU: -
Source: UNAIDS, 2006		
Knowledge of HIV prevention among young people aged 15-24	M: 34.8%	F: 15.9%
Source: DHS, 2006		
Sex before age 15 among young people aged 15-19	M: 13.1%	F: 12.3%
Source: DHS, 2006		
Condom use at last higher-risk sex among young people aged 15-24	M: 44.4%	F: 16.5%
Source: DHS, 2006		
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	
% of men who used a condom the last time they had sex with a male partner	-	

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	0.90
Source: DHS, 2006	
% of orphans receiving external support	-



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Bhutan

Estimated adult HIV prevalence rate

<0.1%

Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	51%
Source: NHS, 2000	
Estimated number of births per year	12,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	<100 [<100]*
Source: UNAIDS/WHO unpublished estimates, 2007	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	3 -
Source: MOH, 2006	

PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	0 -
Source: MOH, 2006	
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	0 -
Source: MOH, 2006	
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	0 -
Source: MOH, 2006	


PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: -	F: -
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-


PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-


Number of HIV+ pregnant women receiving ARVs for PMTCT




Number of children in need receiving ART




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.

* Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Bolivia

Estimated adult HIV prevalence rate

0.1%

Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	79%
<small>Source: DHS, 2003</small>	
Estimated number of births per year	264,000
<small>Source: UN Population Division, 2006</small>	
Estimated number of HIV+ pregnant women	<200
<small>Source: UNAIDS/WHO unpublished estimates, 2007</small>	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	- -
<small>[<200 - <500]*</small>	

PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	- -
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	- -
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	- -


PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: 18.0%	F: 15.0%
<small>Source: DHS, 2003</small>		
Sex before age 15 among young people aged 15-19	M: 15.3%	F: 6.2%
<small>Source: DHS, 2003</small>		
Condom use at last higher-risk sex among young people aged 15-24	M: 37.0%	F: 20.0%
<small>Source: DHS, 2003</small>		
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	
% of men who used a condom the last time they had sex with a male partner	-	


PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	0.74
<small>Source: DHS, 2003</small>	
% of orphans receiving external support	-


Number of HIV+ pregnant women receiving ARVs for PMTCT




Number of children in need receiving ART




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.

* Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Bosnia and Herzegovina

Estimated adult HIV prevalence rate

<0.1%

Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	99%
Source: MICS, 2006	
Estimated number of births per year	35,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	-
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	0 -
Source: MOH, 2006	

PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	0 -
Source: MOH, 2006	
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	0 -
Source: MOH, 2006	
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	0 -
Source: MOH, 2006	


PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: -	F: 48.0%
Source: MICS, 2006		
Sex before age 15 among young people aged 15-19	M: -	F: 0.6%
Source: MICS, 2006		
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: 71.0%
Source: MICS, 2006		
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-


PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	1.01
Source: MICS, 2006	
% of orphans receiving external support	-


Number of HIV+ pregnant women receiving ARVs for PMTCT




Number of children in need receiving ART




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.

* Represents the uncertainty range around the estimates

Rev. 7 March 2008

Children and AIDS fact sheet: Botswana

Estimated adult HIV prevalence rate 25.2%
 Source: UNAIDS/WHO, 2007

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	97%
Source: MICS, 2000	
Estimated number of births per year	47,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	11,370 [10,526 - 12,286]*
Source: UNAIDS/WHO unpublished estimates, 2007	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	12,994 >95.0% [>95.0%]*
Source: MOH, 2006	

PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	8,830 -
Source: MOH, 2006	
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	- -
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	6,831 -
Source: MOH, 2006	


PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: 5.7%	F: 15.3%
Source: UNAIDS, 2006		
HIV prevalence among pregnant women aged 15-24 in capital city	33.5%	
Source: UNAIDS, 2006		
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: 33.0%	F: 40.0%
Source: AIS, 2001		
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: 88.0%	F: 75.0%
Source: AIS, 2001		
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	
% of men who used a condom the last time they had sex with a male partner	-	


PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	0.99
Source: AIS, 2001	
% of orphans receiving external support	-


Number of HIV+ pregnant women receiving ARVs for PMTCT




Number of children in need receiving ART




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Brazil

Estimated adult HIV prevalence rate 0.5%
 Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	97%
Source: SINASC, 2004	
Estimated number of births per year	3,720,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	9,222 [5,829 - 13,361]*
Source: UNAIDS/WHO unpublished estimates, 2007	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	6,510 70.6% [48.7% - >95.0%]*
Source: MOH, 2006	

PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	1,548
Source: MOH, 2006	
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	7,633
Source: MOH-National AIDS Program, 2006	


PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	
HIV prevalence among most-at-risk populations	FSW: 2.9%	IDU: 46.0%
Source: UNAIDS, 2006		
Knowledge of HIV prevention among young people aged 15-24	M: -	F: -
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: 71.3%	F: 58.2%
Source: MOH-National aids Programme (DST/AIDS), 2004		
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	
% of men who used a condom the last time they had sex with a male partner	-	


PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-


Number of HIV+ pregnant women receiving ARVs for PMTCT




Number of children in need receiving ART




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Brunei Darussalam

Estimated adult HIV prevalence rate

<0.1%

Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	-	-
Estimated number of births per year	8,000	-
<small>Source: UN Population Division, 2006</small>		
Estimated number of HIV+ pregnant women	-	-
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	-	-

PROVIDE PAEDIATRIC HIV TREATMENT



Estimated number of children living with HIV	-	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-	-
Estimated number of children in need of ART	-	-
Number and % of children in need receiving ART	-	-

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE


Estimated HIV prevalence among young people aged 15-24	M:	-	F:	-
HIV prevalence among pregnant women aged 15-24 in capital city				-
HIV prevalence among most-at-risk populations	FSW:	-	IDU:	-
Knowledge of HIV prevention among young people aged 15-24	M:	-	F:	-
Sex before age 15 among young people aged 15-19	M:	-	F:	-
Condom use at last higher-risk sex among young people aged 15-24	M:	-	F:	-
% of sex workers who used a condom with their most recent client	M:	-	F:	-
% of IDUs who used sterile equipment at last injection				-
% of men who used a condom the last time they had sex with a male partner				-

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS


Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.

* Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Bulgaria

Estimated adult HIV prevalence rate

<0.1%

Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	-	-
Estimated number of births per year	69,000	-
<small>Source: UN Population Division, 2006</small>		
Estimated number of HIV+ pregnant women	-	-
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	-	-

PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-	-
Estimated number of children in need of ART	-	-
Number and % of children in need receiving ART	3	-
<small>Source: MOH, 2006</small>		


PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: 15.3%	F: 17.2%
<small>Source: National Committee on Prevention of AIDS and STDs at the Council of Ministers UNGASS Indicators country report, 2005</small>		
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: 70.3%	F: 57.0%
<small>Source: National Committee on Prevention of AIDS and STDs at the Council of Ministers UNGASS Indicators country report, 2005</small>		
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-


PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-


Number of HIV+ pregnant women receiving ARVs for PMTCT




Number of children in need receiving ART




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
* Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Burkina Faso

Estimated adult HIV prevalence rate 1.8%
 Source: DHS, 2003

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	85%
Source: MICS, 2006	
Estimated number of births per year	641,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	3,029
Source: UNAIDS/WHO unpublished estimates, 2007 [2,014 - 4,221]*	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	1,615 53.3%
Source: MOH, 2006 [38.3% - 80.2%]*	

PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	534
Source: MOH, 2006	


PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: 0.5%	F: 1.4%
Source: UNAIDS, 2006		
HIV prevalence among pregnant women aged 15-24 in capital city	1.8%	
Source: UNAIDS, 2006		
HIV prevalence among most-at-risk populations	FSW: 20.8%	IDU: -
Source: UNAIDS, 2006		
Knowledge of HIV prevention among young people aged 15-24	M: -	F: 19.0%
Source: MICS, 2006		
Sex before age 15 among young people aged 15-19	M: 4.7%	F: 7.3%
Source: DHS, 2003		
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: 63.7%
Source: MICS, 2006		
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	
% of men who used a condom the last time they had sex with a male partner	-	


PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	0.71
Source: MICS, 2006	
% of orphans receiving external support	-


Number of HIV+ pregnant women receiving ARVs for PMTCT




Number of children in need receiving ART




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.

* Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Burundi

Estimated adult HIV prevalence rate 3.6%
 Source: UNAIDS/WHO, 2007

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	92%
Source: MICS, 2005	
Estimated number of births per year	381,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	8,208
Source: UNAIDS/WHO unpublished estimates, 2007 [5,462 - 10,743]*	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	1,112 13.5%
Source: MOH, 2006 [10.4% - 20.4%]*	

PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	814 -
Source: MOH, 2006	
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	0 -
Source: MOH, 2006	
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	524 -
Source: MOH, 2006	


PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: 0.8%	F: 2.3%
Source: UNAIDS, 2006		
HIV prevalence among pregnant women aged 15-24 in capital city	8.6%	
Source: UNAIDS, 2006		
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: -	F: 30.4%
Source: MICS, 2005		
Sex before age 15 among young people aged 15-19	M: -	F: 3.1%
Source: MICS, 2005		
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: 25.3%
Source: MICS, 2005		
% of sex workers who used a condom with their most recent client	M: -	F: 73.8%
Source: UNAIDS, 2006		
% of IDUs who used sterile equipment at last injection	-	
% of men who used a condom the last time they had sex with a male partner	-	


PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	0.86
Source: MICS, 2005	
% of orphans receiving external support	-


Number of HIV+ pregnant women receiving ARVs for PMTCT




Number of children in need receiving ART




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Cambodia

Estimated adult HIV prevalence rate 0.6%
 Source: DHS, 2005

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	69%
Source: DHS, 2005	
Estimated number of births per year	377,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	1,502
Source: UNAIDS/WHO unpublished estimates, 2007 [1,170 - 1,858]*	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	312 20.8%
Source: MOH, 2006 [16.8% - 26.7%]*	

PROVIDE PAEDIATRIC HIV TREATMENT






Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	86 -
Source: MOH, 2006	
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	- -
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	1,787 -
Source: MOH, 2006	

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: 0.1%	F: 0.3%
Source: DHS, 2005		
HIV prevalence among pregnant women aged 15-24 in capital city	2.7%	
Source: SOWC, 2008		
HIV prevalence among most-at-risk populations	FSW: 26.3%	IDU: -
Source: UNAIDS, 2006		
Knowledge of HIV prevention among young people aged 15-24	M: 45.2%	F: 50.1%
Source: DHS, 2005		
Sex before age 15 among young people aged 15-19	M: 0.4%	F: 0.7%
Source: DHS, 2005		
Condom use at last higher-risk sex among young people aged 15-24	M: 84.4%	F: -
Source: DHS, 2005		
% of sex workers who used a condom with their most recent client	M: -	F: 96.0%
Source: UNAIDS, 2006		
% of IDUs who used sterile equipment at last injection	-	
% of men who used a condom the last time they had sex with a male partner	-	

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	0.83
Source: DHS, 2005	
% of orphans receiving external support	-



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Cameroon

Estimated adult HIV prevalence rate 5.5%
 Source: DHS, 2004

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	82%
Source: MICS, 2006	
Estimated number of births per year	649,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	35,127 [25,363 - 41,913]*
Source: UNAIDS/WHO unpublished estimates, 2007	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	7,588 21.6% [18.1% - 29.9%]*
Source: MOH, 2006	

PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	1,014
Source: MOH, 2006	


PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: 1.4%	F: 4.8%
Source: DHS, 2004		
HIV prevalence among pregnant women aged 15-24 in capital city	7.0%	
Source: SOWC, 2002		
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: -	F: 32.0%
Source: DHS, 2004		
Sex before age 15 among young people aged 15-19	M: -	F: 13.4%
Source: MICS, 2006		
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: 61.8%
Source: MICS, 2006		
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	
% of men who used a condom the last time they had sex with a male partner	-	


PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	0.89
Source: MICS, 2006	
% of orphans receiving external support	9.3%
Source: MICS, 2006	


Number of HIV+ pregnant women receiving ARVs for PMTCT




Number of children in need receiving ART




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Cape Verde

Estimated adult HIV prevalence rate -

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	-
Estimated number of births per year	15,000
<small>Source: UN Population Division, 2006</small>	
Estimated number of HIV+ pregnant women	-
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	43 -
<small>Source: MOH, 2006</small>	

PROVIDE PAEDIATRIC HIV TREATMENT



Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	31 -
<small>Source: MOH, 2006</small>	
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	0 -
<small>Source: MOH, 2006</small>	
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	22 -
<small>Source: MOH, 2006</small>	

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE


Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: -	F: -
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS


Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-



% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.

* Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Central African Republic

Estimated adult HIV prevalence rate 6.2%

Source: UNAIDS/WHO, 2007

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	69%
Source: MICS, 2006	
Estimated number of births per year	157,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	10,913
Source: UNAIDS/WHO unpublished estimates, 2007 [9,791 - 12,218]*	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	1,943 17.8%
Source: MOH, 2006 [15.9% - 19.8%]*	

PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	443 -
Source: MOH, 2006	
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	19 -
Source: MOH, 2006	
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	268 -
Source: MOH, 2006	


PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: 2.5%	F: 7.3%
Source: UNAIDS, 2006		
HIV prevalence among pregnant women aged 15-24 in capital city	14.0%	
Source: SOWC, 2002		
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: -	F: 17.3%
Source: MICS, 2006		
Sex before age 15 among young people aged 15-19	M: -	F: 29.0%
Source: MICS, 2006		
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: 40.6%
Source: MICS, 2006		
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	
% of men who used a condom the last time they had sex with a male partner	-	


PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	0.96
Source: MICS, 2006	
% of orphans receiving external support	7.4%
Source: MICS, 2006	


Number of HIV+ pregnant women receiving ARVs for PMTCT




Number of children in need receiving ART




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.

* Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Chad

Estimated adult HIV prevalence rate 3.3%
 Source: UNAIDS/WHO, 2007

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	39%
Source: DHS, 2004	
Estimated number of births per year	482,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	17,231 [11,000 - 21,586]*
Source: UNAIDS/WHO unpublished estimates, 2007	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	254 1.5% [1.2% - 2.3%]*
Source: MOH, 2006	

PROVIDE PAEDIATRIC HIV TREATMENT






Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	63 -
Source: MOH, 2006	
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	0 -
Source: MOH, 2006	
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	129 -
Source: MOH, 2006	

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: 0.9%	F: 2.2%
Source: UNAIDS, 2006		
HIV prevalence among pregnant women aged 15-24 in capital city	3.6%	
Source: UNAIDS, 2006		
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: 20.2%	F: 8.1%
Source: DHS, 2004		
Sex before age 15 among young people aged 15-19	M: 10.7%	F: 19.0%
Source: DHS, 2004		
Condom use at last higher-risk sex among young people aged 15-24	M: 25.0%	F: 17.0%
Source: DHS, 2004		
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	
% of men who used a condom the last time they had sex with a male partner	-	

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	1.05
Source: DHS, 2004	
% of orphans receiving external support	-



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Chile

Estimated adult HIV prevalence rate

0.3%

Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	-	-
Estimated number of births per year	249,000	
<small>Source: UN Population Division, 2006</small>		
Estimated number of HIV+ pregnant women	<500	
<small>Source: UNAIDS/WHO unpublished estimates, 2007</small>		
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	-	-

PROVIDE PAEDIATRIC HIV TREATMENT



Estimated number of children living with HIV	-	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-	-
Estimated number of children in need of ART	-	-
Number and % of children in need receiving ART	-	-

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE


Estimated HIV prevalence among young people aged 15-24	M:	-	F:	-
HIV prevalence among pregnant women aged 15-24 in capital city				-
HIV prevalence among most-at-risk populations	FSW:	-	IDU:	-
Knowledge of HIV prevention among young people aged 15-24	M:	-	F:	-
Sex before age 15 among young people aged 15-19	M:	-	F:	-
Condom use at last higher-risk sex among young people aged 15-24	M:	-	F:	-
% of sex workers who used a condom with their most recent client	M:	-	F:	-
% of IDUs who used sterile equipment at last injection				-
% of men who used a condom the last time they had sex with a male partner				-

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS


Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.

* Represents the uncertainty range around the estimates

Children and AIDS fact sheet: China

Estimated adult HIV prevalence rate 0.1%
 Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	90%
Source: Social Progress in China, 2005	
Estimated number of births per year	17,309,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	6,899 [4,255 - 10,777]*
Source: UNAIDS/WHO unpublished estimates, 2007	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	650 9.4% [6.0% - 15.3%]*
Source: MOH, 2006	

PROVIDE PAEDIATRIC HIV TREATMENT



Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	650 -
Source: MOH, 2006	
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	- -
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	410 -
Source: MOH, 2006	

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE


Estimated HIV prevalence among young people aged 15-24	M: - F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-
HIV prevalence among most-at-risk populations	FSW: 0.5% IDU: 8.3%
Source: UNAIDS, 2006	
Knowledge of HIV prevention among young people aged 15-24	M: - F: -
Sex before age 15 among young people aged 15-19	M: - F: -
Condom use at last higher-risk sex among young people aged 15-24	M: - F: -
% of sex workers who used a condom with their most recent client	M: - F: 68.5%
Source: UNAIDS, 2006	
% of IDUs who used sterile equipment at last injection	-
% of men who used a condom the last time they had sex with a male partner	41.1%
Source: UNAIDS, 2006	

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS


Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Colombia

Estimated adult HIV prevalence rate 0.6%
 Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	94%
Source: DHS, 2005	
Estimated number of births per year	884,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	2,452
Source: UNAIDS/WHO unpublished estimates, 2007	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	112 4.6%
Source: MOH, 2006	
	[2.6% - 7.5%]*

PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	2
Source: MOH, 2006	
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	-


PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: 0.8%	IDU: -
Source: UNAIDS, 2006		
Knowledge of HIV prevention among young people aged 15-24	M: -	F: -
Sex before age 15 among young people aged 15-19	M: -	F: 13.7%
Source: DHS, 2005		
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: 36.0%
Source: RHS, 2005		
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-


PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	0.85
Source: DHS, 2005	
% of orphans receiving external support	-


Number of HIV+ pregnant women receiving ARVs for PMTCT




Number of children in need receiving ART




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Comoros

Estimated adult HIV prevalence rate <0.1%
Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	75%
<small>Source: EIM, 2004</small>	
Estimated number of births per year	28,000
<small>Source: UN Population Division, 2006</small>	
Estimated number of HIV+ pregnant women	<100 [<100]*
<small>Source: UNAIDS/WHO unpublished estimates, 2007</small>	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	0 -
<small>Source: MOH, 2006</small>	

PROVIDE PAEDIATRIC HIV TREATMENT






Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	0 -
<small>Source: MOH, 2006</small>	
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	0 -
<small>Source: MOH, 2006</small>	
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	0 -
<small>Source: MOH, 2006</small>	

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: <0.1%	F: <0.1%
<small>Source: UNAIDS, 2006</small>		
HIV prevalence among pregnant women aged 15-24 in capital city		-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: -	F: 17.9%
<small>Source: MICS, 2000</small>		
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection		-
% of men who used a condom the last time they had sex with a male partner		-

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	0.59
<small>Source: MICS, 2000</small>	
% of orphans receiving external support	-



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Congo

Estimated adult HIV prevalence rate 5.3%
 Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	86%
Source: DHS, 2005	
Estimated number of births per year	132,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	4,523 [3,707 - 5,457]*
Source: UNAIDS/WHO unpublished estimates, 2007	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	325 7.2% [6.0% - 8.8%]*
Source: MOH, 2006	

PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	447 -
Source: MOH, 2006	
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	264 -
Source: MOH, 2006	


PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: 1.2%	F: 3.7%
Source: UNAIDS, 2006		
HIV prevalence among pregnant women aged 15-24 in capital city	3.0%	
Source: SOWC, 2002		
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: 22.0%	F: 10.0%
Source: DHS, 2005		
Sex before age 15 among young people aged 15-19	M: 25.3%	F: 23.5%
Source: DHS, 2005		
Condom use at last higher-risk sex among young people aged 15-24	M: 36.0%	F: 16.0%
Source: DHS, 2005		
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	
% of men who used a condom the last time they had sex with a male partner	23.3%	
Source: UNAIDS, 2006		


PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	0.88
Source: DHS, 2005	
% of orphans receiving external support	2.0
Source: DHS, 2005	


Number of HIV+ pregnant women receiving ARVs for PMTCT




Number of children in need receiving ART




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Congo, Democratic Republic of the

Estimated adult HIV prevalence rate 3.2%
 Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	68%
Source: MICS, 2001	
Estimated number of births per year	3,026,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	90,077 [70,480 - 112,999]*
Source: UNAIDS/WHO unpublished estimates, 2007	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	3,422 3.8% [3.0% - 4.9%]*
Source: MOH, 2006	

PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	170 -
Source: MOH, 2006	
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	- -
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	124 -
Source: MOH, 2006	


PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: 0.8%	F: 2.2%
Source: UNAIDS, 2006		
HIV prevalence among pregnant women aged 15-24 in capital city	3.2%	
Source: UNAIDS, 2006		
HIV prevalence among most-at-risk populations	FSW: 12.4%	IDU: -
Source: UNAIDS, 2006		
Knowledge of HIV prevention among young people aged 15-24	M: -	F: -
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	
% of men who used a condom the last time they had sex with a male partner	-	


PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	0.72
Source: MICS, 2000	
% of orphans receiving external support	-

Number of HIV+ pregnant women receiving ARVs for PMTCT




Number of children in need receiving ART




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.


* Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Cook Islands

Estimated adult HIV prevalence rate -


PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	-
Estimated number of births per year <small>Source: UN Population Division, 2006</small>	0
Estimated number of HIV+ pregnant women	-
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	-



PROVIDE PAEDIATRIC HIV TREATMENT


Estimated number of children living with HIV <small>Source</small>	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	-




PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: -	F: -
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-

% of young people aged 15-24 who used a condom at last higher-risk sex




HIV prevalence among female sex workers (%)



PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Year refers to the year of data, not publication date. Dash (-) means data not available.
* Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Costa Rica

Estimated adult HIV prevalence rate 0.3%
 Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	92%	
<small>Source: CCSS, 2004</small>		
Estimated number of births per year	80,000	
<small>Source: UN Population Division, 2006</small>		
Estimated number of HIV+ pregnant women	<200	[<100 - <500]*
<small>Source: UNAIDS/WHO unpublished estimates, 2007</small>		
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	21	-
<small>Source: MOH, 2006</small>		

PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-	
<small>Source</small>		
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	40	-
<small>Source: MOH, 2006</small>		
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	40	-
<small>Source: MOH, 2006</small>		
Estimated number of children in need of ART	-	
Number and % of children in need receiving ART	52	-
<small>Source: MOH, 2006</small>		


PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city		5.2%
<small>Source: SOWC, 2008</small>		
HIV prevalence among most-at-risk populations	FSW: 0.1%	IDU: -
<small>Source: UNAIDS, 2006</small>		
Knowledge of HIV prevention among young people aged 15-24	M: -	F: -
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection		-
% of men who used a condom the last time they had sex with a male partner		-


PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-


Number of HIV+ pregnant women receiving ARVs for PMTCT




Number of children in need receiving ART




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Côte d'Ivoire

Estimated adult HIV prevalence rate 4.7%

Source: AIS, 2005

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	85%
Source: MICS, 2006	
Estimated number of births per year	684,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	30,412 [24,278 - 36,289]*
Source: UNAIDS/WHO unpublished estimates, 2007	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	2,773 9.1% [7.6% - 11.4%]*
Source: MOH, 2006	

PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	1,348
Source: MOH, 2006	


PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: 0.3%	F: 2.4%
Source: AIS, 2005		
HIV prevalence among pregnant women aged 15-24 in capital city	5.2%	
Source: SOWC, 2008		
HIV prevalence among most-at-risk populations	FSW: 28.0%	IDU: -
Source: UNAIDS, 2006		
Knowledge of HIV prevention among young people aged 15-24	M: 27.6%	F: 18.0%
Source: AIS, 2005		
Sex before age 15 among young people aged 15-19	M: 16.7%	F: 20.4%
Source: AIS, 2005		
Condom use at last higher-risk sex among young people aged 15-24	M: 52.5%	F: 39.3%
Source: AIS, 2005		
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	
% of men who used a condom the last time they had sex with a male partner	-	


PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	0.79
Source: MICS, 2006	
% of orphans receiving external support	9.3%
Source: AIS, 2005	


Number of HIV+ pregnant women receiving ARVs for PMTCT




Number of children in need receiving ART




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.

* Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Croatia

Estimated adult HIV prevalence rate

<0.1%

Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	-	-
Estimated number of births per year	41,000	-
<small>Source: UN Population Division, 2006</small>		
Estimated number of HIV+ pregnant women	-	-
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	-	-

PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-	-
Estimated number of children in need of ART	-	-
Number and % of children in need receiving ART	5	-
<small>Source: MOH, 2006</small>		

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: -	F: -
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-


PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-


Number of HIV+ pregnant women receiving ARVs for PMTCT




Number of children in need receiving ART




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.

* Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Cuba

Estimated adult HIV prevalence rate 0.1%
 Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	100%
Source: Statistical System of External Consultation, 2005	
Estimated number of births per year	121,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	<100 [<100 - <200]*
Source: UNAIDS/WHO unpublished estimates, 2007	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	34 -
Source: MOH, 2006	

PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	1 -
Source: MOH, 2006	
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	34 -
Source: MOH, 2006	
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	17 -
Source: MOH, 2006	


PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: -	F: -
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-


PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-


Number of HIV+ pregnant women receiving ARVs for PMTCT




Number of children in need receiving ART




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Djibouti

Estimated adult HIV prevalence rate 3.1%
 Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	67%
Source: Programme Sante de la Reproduction, 2003	
Estimated number of births per year	24,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	828 [626 - 1,035]*
Source: UNAIDS/WHO unpublished estimates, 2007	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	52 6.3% [5.0% - 8.3%]*
Source: MOH, 2006	

PROVIDE PAEDIATRIC HIV TREATMENT



Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	52 -
Source: MOH, 2006	
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	- -
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	52 -
Source: MOH, 2006	

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE


Estimated HIV prevalence among young people aged 15-24	M: 0.7%	F: 2.1%
Source: UNAIDS, 2006		
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: -	F: 18.4%
Source: MICS, 2006		
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: 50.0%	F: 26.0%
Source: UNICEF/Ministry of Youth and Sports, 2006		
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS


Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	0.82
Source: MICS, 2006	
% of orphans receiving external support	-




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.

* Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Dominica

Estimated adult HIV prevalence rate -

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	100%
<small>Source: PAHO/WHO, 2004</small>	
Estimated number of births per year	0
<small>Source: UN Population Division, 2006</small>	
Estimated number of HIV+ pregnant women	-
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	1 -
<small>Source: MOH, 2006</small>	

PROVIDE PAEDIATRIC HIV TREATMENT






Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	2 -
<small>Source: MOH, 2006</small>	
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	0 -
<small>Source: MOH, 2006</small>	
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	2 -
<small>Source: MOH, 2006</small>	

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: -	F: -
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Dominican Republic

Estimated adult HIV prevalence rate 1.0%
 Source: DHS, 2002

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	99%
Source: DHS, 2002	
Estimated number of births per year	231,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	1,798
Source: UNAIDS/WHO unpublished estimates, 2007	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	738 41.0%
Source: MOH, 2006	
[31.5% - 54.0%]*	

PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	377
Source: MOH, 2006	


PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: 0.4%	F: 0.7%
Source: DHS, 2002		
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: 3.6%	IDU: -
Source: UNAIDS, 2006		
Knowledge of HIV prevention among young people aged 15-24	M: -	F: 36.1%
Source: MICS, 2006		
Sex before age 15 among young people aged 15-19	M: 18.4%	F: 12.8%
Source: DHS, 2002		
Condom use at last higher-risk sex among young people aged 15-24	M: 52.0%	F: 29.0%
Source: DHS, 2002		
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-


PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	0.70
Source: MICS, 2006	
% of orphans receiving external support	-


Number of HIV+ pregnant women receiving ARVs for PMTCT




Number of children in need receiving ART




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.

* Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Ecuador

Estimated adult HIV prevalence rate 0.3%
 Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	84%
Source: ENDEMAIN (RHS), 2004	
Estimated number of births per year	285,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	<500 [500 - 759]*
Source: UNAIDS/WHO unpublished estimates, 2007	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	213 -
Source: MOH, 2006	

PROVIDE PAEDIATRIC HIV TREATMENT






Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	- -
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	- -
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	300 -
Source: MOH, 2006	

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: -	F: -
Sex before age 15 among young people aged 15-19	M: -	F: 7.0%
Source: RHS, 2001		
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Egypt

Estimated adult HIV prevalence rate

<0.1%

Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	70%
<small>Source: DHS, 2005</small>	
Estimated number of births per year	1,828,000
<small>Source: UN Population Division, 2006</small>	
Estimated number of HIV+ pregnant women	<200
<small>Source: UNAIDS/WHO unpublished estimates, 2007</small>	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	- -

PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	- -
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	- -
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	- -


PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: -	F: 3.7%
<small>Source: DHS, 2005</small>		
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-


PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-


Number of HIV+ pregnant women receiving ARVs for PMTCT




Number of children in need receiving ART




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.

* Represents the uncertainty range around the estimates

Children and AIDS fact sheet: El Salvador

Estimated adult HIV prevalence rate 0.9%
 Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	86%
Source: FESAL, 2002-2003	
Estimated number of births per year	159,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	629
Source: UNAIDS/WHO unpublished estimates, 2007	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	193 30.7%
Source: MOH, 2006	
[18.3% - 53.4%]*	

PROVIDE PAEDIATRIC HIV TREATMENT






Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	122 -
Source: MOH, 2006	
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	147 -
Source: MOH, 2006	
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	297 -
Source: MOH, 2006	

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: 3.9%	IDU: -
Source: UNAIDS, 2006		
Knowledge of HIV prevention among young people aged 15-24	M: -	F: -
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Equatorial Guinea

Estimated adult HIV prevalence rate 3.2%
 Source: UNAIDS/WHO, 2007

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	86%
Source: MICS, 2000	
Estimated number of births per year	19,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	712
Source: UNAIDS/WHO unpublished estimates, 2007 [527 - 953]*	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	103 14.5%
Source: MOH, 2006 [10.8% - 19.5%]*	

PROVIDE PAEDIATRIC HIV TREATMENT






Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	-

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: 0.7%	F: 2.3%
Source: UNAIDS, 2006		
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: -	F: 4.0%
Source: MICS, 2000		
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	0.95
Source: MICS, 2000	
% of orphans receiving external support	-



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Eritrea

Estimated adult HIV prevalence rate 2.4%
 Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	70%
Source: DHS, 2002	
Estimated number of births per year	186,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	2,433
Source: UNAIDS/WHO unpublished estimates, 2007 [1,624 - 3,592]*	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	75 3.1%
Source: MOH, 2006 [2.1% - 4.6%]*	

PROVIDE PAEDIATRIC HIV TREATMENT



Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	0
Source: MOH, 2006	
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	84
Source: MOH, 2006	

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE


Estimated HIV prevalence among young people aged 15-24	M: 0.6%	F: 1.6%
Source: UNAIDS, 2006		
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: -	F: 37.0%
Source: DHS, 2002		
Sex before age 15 among young people aged 15-19	M: -	F: 8.8%
Source: DHS, 2002		
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS


Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	0.83
Source: DHS, 2002	
% of orphans receiving external support	-




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. B Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Ethiopia

Estimated adult HIV prevalence rate 1.4%
 Source: DHS, 2005

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	28%
Source: DHS, 2005	
Estimated number of births per year	3,159,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	65,585
Source: UNAIDS/WHO unpublished estimates, 2007 [57,912 - 74,324]*	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	2,330 3.6%
Source: MOH, 2006 [3.1% - 4.0%]*	

PROVIDE PAEDIATRIC HIV TREATMENT



Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	388 -
Source: MOH, 2006	
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	94 -
Source: MOH, 2006	
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	2,512 -
Source: MOH, 2006	

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE


Estimated HIV prevalence among young people aged 15-24	M: 0.3%	F: 1.2%
Source: DHS, 2005		
HIV prevalence among pregnant women aged 15-24 in capital city	11.5%	
Source: UNAIDS, 2006		
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: 33.3%	F: 20.5%
Source: DHS, 2005		
Sex before age 15 among young people aged 15-19	M: 1.7%	F: 11.1%
Source: DHS, 2005		
Condom use at last higher-risk sex among young people aged 15-24	M: 50.2%	F: 28.4%
Source: DHS, 2005		
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	
% of men who used a condom the last time they had sex with a male partner	-	

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS


Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	0.90
Source: DHS, 2005	
% of orphans receiving external support	-




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Fiji

Estimated adult HIV prevalence rate 0.1%
 Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	-	-
Estimated number of births per year	18,000	-
<small>Source: UN Population Division, 2006</small>		
Estimated number of HIV+ pregnant women	<100	[<100]*
<small>Source: UNAIDS/WHO unpublished estimates, 2007</small>		
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	5	-
<small>Source: MOH, 2006</small>		

PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	2	-
<small>Source: MOH, 2006</small>		
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-	-
Estimated number of children in need of ART	-	-
Number and % of children in need receiving ART	2	-
<small>Source: MOH, 2006</small>		


PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: -	F: -
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	20.0%
<small>Source: UNAIDS, 2006</small>		


PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-


Number of HIV+ pregnant women receiving ARVs for PMTCT




Number of children in need receiving ART




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Gabon

Estimated adult HIV prevalence rate 7.9%
 Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	94%
Source: DHS, 2000	
Estimated number of births per year	34,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	2,300
Source: UNAIDS/WHO unpublished estimates, 2007 [1,651 - 3,316]*	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	89 3.9%
Source: MOH, 2006 [2.7% - 5.4%]*	

PROVIDE PAEDIATRIC HIV TREATMENT






Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	362
Source: MOH, 2006	

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: 1.8%	F: 5.4%
Source: UNAIDS, 2006		
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: 40.4%	IDU: -
Source: UNAIDS, 2006		
Knowledge of HIV prevention among young people aged 15-24	M: 22.0%	F: 24.0%
Source: DHS, 2000		
Sex before age 15 among young people aged 15-19	M: 48.1%	F: 23.8%
Source: DHS, 2000		
Condom use at last higher-risk sex among young people aged 15-24	M: 48.0%	F: 33.0%
Source: DHS, 2000		
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	0.98
Source: DHS, 2000	
% of orphans receiving external support	-



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Gambia

Estimated adult HIV prevalence rate 2.4%
 Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	98%
Source: MICS, 2006	
Estimated number of births per year	60,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	500
Source: UNAIDS/WHO unpublished estimates, 2007	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	103 20.6%
Source: MOH, 2006 [13.2% - 44.0%]*	

PROVIDE PAEDIATRIC HIV TREATMENT



Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	0
Source: MOH, 2006	
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	83
Source: MOH, 2006	

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE


Estimated HIV prevalence among young people aged 15-24	M: 0.6%	F: 1.7%
Source: UNAIDS, 2006		
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: -	F: 39.4%
Source: MICS, 2006		
Sex before age 15 among young people aged 15-19	M: -	F: 3.9%
Source: MICS, 2006		
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: 54.3%
Source: MICS, 2006		
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS


Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	0.87
Source: MICS, 2006	
% of orphans receiving external support	-




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Georgia

Estimated adult HIV prevalence rate 0.2%
 Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	94%	
<small>Source: RHS, 2005</small>		
Estimated number of births per year	48,000	
<small>Source: UN Population Division, 2006</small>		
Estimated number of HIV+ pregnant women	<100	<100*
<small>Source: UNAIDS/WHO unpublished estimates, 2007</small>		
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	15	-
<small>Source: MOH, 2006</small>		

PROVIDE PAEDIATRIC HIV TREATMENT



Estimated number of children living with HIV	-	
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	15	-
<small>Source: MOH, 2006</small>		
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	15	-
<small>Source: MOH, 2006</small>		
Estimated number of children in need of ART	-	
Number and % of children in need receiving ART	8	-
<small>Source: MOH, 2006</small>		

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE


Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city		-
HIV prevalence among most-at-risk populations	FSW: 1.3%	IDU: 1.0%
<small>Source: UNAIDS, 2006</small>		
Knowledge of HIV prevention among young people aged 15-24	M: -	F: -
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: 95.0%
<small>Source: UNAIDS, 2006</small>		
% of IDUs who used sterile equipment at last injection		17.0%
<small>Source: BSS, 2004</small>		
% of men who used a condom the last time they had sex with a male partner		53.6%
<small>Source: UNAIDS, 2006</small>		

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS


Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Ghana

Estimated adult HIV prevalence rate 2.2%
 Source: DHS, 2003

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	92%
Source: MICS, 2006	
Estimated number of births per year	700,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	15,183 [13,146 - 17,450]*
Source: UNAIDS/WHO unpublished estimates, 2007	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	1,239 8.2% [7.1% - 9.4%]*
Source: MOH, 2006	

PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	122
Source: MOH, 2006	


PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: 0.1%	F: 1.2%
Source: DHS, 2003		
HIV prevalence among pregnant women aged 15-24 in capital city	3.9%	
Source: SOWC, 2003		
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: -	F: 25.1%
Source: MICS, 2006		
Sex before age 15 among young people aged 15-19	M: -	F: 6.5%
Source: MICS, 2006		
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: 41.8%
Source: MICS, 2006		
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	
% of men who used a condom the last time they had sex with a male partner	-	


PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	0.90
Source: MICS, 2006	
% of orphans receiving external support	-


Number of HIV+ pregnant women receiving ARVs for PMTCT




Number of children in need receiving ART




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Grenada

Estimated adult HIV prevalence rate -

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage <small>Source: PAHO/WHO, 2005</small>	100%
Estimated number of births per year <small>Source: UN Population Division, 2006</small>	2,000
Estimated number of HIV+ pregnant women	-
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	-

PROVIDE PAEDIATRIC HIV TREATMENT



Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	-

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE


Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: -	F: -
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS


Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
* Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Guatemala

Estimated adult HIV prevalence rate 0.9%
 Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	84%
Source: ENSMI, 2002	
Estimated number of births per year	445,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	1,925 [1,185 - 3,472]*
Source: UNAIDS/WHO unpublished estimates, 2007	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	167 8.7% [4.8% - 14.1%]*
Source: MOH, 2006	

PROVIDE PAEDIATRIC HIV TREATMENT



Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	840
Source: MOH, 2006	

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE


Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	
HIV prevalence among most-at-risk populations	FSW: 8.7%	IDU: -
Source: UNAIDS, 2006		
Knowledge of HIV prevention among young people aged 15-24	M: -	F: -
Sex before age 15 among young people aged 15-19	M: 15.0%	F: 7.0%
Source: RHS, 2002		
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	
% of men who used a condom the last time they had sex with a male partner	-	

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS


Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Guinea

Estimated adult HIV prevalence rate 1.5%
 Source: DHS, 2005

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage Source: DHS, 2005	82%
Estimated number of births per year Source: UN Population Division, 2006	374,000
Estimated number of HIV+ pregnant women Source: UNAIDS/WHO unpublished estimates, 2007	6,074 [4,905 - 8,036]*
Number and % of HIV+ pregnant women receiving ARVs for PMTCT Source: MOH, 2006	486 8.0% [6.0% - 9.9%]*

PROVIDE PAEDIATRIC HIV TREATMENT



Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth Source: MOH, 2006	416 -
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	- -
Estimated number of children in need of ART	-
Number and % of children in need receiving ART Source: MOH, 2006	100 -

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE


Estimated HIV prevalence among young people aged 15-24 Source: DHS, 2005	M: 0.6%	F: 1.2%
HIV prevalence among pregnant women aged 15-24 in capital city Source: UNAIDS, 2006	4.4%	
HIV prevalence among most-at-risk populations Source: UNAIDS, 2006	FSW: 39.7%	IDU: -
Knowledge of HIV prevention among young people aged 15-24 Source: DHS, 2005	M: 22.8%	F: 16.9%
Sex before age 15 among young people aged 15-19 Source: DHS, 2005	M: 17.9%	F: 19.7%
Condom use at last higher-risk sex among young people aged 15-24 Source: DHS, 2005	M: 37.2%	F: 25.8%
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	
% of men who used a condom the last time they had sex with a male partner	-	

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS


Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio Source: DHS, 2005	0.73
% of orphans receiving external support	-




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.

* Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Guinea-Bissau

Estimated adult HIV prevalence rate 3.8%
 Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	78%
Source: MICS, 2006	
Estimated number of births per year	82,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	1,454
Source: UNAIDS/WHO unpublished estimates, 2007 [1,020 - 2,082]*	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	195 13.4%
Source: MOH, 2006 [9.4% - 19.1%]*	

PROVIDE PAEDIATRIC HIV TREATMENT






Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	0
Source: MOH, 2006	
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	24
Source: MOH, 2006	

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: 0.9%	F: 2.5%
Source: UNAIDS, 2006		
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: -	F: 17.7%
Source: MICS, 2006		
Sex before age 15 among young people aged 15-19	M: -	F: 21.8%
Source: MICS, 2006		
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: 38.8%
Source: MICS, 2006		
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	0.97
Source: MICS, 2006	
% of orphans receiving external support	-



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Guyana

Estimated adult HIV prevalence rate 2.4%
 Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage <small>Source: PAHO/WHO, 2003</small>	90%	
Estimated number of births per year <small>Source: UN Population Division, 2006</small>	13,000	
Estimated number of HIV+ pregnant women <small>Source: UNAIDS/WHO unpublished estimates, 2007</small>	<500	[<500 - 519]*
Number and % of HIV+ pregnant women receiving ARVs for PMTCT <small>Source: MOH, 2006</small>	144	-

PROVIDE PAEDIATRIC HIV TREATMENT






Estimated number of children living with HIV	-	
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth <small>Source: MOH, 2006</small>	90	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth <small>Source: MOH, 2006</small>	0	-
Estimated number of children in need of ART	-	
Number and % of children in need receiving ART <small>Source: MOH, 2006</small>	120	-

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city		-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24 <small>Source: AIS, 2005</small>	M: 47.3%	F: 52.6%
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24 <small>Source: AIS, 2005</small>	M: 67.6%	F: 61.6%
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection		-
% of men who used a condom the last time they had sex with a male partner		-

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio <small>Source: MICS, 2006</small>	1.08
% of orphans receiving external support <small>Source: MICS, 2006</small>	12.5



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Haiti

Estimated adult HIV prevalence rate 2.2%
 Source: DHS, 2005-2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	85%
Source: DHS, 2005-2006	
Estimated number of births per year	269,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	4,946 [4,099 - 5,886]*
Source: UNAIDS/WHO unpublished estimates, 2007	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	991 20.0% [16.8% - 24.2%]*
Source: MOH, 2006	

PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	867
Source: MOH, 2006	


PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: 0.6%	F: 1.5%
Source: DHS, 2005-2006		
HIV prevalence among pregnant women aged 15-24 in capital city	3.7%	
Source: UNAIDS, 2006		
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: 40.4%	F: 31.9%
Source: DHS, 2005-2006		
Sex before age 15 among young people aged 15-19	M: 41.9%	F: 15.3%
Source: DHS, 2005-2006		
Condom use at last higher-risk sex among young people aged 15-24	M: 43.3%	F: 29.2%
Source: DHS, 2005-2006		
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	
% of men who used a condom the last time they had sex with a male partner	-	


PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	0.86
Source: DHS, 2005-2006	
% of orphans receiving external support	5.2%
Source: DHS, 2005-2006	


Number of HIV+ pregnant women receiving ARVs for PMTCT




Number of children in need receiving ART




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Honduras

Estimated adult HIV prevalence rate 1.5%
 Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	92%
Source: DHS, 2005-2006	
Estimated number of births per year	199,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	656
Source: UNAIDS/WHO unpublished estimates, 2007	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	200 30.5%
Source: MOH, 2006	
	[19.8% - 45.6%]*

PROVIDE PAEDIATRIC HIV TREATMENT



Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	378
Source: MOH, 2006	
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	672
Source: MOH, 2006	

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE


Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: 9.7%	IDU: -
Source: UNAIDS, 2006		
Knowledge of HIV prevention among young people aged 15-24	M: -	F: 29.9%
Source: DHS, 2005-2006		
Sex before age 15 among young people aged 15-19	M: -	F: 9.8%
Source: DHS, 2005-2006		
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: 23.7%
Source: DHS, 2005-2006		
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS


Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	1.08
Source: DHS, 2005-2006	
% of orphans receiving external support	-




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.

* Represents the uncertainty range around the estimates

Children and AIDS fact sheet: India

Estimated adult HIV prevalence rate 0.28%
 Source: NFHS-3, 2005-2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	74%
Source: NFHS-3, 2005-2006	
Estimated number of births per year	27,195,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	70,489 [43,952 - 111,189]*
Source: UNAIDS/WHO unpublished estimates, 2007	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	7,007 9.9% [6.3% - 15.9%]*
Source: MOH, 2006	

PROVIDE PAEDIATRIC HIV TREATMENT



Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	1,200 -
Source: MOH, 2006	
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	- -
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	2,959 -
Source: MOH, 2006	

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE


Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: 9.4%	IDU: -
Source: UNAIDS, 2006		
Knowledge of HIV prevention among young people aged 15-24	M: 36.1%	F: 19.9%
Source: NFHS-3, 2005-2006		
Sex before age 15 among young people aged 15-19	M: 2.7%	F: 8.0%
Source: NFHS-3, 2005-2006		
Condom use at last higher-risk sex among young people aged 15-24	M: 36.5%	F: 22.2%
Source: NFHS-3, 2005-2006		
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS


Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Indonesia

Estimated adult HIV prevalence rate 0.1%
 Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	92%
Source: DHS, 2002-2003	
Estimated number of births per year	4,427,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	2,724 [1,382 - 5,802]*
Source: UNAIDS/WHO unpublished estimates, 2007	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	40 1.5% [<0.7% - 2.9%]*
Source: MOH, 2006	

PROVIDE PAEDIATRIC HIV TREATMENT



Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	20 -
Source: MOH, 2006	
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	18 -
Source: MOH, 2006	
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	1 -
Source: MOH, 2006	

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE


Estimated HIV prevalence among young people aged 15-24	M: - F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-
HIV prevalence among most-at-risk populations	FSW: 0.0% IDU: 65.5%
Source: UNAIDS, 2006	
Knowledge of HIV prevention among young people aged 15-24	M: 0.0% F: 1.0%
Source: DHS, 2002-2003	
Sex before age 15 among young people aged 15-19	M: - F: -
Condom use at last higher-risk sex among young people aged 15-24	M: - F: -
% of sex workers who used a condom with their most recent client	M: 47.5% F: 56.2%
Source: UNAIDS, 2006	
% of IDUs who used sterile equipment at last injection	18.9%
Source: UNAIDS, 2006	
% of men who used a condom the last time they had sex with a male partner	47.6%
Source: UNAIDS, 2006	

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS


Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	0.82
Source: DHS, 2002-2003	
% of orphans receiving external support	-




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Iran (Islamic Republic of)

Estimated adult HIV prevalence rate

0.2%

Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	-	-
Estimated number of births per year	1,407,000	
<small>Source: UN Population Division, 2006</small>		
Estimated number of HIV+ pregnant women	1,222	[805 - 1,863]*
<small>Source: UNAIDS/WHO unpublished estimates, 2007</small>		
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	-	-

PROVIDE PAEDIATRIC HIV TREATMENT



Estimated number of children living with HIV	-	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-	-
Estimated number of children in need of ART	-	-
Number and % of children in need receiving ART	20	-
<small>Source: MOH, 2006</small>		

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE


Estimated HIV prevalence among young people aged 15-24	M:	-	F:	-
HIV prevalence among pregnant women aged 15-24 in capital city				-
HIV prevalence among most-at-risk populations	FSW:	-	IDU:	-
Knowledge of HIV prevention among young people aged 15-24	M:	-	F:	-
Sex before age 15 among young people aged 15-19	M:	-	F:	-
Condom use at last higher-risk sex among young people aged 15-24	M:	-	F:	-
% of sex workers who used a condom with their most recent client	M:	-	F:	-
% of IDUs who used sterile equipment at last injection				-
% of men who used a condom the last time they had sex with a male partner				-

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS


Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.

* Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Iraq

Estimated adult HIV prevalence rate -

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage <small>Source: MICS, 2006</small>	84%
Estimated number of births per year <small>Source: UN Population Division, 2006</small>	937,000
Estimated number of HIV+ pregnant women	-
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	-

PROVIDE PAEDIATRIC HIV TREATMENT



Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	-

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE


Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24 <small>Source: MICS, 2006</small>	M: -	F: 2.9%
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS


Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio <small>Source: MICS, 2006</small>	0.84
% of orphans receiving external support	-




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
* Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Jamaica

Estimated adult HIV prevalence rate 1.5%
 Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage <small>Source: MICS, 2005</small>	91%
Estimated number of births per year <small>Source: UN Population Division, 2006</small>	55,000
Estimated number of HIV+ pregnant women <small>Source: UNAIDS/WHO unpublished estimates, 2007</small>	<500 [<500 - 636]*
Number and % of HIV+ pregnant women receiving ARVs for PMTCT <small>Source: MOH, 2006</small>	371 -

PROVIDE PAEDIATRIC HIV TREATMENT



Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	- -
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	- -
Estimated number of children in need of ART	-
Number and % of children in need receiving ART <small>Source: MOH, 2006</small>	256 -

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE


Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations <small>Source: UNAIDS, 2006</small>	FSW: 9.0%	IDU: -
Knowledge of HIV prevention among young people aged 15-24 <small>Source: MICS, 2005</small>	M: -	F: 59.8%
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS


Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio <small>Source: MICS, 2005</small>	1.01
% of orphans receiving external support	-




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Jordan

Estimated adult HIV prevalence rate -
 Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	99%	
<small>Source: DHS, 2002</small>		
Estimated number of births per year	152,000	
<small>Source: UN Population Division, 2006</small>		
Estimated number of HIV+ pregnant women	-	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	0	-
<small>Source: MOH, 2006</small>		

PROVIDE PAEDIATRIC HIV TREATMENT






Estimated number of children living with HIV	-	
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	0	-
<small>Source: MOH, 2006</small>		
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	1	-
<small>Source: MOH, 2006</small>		
Estimated number of children in need of ART	-	
Number and % of children in need receiving ART	3	-
<small>Source: MOH, 2006</small>		

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M:	-	F:	-
HIV prevalence among pregnant women aged 15-24 in capital city				-
HIV prevalence among most-at-risk populations	FSW:	-	IDU:	-
Knowledge of HIV prevention among young people aged 15-24	M:	-	F:	3.0%
<small>Source: PFHS, 2002</small>				
Sex before age 15 among young people aged 15-19	M:	-	F:	-
Condom use at last higher-risk sex among young people aged 15-24	M:	-	F:	-
% of sex workers who used a condom with their most recent client	M:	-	F:	-
% of IDUs who used sterile equipment at last injection				-
% of men who used a condom the last time they had sex with a male partner				-

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Kazakhstan

Estimated adult HIV prevalence rate 0.1%
 Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	100%
Source: MICS, 2006	
Estimated number of births per year	289,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	<500 [<200 - 566]*
Source: UNAIDS/WHO unpublished estimates, 2007	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	80 -
Source: MOH, 2006	

PROVIDE PAEDIATRIC HIV TREATMENT



Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	85 -
Source: MOH, 2006	
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	85 -
Source: MOH, 2006	
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	25 -
Source: MOH, 2006	

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE


Estimated HIV prevalence among young people aged 15-24	M: - F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-
HIV prevalence among most-at-risk populations	FSW: - IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: - F: 22.4%
Source: MICS, 2006	
Sex before age 15 among young people aged 15-19	M: - F: -
Condom use at last higher-risk sex among young people aged 15-24	M: - F: -
% of sex workers who used a condom with their most recent client	M: - F: -
% of IDUs who used sterile equipment at last injection	-
% of men who used a condom the last time they had sex with a male partner	-

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS


Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	0.98
Source: MICS, 2006	
% of orphans receiving external support	-




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Kenya

Estimated adult HIV prevalence rate 6.7%
 Source: DHS, 2003

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	88%
Source: DHS, 2003	
Estimated number of births per year	1,447,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	77,799
Source: UNAIDS/WHO unpublished estimates, 2007 [66,593 - 88,223]*	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	37,130 47.7%
Source: MOH, 2006 [42.1% - 55.8%]*	

PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	10,000
Source: MOH, 2006	


PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: 1.3%	F: 5.9%
Source: DHS, 2003		
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: 25.5%	IDU: -
Source: UNAIDS, 2006		
Knowledge of HIV prevention among young people aged 15-24	M: 47.0%	F: 34.0%
Source: DHS, 2003		
Sex before age 15 among young people aged 15-19	M: 30.9%	F: 14.5%
Source: DHS, 2003		
Condom use at last higher-risk sex among young people aged 15-24	M: 47.0%	F: 25.0%
Source: DHS, 2003		
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-


PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	0.95
Source: DHS, 2003	
% of orphans receiving external support	10.3%
Source: UNAIDS, 2006	


Number of HIV+ pregnant women receiving ARVs for PMTCT




Number of children in need receiving ART




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.

* Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Kiribati

Estimated adult HIV prevalence rate -

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	-
Estimated number of births per year <small>Source: UN Population Division, 2006</small>	0
Estimated number of HIV+ pregnant women	-
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	-

PROVIDE PAEDIATRIC HIV TREATMENT






Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	-

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: -	F: -
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-




M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
* Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Korea, Democratic People's Republic of

Estimated adult HIV prevalence rate -


PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	-
Estimated number of births per year	321,000
<small>Source: UN Population Division, 2006</small>	
Estimated number of HIV+ pregnant women	<100
<small>Source: UNAIDS/WHO unpublished estimates, 2007</small>	[<100 - <200]*
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	-





PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	-




PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: -	F: -
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-



PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Korea, Republic of

Estimated adult HIV prevalence rate

<0.1%

Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	-	-
Estimated number of births per year	455,000	
<small>Source: UN Population Division, 2006</small>		
Estimated number of HIV+ pregnant women	<100	[<100 - <500]*
<small>Source: UNAIDS/WHO unpublished estimates, 2007</small>		
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	-	-

PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-	-
Estimated number of children in need of ART	-	-
Number and % of children in need receiving ART	-	-


PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: -	F: -
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-


PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-


Number of HIV+ pregnant women receiving ARVs for PMTCT




Number of children in need receiving ART




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.


* Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Kuwait

Estimated adult HIV prevalence rate -


PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	-
Estimated number of births per year	50,000
<small>Source: UN Population Division, 2006</small>	
Estimated number of HIV+ pregnant women	-
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	-





PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	-




PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: -	F: -
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-



PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Kyrgyzstan

Estimated adult HIV prevalence rate 0.1%
 Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	97%
Source: MICS, 2006	
Estimated number of births per year	113,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	<100
Source: UNAIDS/WHO unpublished estimates, 2007	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	9 -
Source: MOH, 2006	

PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	- -
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	- -
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	3 -
Source: MOH, 2006	


PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	
HIV prevalence among most-at-risk populations	FSW: 1.7%	IDU: 6.2%
Source: UNAIDS, 2006		
Knowledge of HIV prevention among young people aged 15-24	M: -	F: 20.3%
Source: MICS, 2006		
Sex before age 15 among young people aged 15-19	M: -	F: 0.2%
Source: MICS, 2006		
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: 56.0%
Source: MICS, 2006		
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	
% of men who used a condom the last time they had sex with a male partner	68.3%	
Source: UNAIDS, 2006		


PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	1.03
Source: MICS, 2006	
% of orphans receiving external support	-


Number of HIV+ pregnant women receiving ARVs for PMTCT




Number of children in need receiving ART




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates


Children and AIDS fact sheet: Lao People's Democratic Republic

Estimated adult HIV prevalence rate 0.1%
 Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	27%
Source: MICS, 2001	
Estimated number of births per year	156,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	<200
Source: UNAIDS/WHO unpublished estimates, 2007	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	-


Number of HIV+ pregnant women receiving ARVs for PMTCT



PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	3
Source: MOH, 2006	


Number of children in need receiving ART




PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: 1.1%	IDU: -
Source: UNAIDS, 2006		
Knowledge of HIV prevention among young people aged 15-24	M: -	F: -
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: 58.8%	F: 88.9%
Source: UNAIDS, 2006		
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-

% of young people aged 15-24 who used a condom at last higher-risk sex




HIV prevalence among female sex workers (%)



PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-

Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.

* Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Lebanon

Estimated adult HIV prevalence rate 0.1%
 Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	96%
Source: PAPFAM, 2002	
Estimated number of births per year	74,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	<100
Source: UNAIDS/WHO unpublished estimates, 2007	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	- -
[<100 - <500]*	

PROVIDE PAEDIATRIC HIV TREATMENT






Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	- -
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	- -
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	- -

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: -	F: -
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Lesotho

Estimated adult HIV prevalence rate 23.5%
 Source: DHS, 2004

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	90%
Source: DHS, 2004	
Estimated number of births per year	59,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	11,985 [10,902 - 13,292]*
Source: UNAIDS/WHO unpublished estimates, 2007	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	2,005 16.7% [15.1% - 18.4%]*
Source: MOH, 2006	

PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	1,143
Source: MOH, 2006	


PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: 6.0%	F: 15.4%
Source: DHS, 2004		
HIV prevalence among pregnant women aged 15-24 in capital city	27.3%	
Source: UNAIDS, 2006		
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: 18.4%	F: 25.8%
Source: DHS, 2004		
Sex before age 15 among young people aged 15-19	M: 29.7%	F: 16.3%
Source: DHS, 2004		
Condom use at last higher-risk sex among young people aged 15-24	M: 52.8%	F: 53.0%
Source: DHS, 2004		
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	
% of men who used a condom the last time they had sex with a male partner	-	


PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	0.95
Source: DHS, 2004	
% of orphans receiving external support	25.0%
Source: UNAIDS, 2004	


Number of HIV+ pregnant women receiving ARVs for PMTCT




Number of children in need receiving ART




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Liberia

Estimated adult HIV prevalence rate -

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	85%
<small>Source: National Nutrition Survey, 1999-2000</small>	
Estimated number of births per year	184,000
<small>Source: UN Population Division, 2006</small>	
Estimated number of HIV+ pregnant women	2,830 [2,188 - 3,481]*
<small>Source: UNAIDS/WHO unpublished estimates, 2007</small>	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	76 2.7% [2.2% - 3.5%]*
<small>Source: MOH, 2006</small>	

PROVIDE PAEDIATRIC HIV TREATMENT



Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	0
<small>Source: MOH, 2006</small>	
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	-

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE


Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: -	F: -
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS


Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.


* Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Libyan Arab Jamahiriya

Estimated adult HIV prevalence rate -


PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	-
Estimated number of births per year <small>Source: UN Population Division, 2006</small>	144,000
Estimated number of HIV+ pregnant women	-
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	-





PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	-




PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: -	F: -
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-



PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
* Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Madagascar

Estimated adult HIV prevalence rate 0.5%
 Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage Source: DHS, 2003-2004	80%	
Estimated number of births per year Source: UN Population Division, 2006	714,000	
Estimated number of HIV+ pregnant women Source: UNAIDS/WHO unpublished estimates, 2007	<500	[<500 - 904]*
Number and % of HIV+ pregnant women receiving ARVs for PMTCT Source: MOH, 2006	7	-

PROVIDE PAEDIATRIC HIV TREATMENT






Estimated number of children living with HIV	-	
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth Source: MOH, 2006	2	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth Source: MOH, 2006	2	-
Estimated number of children in need of ART	-	
Number and % of children in need receiving ART Source: MOH, 2006	0	-

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24 Source: UNAIDS, 2006	M: 0.6%	F: 0.3%
HIV prevalence among pregnant women aged 15-24 in capital city	-	
HIV prevalence among most-at-risk populations Source: UNAIDS, 2006	FSW: 0.2%	IDU: -
Knowledge of HIV prevention among young people aged 15-24 Source: DHS, 2003-2004	M: 16.0%	F: 19.0%
Sex before age 15 among young people aged 15-19 Source: DHS, 2003-2004	M: 19.5%	F: 31.9%
Condom use at last higher-risk sex among young people aged 15-24 Source: DHS, 2003-2004	M: 12.0%	F: 5.0%
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	
% of men who used a condom the last time they had sex with a male partner	-	

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio Source: DHS, 2003-2004	0.75
% of orphans receiving external support	-



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Malawi

Estimated adult HIV prevalence rate 12.7%
Source: DHS, 2004

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	92%
Source: MICS, 2006	
Estimated number of births per year	566,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	67,316
Source: UNAIDS/WHO unpublished estimates, 2007 [58,121 - 76,957]*	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	9,231 13.7%
Source: MOH, 2006 [12.0% - 15.9%]*	

PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	60
Source: MOH, 2006	
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	5,763
Source: MOH, 2006	


PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: 2.1%	F: 9.1%
Source: DHS, 2004		
HIV prevalence among pregnant women aged 15-24 in capital city	15.0%	
Source: UNAIDS, 2006		
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: -	F: 42.1%
Source: MICS, 2006		
Sex before age 15 among young people aged 15-19	M: -	F: 13.9%
Source: MICS, 2006		
Condom use at last higher-risk sex among young people aged 15-24	M: 59.6%	F: 39.6%
Source: MICS, 2006		
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	
% of men who used a condom the last time they had sex with a male partner	-	


PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	0.97
Source: MICS, 2006	
% of orphans receiving external support	18.5%
Source: MICS, 2006	


Number of HIV+ pregnant women receiving ARVs for PMTCT




Number of children in need receiving ART




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.

* Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Malaysia

Estimated adult HIV prevalence rate 0.5%
 Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	79%
Source: MOH, 2005	
Estimated number of births per year	556,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	1,159 [737 - 1,813]*
Source: UNAIDS/WHO unpublished estimates, 2007	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	170 14.7% [9.4% - 23.1%]*
Source: MOH, 2006	

PROVIDE PAEDIATRIC HIV TREATMENT



Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	62
Source: MOH, 2006	
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	510
Source: MOH, 2006	

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE


Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	
HIV prevalence among most-at-risk populations	FSW: 6.9%	IDU: -
Source: UNAIDS, 2006		
Knowledge of HIV prevention among young people aged 15-24	M: -	F: -
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	
% of men who used a condom the last time they had sex with a male partner	-	

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS


Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Maldives

Estimated adult HIV prevalence rate -

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage <small>Source: MICS, 2001</small>	81%
Estimated number of births per year <small>Source: UN Population Division, 2006</small>	7,000
Estimated number of HIV+ pregnant women <small>Source: UNAIDS/WHO unpublished estimates, 2007</small>	<100 [<100]*
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	- -

PROVIDE PAEDIATRIC HIV TREATMENT



Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	- -
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	- -
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	- -

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE


Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: -	F: -
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS


Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
* Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Mali

Estimated adult HIV prevalence rate 1.3%
 Source: UNAIDS/WHO, 2007

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	57%
Source: DHS, 2001	
Estimated number of births per year	579,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	8,426 [6,893 - 10,346]*
Source: UNAIDS/WHO unpublished estimates, 2007	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	656 7.8% [6.3% - 9.5%]*
Source: MOH, 2006	

PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	691
Source: MOH, 2006	


PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: 0.4%	F: 1.2%
Source: UNAIDS, 2006		
HIV prevalence among pregnant women aged 15-24 in capital city	2.5%	
Source: SOWC, 2002		
HIV prevalence among most-at-risk populations	FSW: 31.6%	IDU: -
Source: UNAIDS, 2006		
Knowledge of HIV prevention among young people aged 15-24	M: 15.0%	F: 9.0%
Source: DHS, 2001		
Sex before age 15 among young people aged 15-19	M: 10.6%	F: 26.0%
Source: DHS, 2001		
Condom use at last higher-risk sex among young people aged 15-24	M: 30.0%	F: 14.0%
Source: DHS, 2001		
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	
% of men who used a condom the last time they had sex with a male partner	-	


PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	1.04
Source: DHS, 2001	
% of orphans receiving external support	-

Number of HIV+ pregnant women receiving ARVs for PMTCT




Number of children in need receiving ART




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Marshall Islands

Estimated adult HIV prevalence rate -

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	-
Estimated number of births per year <small>Source: UN Population Division, 2006</small>	0
Estimated number of HIV+ pregnant women	-
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	-

PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	-


PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: -	F: -
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-


PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-


Number of HIV+ pregnant women receiving ARVs for PMTCT




Number of children in need receiving ART




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
* Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Mauritania

Estimated adult HIV prevalence rate 0.7%
 Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	64%	
<small>Source: DHS, 2000-2001</small>		
Estimated number of births per year	102,000	
<small>Source: UN Population Division, 2006</small>		
Estimated number of HIV+ pregnant women	<500	[<500 - 787]*
<small>Source: UNAIDS/WHO unpublished estimates, 2007</small>		
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	12	-
<small>Source: MOH, 2006</small>		

PROVIDE PAEDIATRIC HIV TREATMENT






Estimated number of children living with HIV	-	
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	14	-
<small>Source: MOH, 2006</small>		
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	0	-
<small>Source: MOH, 2006</small>		
Estimated number of children in need of ART	-	
Number and % of children in need receiving ART	7	-
<small>Source: MOH, 2006</small>		

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: 0.2%	F: 0.5%
<small>Source: UNAIDS, 2006</small>		
HIV prevalence among pregnant women aged 15-24 in capital city	-	
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: -	F: -
Sex before age 15 among young people aged 15-19	M: 2.1%	F: 13.2%
<small>Source: DHS, 2000-2001</small>		
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	
% of men who used a condom the last time they had sex with a male partner	-	

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Mauritius

Estimated adult HIV prevalence rate


0.6%

Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	-	-
Estimated number of births per year	19,000	
<small>Source: UN Population Division, 2006</small>		
Estimated number of HIV+ pregnant women	<200	[<100 - <500]*
<small>Source: UNAIDS/WHO unpublished estimates, 2007</small>		
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	-	-


Number of HIV+ pregnant women receiving ARVs for PMTCT



PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-	-
Estimated number of children in need of ART	-	-
Number and % of children in need receiving ART	-	-


Number of children in need receiving ART




PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: -	F: -
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-

% of young people aged 15-24 who used a condom at last higher-risk sex




HIV prevalence among female sex workers (%)



PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-

Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.

* Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Mexico

Estimated adult HIV prevalence rate

0.3%

Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	-
Estimated number of births per year	2,109,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	3,073
Source: UNAIDS/WHO unpublished estimates, 2007	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	146 4.8%
Source: MOH, 2006	
	[2.5% - 8.8%]*

PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	176
Source: MOH, 2006	
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	176
Source: MOH, 2006	


PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M:	-	F:	-
HIV prevalence among pregnant women aged 15-24 in capital city				-
HIV prevalence among most-at-risk populations	FSW:	-	IDU:	-
Knowledge of HIV prevention among young people aged 15-24	M:	-	F:	-
Sex before age 15 among young people aged 15-19	M:	-	F:	-
Condom use at last higher-risk sex among young people aged 15-24	M:	-	F:	-
% of sex workers who used a condom with their most recent client	M:	-	F:	-
% of IDUs who used sterile equipment at last injection				-
% of men who used a condom the last time they had sex with a male partner				-


PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-

Number of HIV+ pregnant women receiving ARVs for PMTCT




Number of children in need receiving ART




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.


* Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Micronesia (Federated States of)

Estimated adult HIV prevalence rate -


PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	-
Estimated number of births per year	3,000
<small>Source: UN Population Division, 2006</small>	
Estimated number of HIV+ pregnant women	-
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	-





PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	-




PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: -	F: -
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-



PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.

* Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Moldova

Estimated adult HIV prevalence rate 1.1%
 Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage Source: DHS, 2005	98%	
Estimated number of births per year Source: UN Population Division, 2006	44,000	
Estimated number of HIV+ pregnant women Source: UNAIDS/WHO unpublished estimates, 2007	<100	[<100 - <200]*
Number and % of HIV+ pregnant women receiving ARVs for PMTCT Source: MOH, 2006	62	-

PROVIDE PAEDIATRIC HIV TREATMENT



Estimated number of children living with HIV	-	
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth Source: MOH, 2006	0	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth Source: MOH, 2006	60	-
Estimated number of children in need of ART	-	
Number and % of children in need receiving ART Source: MOH, 2006	14	-

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE


Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city		-
HIV prevalence among most-at-risk populations Source: BSS, 2004	FSW: 4.4%	IDU: -
Knowledge of HIV prevention among young people aged 15-24 Source: DHS, 2005	M: 54.3%	F: 41.6%
Sex before age 15 among young people aged 15-19 Source: DHS, 2005	M: 10.1%	F: 1.7%
Condom use at last higher-risk sex among young people aged 15-24 Source: DHS, 2005	M: 63.0%	F: 44.0%
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection Source: BSS, 2004		81.0%
% of men who used a condom the last time they had sex with a male partner Source: BSS, 2004		63.0%

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS


Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio Source: DHS, 2005	0.90
% of orphans receiving external support	-




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Mongolia

Estimated adult HIV prevalence rate <0.1%
Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage Source: MICS, 2005-2006	99%
Estimated number of births per year Source: UN Population Division, 2006	49,000
Estimated number of HIV+ pregnant women Source: UNAIDS/WHO unpublished estimates, 2007	<100 [<100]*
Number and % of HIV+ pregnant women receiving ARVs for PMTCT Source: MOH, 2006	1 -

PROVIDE PAEDIATRIC HIV TREATMENT



Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth Source: MOH, 2006	0 -
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth Source: MOH, 2006	0 -
Estimated number of children in need of ART	-
Number and % of children in need receiving ART Source: MOH, 2006	0 -

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE


Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24 Source: MICS, 2005	M: -	F: 34.5%
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client Source: UNAIDS, 2006	M: 50.0%	F: 96.4%
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner Source: UNAIDS, 2006	-	13.0%

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS


Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio Source: MICS, 2005	0.96
% of orphans receiving external support	-




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.

* Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Montenegro

Estimated adult HIV prevalence rate -

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage <small>Source: MICS, 2005</small>	97%
Estimated number of births per year <small>Source: UN Population Division, 2006</small>	8,000
Estimated number of HIV+ pregnant women	-
Number and % of HIV+ pregnant women receiving ARVs for PMTCT <small>Source: MOH, 2006</small>	0 -

PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	- -
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	- -
Estimated number of children in need of ART	-
Number and % of children in need receiving ART <small>Source: MOH, 2006</small>	2 -


PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24 <small>Source: MICS, 2006</small>	M: -	F: 29.6%
Sex before age 15 among young people aged 15-19 <small>Source: MICS, 2006</small>	M: -	F: 0.4%
Condom use at last higher-risk sex among young people aged 15-24 <small>Source: MICS, 2006</small>	M: -	F: 66.4%
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-


PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio <small>Source: MICS, 2005</small>	1.00
% of orphans receiving external support	-


Number of HIV+ pregnant women receiving ARVs for PMTCT




Number of children in need receiving ART




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
* Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Morocco

Estimated adult HIV prevalence rate 0.1%
 Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage <small>Source: DHS, 2003-2004</small>	68%
Estimated number of births per year <small>Source: UN Population Division, 2006</small>	635,000
Estimated number of HIV+ pregnant women <small>Source: UNAIDS/WHO unpublished estimates, 2007</small>	<500 [<200 - 585]*
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	- -

PROVIDE PAEDIATRIC HIV TREATMENT



Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	- -
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	- -
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	- -

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE


Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations <small>Source: UNAIDS, 2006</small>	FSW: 1.9%	IDU: -
Knowledge of HIV prevention among young people aged 15-24 <small>Source: DHS, 2003-2004</small>	M: -	F: 12.0%
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client <small>Source: UNAIDS, 2006</small>	M: -	F: 37.8%
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS


Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Mozambique

Estimated adult HIV prevalence rate 16.1%
 Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	85%
Source: DHS, 2003	
Estimated number of births per year	856,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	95,482 [79,656 - 112,694]*
Source: UNAIDS/WHO unpublished estimates, 2007	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	12,150 12.7% [10.8% - 15.3%]*
Source: MOH, 2006	

PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	3,443
Source: MOH, 2006	


PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: 3.6%	F: 10.7%
Source: UNAIDS, 2006		
HIV prevalence among pregnant women aged 15-24 in capital city	14.7%	
Source: SOWC, 2002		
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: 33.0%	F: 20.0%
Source: DHS, 2003		
Sex before age 15 among young people aged 15-19	M: 31.3%	F: 27.8%
Source: DHS, 2003		
Condom use at last higher-risk sex among young people aged 15-24	M: 33.0%	F: 29.0%
Source: DHS, 2003		
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	
% of men who used a condom the last time they had sex with a male partner	-	


PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	0.80
Source: DHS, 2003	
% of orphans receiving external support	-


Number of HIV+ pregnant women receiving ARVs for PMTCT




Number of children in need receiving ART




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Myanmar

Estimated adult HIV prevalence rate 1.3%
 Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	76%
Source: FRHS, 2001	
Estimated number of births per year	897,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	2,907 [2,077 - 3,952]*
Source: UNAIDS/WHO unpublished estimates, 2007	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	993 34.2% [25.1% - 47.8%]*
Source: MOH, 2006	

PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	287
Source: MOH, 2006	


PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: -	F: -
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-


PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-


Number of HIV+ pregnant women receiving ARVs for PMTCT




Number of children in need receiving ART




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Namibia

Estimated adult HIV prevalence rate 19.6%
 Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	91%
Source: DHS, 2000	
Estimated number of births per year	53,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	9,316 [7,615 - 11,151]*
Source: UNAIDS/WHO unpublished estimates, 2007	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	6,022 64.6% [54.0% - 79.1%]*
Source: MOH, 2006	

PROVIDE PAEDIATRIC HIV TREATMENT






Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	3,932
Source: MOH, 2006	

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: 4.4%	F: 13.4%
Source: UNAIDS, 2006		
HIV prevalence among pregnant women aged 15-24 in capital city	7.5%	
Source: UNAIDS, 2006		
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: 41.0%	F: 31.0%
Source: DHS, 2000		
Sex before age 15 among young people aged 15-19	M: 31.3%	F: 9.8%
Source: DHS, 2000		
Condom use at last higher-risk sex among young people aged 15-24	M: 69.0%	F: 48.0%
Source: DHS, 2000		
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	
% of men who used a condom the last time they had sex with a male partner	-	

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	0.92
Source: DHS, 2000	
% of orphans receiving external support	16.5%
Source: DHS, 2006	



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Nauru

Estimated adult HIV prevalence rate -

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	-
Estimated number of births per year <small>Source: UN Population Division, 2006</small>	0
Estimated number of HIV+ pregnant women	-
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	-

PROVIDE PAEDIATRIC HIV TREATMENT






Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	-

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: -	F: -
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
* Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Nepal

Estimated adult HIV prevalence rate 0.5%
 Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	44%
Source: DHS, 2006	
Estimated number of births per year	791,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	1,738 [919 - 3,083]*
Source: UNAIDS/WHO unpublished estimates, 2007	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	35 2.0% [1.1% - 3.8%]*
Source: MOH, 2006	

PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	9 -
Source: MOH, 2006	
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	0 -
Source: MOH, 2006	
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	12 -
Source: MOH, 2006	


PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	
HIV prevalence among most-at-risk populations	FSW: 2.0%	IDU: 50.0%
Source: UNAIDS, 2006		
Knowledge of HIV prevention among young people aged 15-24	M: 43.6%	F: 27.6%
Source: DHS, 2006		
Sex before age 15 among young people aged 15-19	M: 3.1%	F: 5.5%
Source: DHS, 2006		
Condom use at last higher-risk sex among young people aged 15-24	M: 78.4%	F: -
Source: DHS, 2006		
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	
% of men who used a condom the last time they had sex with a male partner	-	


PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-


Number of HIV+ pregnant women receiving ARVs for PMTCT




Number of children in need receiving ART




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Nicaragua

Estimated adult HIV prevalence rate

0.2%

Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	86%
Source: DHS, 2001	
Estimated number of births per year	139,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	<200
Source: UNAIDS/WHO unpublished estimates, 2007	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	26 -
Source: MOH, 2006	

PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	26 -
Source: MOH, 2006	
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	3 -
Source: MOH, 2006	
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	26 -
Source: MOH, 2006	


PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: 0.9%	IDU: -
Source: UNAIDS, 2006		
Knowledge of HIV prevention among young people aged 15-24	M: -	F: 22.0%
Source: DHS, 2001		
Sex before age 15 among young people aged 15-19	M: -	F: 10.9%
Source: DHS, 2001		
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: 17.0%
Source: DHS, 2001		
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-


PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	1.06
Source: DHS, 2001	
% of orphans receiving external support	-


Number of HIV+ pregnant women receiving ARVs for PMTCT




Number of children in need receiving ART




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.

* Represents the uncertainty range around the estimates

Rev. 7 March 2008

Children and AIDS fact sheet: Niger

Estimated adult HIV prevalence rate 0.7%
Source: DHS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	46%
<small>Source: DHS, 2006</small>	
Estimated number of births per year	683,000
<small>Source: UN Population Division, 2006</small>	
Estimated number of HIV+ pregnant women	3,132
<small>Source: UNAIDS/WHO unpublished estimates, 2007</small>	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	214 6.8%
<small>Source: MOH, 2006</small>	

PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	46
<small>Source: MOH, 2006</small>	


PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: 0.1%	F: 0.5%
<small>Source: DHS, 2006</small>		
HIV prevalence among pregnant women aged 15-24 in capital city	-	
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: 15.9%	F: 13.4%
<small>Source: DHS, 2006</small>		
Sex before age 15 among young people aged 15-19	M: 5.3%	F: 25.7%
<small>Source: DHS, 2006</small>		
Condom use at last higher-risk sex among young people aged 15-24	M: 37.2%	F: 17.8%
<small>Source: DHS, 2006</small>		
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	
% of men who used a condom the last time they had sex with a male partner	-	


PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	0.67
<small>Source: DHS, 2006</small>	
% of orphans receiving external support	-


Number of HIV+ pregnant women receiving ARVs for PMTCT




Number of children in need receiving ART




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Year refers to the year of data, not publication date. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Nigeria

Estimated adult HIV prevalence rate 3.9%
 Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	58%
Source: DHS, 2003	
Estimated number of births per year	5,909,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	187,544
Source: UNAIDS/WHO unpublished estimates, 2007 [133,734 - 231,696]*	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	6,168 3.3%
Source: MOH, 2006 [2.7% - 4.6%]*	

PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	5,279
Source: MOH, 2006	


PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: 0.9%	F: 2.7%
Source: UNAIDS, 2006		
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: 21.0%	F: 18.0%
Source: DHS, 2003		
Sex before age 15 among young people aged 15-19	M: 7.9%	F: 20.3%
Source: DHS, 2003		
Condom use at last higher-risk sex among young people aged 15-24	M: 46.0%	F: 24.0%
Source: DHS, 2003		
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-


PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	0.64
Source: DHS, 2003	
% of orphans receiving external support	-


Number of HIV+ pregnant women receiving ARVs for PMTCT




Number of children in need receiving ART




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Niue

Estimated adult HIV prevalence rate -

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	-
Estimated number of births per year <small>Source: UN Population Division, 2006</small>	0
Estimated number of HIV+ pregnant women	-
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	-

PROVIDE PAEDIATRIC HIV TREATMENT






Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	-

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: -	F: -
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
* Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Occupied Palestinian Territory

Estimated adult HIV prevalence rate -

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage <small>Source: FHS, 2006</small>	99%
Estimated number of births per year <small>Source: UN Population Division, 2006</small>	143,000
Estimated number of HIV+ pregnant women	-
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	-

PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	-


PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: -	F: -
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-


PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-


Number of HIV+ pregnant women receiving ARVs for PMTCT




Number of children in need receiving ART




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
* Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Oman

Estimated adult HIV prevalence rate -

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage <small>Source: NHS, Part II, 2000</small>	100%
Estimated number of births per year <small>Source: UN Population Division, 2006</small>	57,000
Estimated number of HIV+ pregnant women	-
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	-

PROVIDE PAEDIATRIC HIV TREATMENT






Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-
Estimated number of children in need of ART	-
Number and % of children in need receiving ART <small>Source: MOH, 2006</small>	25

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: -	F: -
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
* Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Pakistan

Estimated adult HIV prevalence rate 0.1%
 Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	36%
Source: PSLM, 2004-2005	
Estimated number of births per year	4,358,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	2,089 [1,137 - 4,265]*
Source: UNAIDS/WHO unpublished estimates, 2007	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	8 0.4% [0.2% - 0.7%]*
Source: MOH, 2006	

PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	0 -
Source: MOH, 2006	
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	4 -
Source: MOH, 2006	
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	10 -
Source: MOH, 2006	


PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	
HIV prevalence among most-at-risk populations	FSW: -	IDU: 22.9%
Source: UNAIDS, 2006		
Knowledge of HIV prevention among young people aged 15-24	M: -	F: -
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: 6.6%	F: 37.0%
Source: UNAIDS, 2006		
% of IDUs who used sterile equipment at last injection	-	
% of men who used a condom the last time they had sex with a male partner	7.6%	
Source: UNAIDS, 2006		


PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-


Number of HIV+ pregnant women receiving ARVs for PMTCT




Number of children in need receiving ART




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Palau

Estimated adult HIV prevalence rate -

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	-
Estimated number of births per year <small>Source: UN Population Division, 2006</small>	0
Estimated number of HIV+ pregnant women	-
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	-

PROVIDE PAEDIATRIC HIV TREATMENT



Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	-

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE


Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: -	F: -
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS


Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
* Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Panama

Estimated adult HIV prevalence rate -
Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	-	-
Estimated number of births per year	70,000	
<small>Source: UN Population Division, 2006</small>		
Estimated number of HIV+ pregnant women	<500	[<500 - 513]*
<small>Source: UNAIDS/WHO unpublished estimates, 2007</small>		
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	-	-

PROVIDE PAEDIATRIC HIV TREATMENT






Estimated number of children living with HIV	-	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-	-
Estimated number of children in need of ART	-	-
Number and % of children in need receiving ART	214	-
<small>Source: MOH, 2006</small>		

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: 1.8%	IDU: -
<small>Source: UNAIDS, 2006</small>		
Knowledge of HIV prevention among young people aged 15-24	M: -	F: -
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: 90.8%	F: 91.6%
<small>Source: UNAIDS, 2006</small>		
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	84.2%
<small>Source: UNAIDS, 2006</small>		

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Papua New Guinea

Estimated adult HIV prevalence rate 1.8%
 Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	-
Estimated number of births per year	191,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	1,546 [1,430 - 1,672]*
Source: UNAIDS/WHO unpublished estimates, 2007	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	31 -
Source: MOH, 2006	

PROVIDE PAEDIATRIC HIV TREATMENT



Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	- -
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	0 -
Source: MOH, 2006	
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	23 -
Source: MOH, 2006	

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE


Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: 16.0%	IDU: -
Source: UNAIDS, 2006		
Knowledge of HIV prevention among young people aged 15-24	M: -	F: -
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS


Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Paraguay

Estimated adult HIV prevalence rate 0.4%
Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	94%	
<small>Source: Encuesta Nacional de Demografía y Salud Sexual y Reproductiva, 2004</small>		
Estimated number of births per year	153,000	
<small>Source: UN Population Division, 2006</small>		
Estimated number of HIV+ pregnant women	<500	[<500 - 769]*
<small>Source: UNAIDS/WHO unpublished estimates, 2007</small>		
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	73	-
<small>Source: MOH, 2006</small>		

PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-	
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	42	-
<small>Source: MOH, 2006</small>		
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	54	-
<small>Source: MOH, 2006</small>		
Estimated number of children in need of ART	-	
Number and % of children in need receiving ART	160	-
<small>Source: MOH, 2006</small>		


PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M:	-	F:	-
HIV prevalence among pregnant women aged 15-24 in capital city				-
HIV prevalence among most-at-risk populations	FSW:	3.1%	IDU:	-
<small>Source: UNAIDS, 2006</small>				
Knowledge of HIV prevention among young people aged 15-24	M:	-	F:	-
Sex before age 15 among young people aged 15-19	M:	-	F:	-
Condom use at last higher-risk sex among young people aged 15-24	M:	-	F:	-
% of sex workers who used a condom with their most recent client	M:	-	F:	-
% of IDUs who used sterile equipment at last injection				-
% of men who used a condom the last time they had sex with a male partner				-


PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-


Number of HIV+ pregnant women receiving ARVs for PMTCT




Number of children in need receiving ART




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
* Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Peru

Estimated adult HIV prevalence rate 0.6%
 Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	92%
Source: DHS, 2004	
Estimated number of births per year	584,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	1,653 [902 - 2,685]*
Source: UNAIDS/WHO unpublished estimates, 2007	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	560 33.9% [20.9% - 62.1%]*
Source: MOH, 2006	

PROVIDE PAEDIATRIC HIV TREATMENT






Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	280
Source: MOH, 2006	

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: -	F: -
Sex before age 15 among young people aged 15-19	M: -	F: 5.1%
Source: DHS, 2000		
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: 31.6%
Source: DHS, 2004		
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	46.3%
Source: UNAIDS, 2006		

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	0.85
Source: DHS, 2000	
% of orphans receiving external support	-



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Philippines

Estimated adult HIV prevalence rate

<0.1%

Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	88%
Source: DHS, 2003	
Estimated number of births per year	2,295,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	<500
Source: UNAIDS/WHO unpublished estimates, 2007	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	-

PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	-


PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: 1.0%
Source: UNAIDS, 2006		
Knowledge of HIV prevention among young people aged 15-24	M: 18.0%	F: 12.0%
Source: DHS, 2003		
Sex before age 15 among young people aged 15-19	M: 3.0%	F: 1.0%
Source: DHS, 2003		
Condom use at last higher-risk sex among young people aged 15-24	M: 25.0%	F: 11.0%
Source: DHS, 2003		
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-


PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-


Number of HIV+ pregnant women receiving ARVs for PMTCT




Number of children in need receiving ART




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.

* Represents the uncertainty range around the estimates

Rev. 7 March 2008

Children and AIDS fact sheet: Qatar

Estimated adult HIV prevalence rate -

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	-
Estimated number of births per year	13,000
<small>Source: UN Population Division, 2006</small>	
Estimated number of HIV+ pregnant women	-
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	-

PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	-

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: -	F: -
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-


PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-


Number of HIV+ pregnant women receiving ARVs for PMTCT




Number of children in need receiving ART




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Romania

Estimated adult HIV prevalence rate

<0.1%

Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	94%
Source: RHS, 2004	
Estimated number of births per year	213,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	<500
Source: UNAIDS/WHO unpublished estimates, 2007	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	75 -
Source: MOH, 2006	

PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	59 -
Source: MOH, 2006	
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	280 -
Source: MOH, 2006	


PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: 1.4%	IDU: -
Source: UNAIDS, 2006		
Knowledge of HIV prevention among young people aged 15-24	M: 3.0%	F: 6.0%
Source: RHS, 2004		
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: 85.0%
Source: UNAIDS, 2006		
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-


PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-


Number of HIV+ pregnant women receiving ARVs for PMTCT




Number of children in need receiving ART




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.

* Represents the uncertainty range around the estimates

Rev. 7 March 2008

Children and AIDS fact sheet: Russian Federation

Estimated adult HIV prevalence rate 1.1%
 Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	-
Estimated number of births per year	1,506,000
<small>Source: UN Population Division, 2006</small>	
Estimated number of HIV+ pregnant women	7,223 [3,525 - 15,697]*
<small>Source: UNAIDS/WHO unpublished estimates, 2007</small>	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	6,224 86.2% [39.7% - >95.0%]*
<small>Source: MOH, 2006</small>	

PROVIDE PAEDIATRIC HIV TREATMENT



Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	330
<small>Source: MOH, 2006</small>	

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE


Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: 3.1%	IDU: 12.4%
<small>Source: UNAIDS, 2006</small>		
Knowledge of HIV prevention among young people aged 15-24	M: -	F: -
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	38.8%
<small>Source: UNAIDS, 2006</small>		

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS


Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Rwanda

Estimated adult HIV prevalence rate 3.0%
 Source: DHS, 2005

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	94%
Source: DHS, 2005	
Estimated number of births per year	420,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	14,107 [12,522 - 16,777]*
Source: UNAIDS/WHO unpublished estimates, 2007	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	7,768 55.1% [46.3% - 62.0%]*
Source: MOH, 2006	

PROVIDE PAEDIATRIC HIV TREATMENT



Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	1,209 -
Source: MOH, 2006	
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	1,032 -
Source: MOH, 2006	
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	2,757 -
Source: MOH, 2006	

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE


Estimated HIV prevalence among young people aged 15-24	M: 0.4%	F: 1.5%
Source: DHS, 2005		
HIV prevalence among pregnant women aged 15-24 in capital city	10.3%	
Source: SOWC, 2003		
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: 53.6%	F: 50.9%
Source: DHS, 2005		
Sex before age 15 among young people aged 15-19	M: 15.3%	F: 5.2%
Source: DHS, 2005		
Condom use at last higher-risk sex among young people aged 15-24	M: 39.5%	F: 26.4%
Source: DHS, 2005		
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	
% of men who used a condom the last time they had sex with a male partner	-	

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS


Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	0.82
Source: DHS, 2005	
% of orphans receiving external support	-




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Saint Kitts and Nevis

Estimated adult HIV prevalence rate -

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage <small>Source: PAHO/WHO, 2005</small>	100%
Estimated number of births per year <small>Source: UN Population Division, 2006</small>	1,000
Estimated number of HIV+ pregnant women	-
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	-

PROVIDE PAEDIATRIC HIV TREATMENT






Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	-

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: -	F: -
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
* Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Saint Lucia

Estimated adult HIV prevalence rate -

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage <small>Source: PAHO/WHO, 2004</small>	99%
Estimated number of births per year <small>Source: UN Population Division, 2006</small>	3,000
Estimated number of HIV+ pregnant women	-
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	-

PROVIDE PAEDIATRIC HIV TREATMENT






Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	-

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: -	F: -
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
* Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Saint Vincent and the Grenadines

Estimated adult HIV prevalence rate -

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage <small>Source: PAHO/WHO, 2005</small>	95%
Estimated number of births per year <small>Source: UN Population Division, 2006</small>	2,000
Estimated number of HIV+ pregnant women	-
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	-

PROVIDE PAEDIATRIC HIV TREATMENT






Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	-

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: -	F: -
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-




M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
* Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Samoa

Estimated adult HIV prevalence rate -


PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	-
Estimated number of births per year	5,000
<small>Source: UN Population Division, 2006</small>	
Estimated number of HIV+ pregnant women	-
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	-





PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	-




PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: -	F: -
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-



PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Sao Tome and Principe

Estimated adult HIV prevalence rate -

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage <small>Source: MICS, 2006</small>	97%
Estimated number of births per year <small>Source: UN Population Division, 2006</small>	5,000
Estimated number of HIV+ pregnant women	-
Number and % of HIV+ pregnant women receiving ARVs for PMTCT <small>Source: MOH, 2006</small>	11 -

PROVIDE PAEDIATRIC HIV TREATMENT



Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth <small>Source: MOH, 2006</small>	3 -
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	- -
Estimated number of children in need of ART	-
Number and % of children in need receiving ART <small>Source: MOH, 2006</small>	3 -

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE


Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24 <small>Source: MICS, 2006</small>	M: -	F: 43.6%
Sex before age 15 among young people aged 15-19 <small>Source: MICS, 2006</small>	M: -	F: 8.9%
Condom use at last higher-risk sex among young people aged 15-24 <small>Source: MICS, 2006</small>	M: -	F: 56.3%
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS


Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio <small>Source: MICS, 2006</small>	1.09
% of orphans receiving external support	-




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
* Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Saudi Arabia

Estimated adult HIV prevalence rate -

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	-
Estimated number of births per year <small>Source: UN Population Division, 2006</small>	612,000
Estimated number of HIV+ pregnant women	-
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	-

PROVIDE PAEDIATRIC HIV TREATMENT






Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	-

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M:	-	F:	-
HIV prevalence among pregnant women aged 15-24 in capital city				-
HIV prevalence among most-at-risk populations	FSW:	-	IDU:	-
Knowledge of HIV prevention among young people aged 15-24	M:	-	F:	-
Sex before age 15 among young people aged 15-19	M:	-	F:	-
Condom use at last higher-risk sex among young people aged 15-24	M:	-	F:	-
% of sex workers who used a condom with their most recent client	M:	-	F:	-
% of IDUs who used sterile equipment at last injection				-
% of men who used a condom the last time they had sex with a male partner				-

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
* Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Senegal

Estimated adult HIV prevalence rate 0.7%
 Source: DHS, 2005

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	87%
Source: DHS, 2005	
Estimated number of births per year	435,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	3,818
Source: UNAIDS/WHO unpublished estimates, 2007	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	-

PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	228
Source: MOH, 2006	


PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: 0.2%	F: 0.6%
Source: DHS, 2005		
HIV prevalence among pregnant women aged 15-24 in capital city	0.9%	
Source: SOWC, 2005		
HIV prevalence among most-at-risk populations	FSW: 27.1%	IDU: -
Source: UNAIDS, 2006		
Knowledge of HIV prevention among young people aged 15-24	M: 23.7%	F: 19.3%
Source: DHS, 2005		
Sex before age 15 among young people aged 15-19	M: 13.0%	F: 9.0%
Source: DHS, 2005		
Condom use at last higher-risk sex among young people aged 15-24	M: 52.4%	F: 35.6%
Source: DHS, 2005		
% of sex workers who used a condom with their most recent client	M: -	F: 86.6%
Source: UNAIDS, 2006		
% of IDUs who used sterile equipment at last injection	-	
% of men who used a condom the last time they had sex with a male partner	44.5%	
Source: UNAIDS, 2006		


PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	0.83
Source: National Report on the Situation of Children and Women in Senegal, 2005	
% of orphans receiving external support	-


Number of HIV+ pregnant women receiving ARVs for PMTCT




Number of children in need receiving ART




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Serbia

Estimated adult HIV prevalence rate -

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	98%
<small>Source: MICS, 2005</small>	
Estimated number of births per year	126,000
<small>Source: UN Population Division, 2006</small>	
Estimated number of HIV+ pregnant women	-
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	2 -
<small>Source: MOH, 2006</small>	

PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	0 -
<small>Source: MOH, 2006</small>	
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	2 -
<small>Source: MOH, 2006</small>	
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	14 -
<small>Source: MOH, 2006</small>	


PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: -	F: 42.3%
<small>Source: MICS, 2006</small>		
Sex before age 15 among young people aged 15-19	M: -	F: 1.1%
<small>Source: MICS, 2006</small>		
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: 74.4%
<small>Source: MICS, 2006</small>		
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-


PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-


Number of HIV+ pregnant women receiving ARVs for PMTCT




Number of children in need receiving ART




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.


* Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Seychelles

Estimated adult HIV prevalence rate -


PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	-
Estimated number of births per year	3,000
<small>Source: UN Population Division, 2006</small>	
Estimated number of HIV+ pregnant women	-
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	-



PROVIDE PAEDIATRIC HIV TREATMENT


Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	-




PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: -	F: -
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-

% of young people aged 15-24 who used a condom at last higher-risk sex




HIV prevalence among female sex workers (%)



PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.

* Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Sierra Leone

Estimated adult HIV prevalence rate 1.5%
 Source: UNAIDS/WHO, 2007

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage Source: MICS, 2005	81%
Estimated number of births per year Source: UN Population Division, 2006	262,000
Estimated number of HIV+ pregnant women Source: UNAIDS/WHO unpublished estimates, 2007	4,212 [3,233 - 5,535]*
Number and % of HIV+ pregnant women receiving ARVs for PMTCT Source: MOH, 2006	354 8.4% [6.4% - 10.9%]*

PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth Source: MOH, 2006	60 -
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	- -
Estimated number of children in need of ART	-
Number and % of children in need receiving ART Source: MOH, 2006	12 -


PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24 Source: UNAIDS, 2006	M: 0.4%	F: 1.1%
HIV prevalence among pregnant women aged 15-24 in capital city Source: SOWC, 2003	3.2%	
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24 Source: MICS, 2005	M: -	F: 17.1%
Sex before age 15 among young people aged 15-19 Source: MICS, 2005	M: -	F: 24.9%
Condom use at last higher-risk sex among young people aged 15-24 Source: MICS, 2005	M: -	F: 20.4%
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	
% of men who used a condom the last time they had sex with a male partner	-	


PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio Source: MICS, 2005	0.83
% of orphans receiving external support	-


Number of HIV+ pregnant women receiving ARVs for PMTCT




Number of children in need receiving ART




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Singapore

Estimated adult HIV prevalence rate

0.3%

Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	-	-
Estimated number of births per year	37,000	-
<small>Source: UN Population Division, 2006</small>		
Estimated number of HIV+ pregnant women	-	-
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	-	-

PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-	-
Estimated number of children in need of ART	-	-
Number and % of children in need receiving ART	-	-


PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M:	-	F:	-
HIV prevalence among pregnant women aged 15-24 in capital city				-
HIV prevalence among most-at-risk populations	FSW:	-	IDU:	-
Knowledge of HIV prevention among young people aged 15-24	M:	-	F:	-
Sex before age 15 among young people aged 15-19	M:	-	F:	-
Condom use at last higher-risk sex among young people aged 15-24	M:	-	F:	-
% of sex workers who used a condom with their most recent client	M:	-	F:	-
% of IDUs who used sterile equipment at last injection				-
% of men who used a condom the last time they had sex with a male partner				-


PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-


Number of HIV+ pregnant women receiving ARVs for PMTCT




Number of children in need receiving ART




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.

* Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Solomon Islands

Estimated adult HIV prevalence rate -

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	-
Estimated number of births per year	15,000
<small>Source: UN Population Division, 2006</small>	
Estimated number of HIV+ pregnant women	-
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	-

PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	-


PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: -	F: -
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-


PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-


Number of HIV+ pregnant women receiving ARVs for PMTCT




Number of children in need receiving ART




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.

* Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Somalia

Estimated adult HIV prevalence rate 0.9%
 Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	26%
Source: MICS, 2006	
Estimated number of births per year	371,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	900
Source: UNAIDS/WHO unpublished estimates, 2007	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	5 0.6%
Source: MOH, 2006	
	[<500 - 1,915]*
	[0.3% - 1.0%]*

PROVIDE PAEDIATRIC HIV TREATMENT



Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	0
Source: MOH, 2006	
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	-

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE


Estimated HIV prevalence among young people aged 15-24	M: 0.2%	F: 0.6%
Source: UNAIDS, 2006		
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: -	F: 3.7%
Source: MICS, 2006		
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS


Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	0.87
Source: MICS, 2006	
% of orphans receiving external support	-




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: South Africa

Estimated adult HIV prevalence rate 16.2%
 Source: UNAIDS/WHO, 2007

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	92%
Source: DHS, 2003	
Estimated number of births per year	1,102,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	222,415 [186,658 - 258,165]*
Source: UNAIDS/WHO unpublished estimates, 2007	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	111,357 50.1% [43.1% - 59.7%]*
Source: MOH, 2006	

PROVIDE PAEDIATRIC HIV TREATMENT



Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	23,369
Source: MOH, 2006	

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE


Estimated HIV prevalence among young people aged 15-24	M: 4.4%	F: 16.9%
Source: National HIV Prevalence, HIV Incidence, Behavior and Communication Survey, 2005		
HIV prevalence among pregnant women aged 15-24 in capital city	25.2%	
Source: National HIV and Syphilis Antenatal Sero-prevalence Survey in South Africa, 2004		
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: -	F: -
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	
% of men who used a condom the last time they had sex with a male partner	-	

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS


Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Sri Lanka

Estimated adult HIV prevalence rate <0.1%
Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	95%
<small>Source: DHS, 2000</small>	
Estimated number of births per year	295,000
<small>Source: UN Population Division, 2006</small>	
Estimated number of HIV+ pregnant women	<100 [<100]*
<small>Source: UNAIDS/WHO unpublished estimates, 2007</small>	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	1 -
<small>Source: MOH, 2006</small>	

PROVIDE PAEDIATRIC HIV TREATMENT






Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	1 -
<small>Source: MOH, 2006</small>	
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	0 -
<small>Source: MOH, 2006</small>	
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	0 -
<small>Source: MOH, 2006</small>	

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	
HIV prevalence among most-at-risk populations	FSW: 0.0%	IDU: -
<small>Source: UNAIDS, 2006</small>		
Knowledge of HIV prevention among young people aged 15-24	M: -	F: -
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: 64.9%
<small>Source: UNAIDS, 2006</small>		
% of IDUs who used sterile equipment at last injection	-	
% of men who used a condom the last time they had sex with a male partner	-	

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Sudan

Estimated adult HIV prevalence rate 1.6%
Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage <small>Source: MICS, 2000</small>	60%
Estimated number of births per year <small>Source: UN Population Division, 2006</small>	1,225,000
Estimated number of HIV+ pregnant women <small>Source: UNAIDS/WHO unpublished estimates, 2007</small>	18,353 [11,981 - 24,788]*
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	- -

PROVIDE PAEDIATRIC HIV TREATMENT






Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	- -
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	- -
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	- -

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: -	F: -
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio <small>Source: MICS, 2000</small>	0.96
% of orphans receiving external support	-



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Suriname

Estimated adult HIV prevalence rate 1.9%
Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage <small>Source: MICS, 2000</small>	91%
Estimated number of births per year <small>Source: UN Population Division, 2006</small>	9,000
Estimated number of HIV+ pregnant women <small>Source: UNAIDS/WHO unpublished estimates, 2007</small>	<100 [<100 - <200]*
Number and % of HIV+ pregnant women receiving ARVs for PMTCT <small>Source: MOH, 2006</small>	35 -

PROVIDE PAEDIATRIC HIV TREATMENT



Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	- -
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	- -
Estimated number of children in need of ART	-
Number and % of children in need receiving ART <small>Source: MOH, 2006</small>	61 -

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE


Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24 <small>Source: MICS, 2006</small>	M: -	F: 41.0%
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS


Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio <small>Source: MICS, 2006</small>	0.97
% of orphans receiving external support	-




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Swaziland

Estimated adult HIV prevalence rate 25.9%
 Source: UNAIDS/WHO, 2007

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	90%
Source: CHS, 2002	
Estimated number of births per year	33,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	13,209
Source: UNAIDS/WHO unpublished estimates, 2007 [11,976 - 14,546]*	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	8,221 62.2%
Source: MOH, 2006 [56.5% - 68.6%]*	

PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	725 -
Source: MOH, 2006	
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	203 -
Source: MOH, 2006	
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	1,155 -
Source: MOH, 2006	


PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: 7.7%	F: 22.7%
Source: UNAIDS, 2006		
HIV prevalence among pregnant women aged 15-24 in capital city	37.3%	
Source: UNAIDS, 2006		
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: 52.3%	F: 52.1%
Source: DHS, 2006-2007		
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: 70.4%	F: 54.2%
Source: DHS, 2006-2007		
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	
% of men who used a condom the last time they had sex with a male partner	-	


PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	0.97
Source: MICS, 2007	
% of orphans receiving external support	41.2%
Source: DHS, 2006-2007	


Number of HIV+ pregnant women receiving ARVs for PMTCT




Number of children in need receiving ART




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Syrian Arab Republic

Estimated adult HIV prevalence rate -

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	84%
<small>Source: MICS, 2006</small>	
Estimated number of births per year	529,000
<small>Source: UN Population Division, 2006</small>	
Estimated number of HIV+ pregnant women	-
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	-

PROVIDE PAEDIATRIC HIV TREATMENT



Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	-

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE


Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: -	F: 7.2%
<small>Source: MICS, 2006</small>		
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS


Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	1.06
<small>Source: MICS, 2006</small>	
% of orphans receiving external support	-




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
* Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Tajikistan

Estimated adult HIV prevalence rate

0.1%

Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	77%
Source: MICS, 2005	
Estimated number of births per year	185,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	<200
Source: UNAIDS/WHO unpublished estimates, 2007	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	4 -
Source: MOH, 2006	

PROVIDE PAEDIATRIC HIV TREATMENT



Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	0 -
Source: MOH, 2006	
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	0 -
Source: MOH, 2006	
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	0 -
Source: MOH, 2006	

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE


Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: 12.1%
Source: UNAIDS, 2006		
Knowledge of HIV prevention among young people aged 15-24	M: -	F: 2.9%
Source: MICS, 2005		
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS


Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	0.92
Source: MICS, 2005	
% of orphans receiving external support	-




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.

* Represents the uncertainty range around the estimates

Rev. 7 March 2008

Children and AIDS fact sheet: Tanzania, United Republic of

Estimated adult HIV prevalence rate 7.0%
 Source: UNAIDS/WHO, 2007

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	78%
Source: DHS, 2004-2005	
Estimated number of births per year	1,589,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	99,775
Source: UNAIDS/WHO unpublished estimates, 2007 [92,091 - 107,699]*	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	14,758 14.8%
Source: MOH, 2006 [13.7% - 16.0%]*	

PROVIDE PAEDIATRIC HIV TREATMENT



Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	3,576
Source: MOH, 2006	

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE


Estimated HIV prevalence among young people aged 15-24	M: 3.0%	F: 4.0%
Source: AIS, 2003-2004		
HIV prevalence among pregnant women aged 15-24 in capital city	8.2%	
Source: UNAIDS, 2006		
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: 40.3%	F: 45.2%
Source: DHS, 2004-2005		
Sex before age 15 among young people aged 15-19	M: 13.0%	F: 11.4%
Source: DHS, 2004-2005		
Condom use at last higher-risk sex among young people aged 15-24	M: 45.5%	F: 33.8%
Source: DHS, 2004-2005		
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	
% of men who used a condom the last time they had sex with a male partner	-	

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS


Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	1.02
Source: DHS, 2004-2005	
% of orphans receiving external support	-




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.

* Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Thailand

Estimated adult HIV prevalence rate 1.4%
 Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	98%
Source: MICS, 2005-2006	
Estimated number of births per year	936,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	5,796 [4,118 - 7,983]*
Source: UNAIDS/WHO unpublished estimates, 2007	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	4,305 74.3% [53.9% - >95.0%]*
Source: MOH, 2006	

PROVIDE PAEDIATRIC HIV TREATMENT



Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	6,298
Source: MOH, 2006	

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE


Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	
HIV prevalence among most-at-risk populations	FSW: 4.3%	IDU: 38.0%
Source: UNAIDS, 2006		
Knowledge of HIV prevention among young people aged 15-24	M: -	F: 46.1%
Source: MICS, 2005-2006		
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	18.8%	
Source: UNAIDS, 2006		
% of men who used a condom the last time they had sex with a male partner	-	

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS


Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	0.99
Source: MICS, 2005-2006	
% of orphans receiving external support	-




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: The former Yugoslav Republic of Macedonia

Estimated adult HIV prevalence rate <0.1%
 Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	81%
<small>Source: Safe Motherhood Needs Assessment, 2000</small>	
Estimated number of births per year	23,000
<small>Source: UN Population Division, 2006</small>	
Estimated number of HIV+ pregnant women	-
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	-

PROVIDE PAEDIATRIC HIV TREATMENT



Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	-

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE


Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: -	F: 26.6%
<small>Source: MICS, 2005-2006</small>		
Sex before age 15 among young people aged 15-19	M: -	F: 0.6%
<small>Source: MICS, 2005-2006</small>		
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: 69.8%
<small>Source: MICS, 2005-2006</small>		
% of sex workers who used a condom with their most recent client	M: 87.5%	F: 84.4%
<small>Source: UNAIDS, 2006</small>		
% of IDUs who used sterile equipment at last injection	-	28.7%
<small>Source: UNAIDS, 2006</small>		
% of men who used a condom the last time they had sex with a male partner	-	29.2%
<small>Source: UNAIDS, 2006</small>		

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS


Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	0.98
<small>Source: MICS, 2005-2006</small>	
% of orphans receiving external support	-




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio




M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Year refers to the year of data, not publication date. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Timor-Leste

Estimated adult HIV prevalence rate -


PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage <small>Source: DHS, 2003</small>	61%
Estimated number of births per year <small>Source: UN Population Division, 2006</small>	46,000
Estimated number of HIV+ pregnant women	-
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	-





PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	-




PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: -	F: -
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-



PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
* Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Togo

Estimated adult HIV prevalence rate 3.2%

Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	84%
Source: MICS, 2006	
Estimated number of births per year	242,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	7,937
Source: UNAIDS/WHO unpublished estimates, 2007 [6,535 - 10,011]*	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	910 11.5%
Source: MOH, 2006 [9.1% - 13.9%]*	

PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	462 -
Source: MOH, 2006	
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	0 -
Source: MOH, 2006	
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	258 -
Source: MOH, 2006	


PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: 0.8%	F: 2.2%
Source: UNAIDS, 2006		
HIV prevalence among pregnant women aged 15-24 in capital city	9.3%	
Source: UNAIDS, 2006		
HIV prevalence among most-at-risk populations	FSW: 53.9%	IDU: -
Source: UNAIDS, 2006		
Knowledge of HIV prevention among young people aged 15-24	M: -	F: 28.3%
Source: MICS, 2006		
Sex before age 15 among young people aged 15-19	M: -	F: 12.4%
Source: MICS, 2006		
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: 49.9%
Source: MICS, 2006		
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	
% of men who used a condom the last time they had sex with a male partner	-	


PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	0.94
Source: MICS, 2006	
% of orphans receiving external support	-


Number of HIV+ pregnant women receiving ARVs for PMTCT




Number of children in need receiving ART




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.

* Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Tonga

Estimated adult HIV prevalence rate -

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	-	-
Estimated number of births per year	3,000	-
<small>Source: UN Population Division, 2006</small>		
Estimated number of HIV+ pregnant women	-	-
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	-	-

PROVIDE PAEDIATRIC HIV TREATMENT



Estimated number of children living with HIV	-	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-	-
Estimated number of children in need of ART	-	-
Number and % of children in need receiving ART	-	-

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE


Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: -	F: -
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS


Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.

* Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Trinidad and Tobago

Estimated adult HIV prevalence rate 2.6%
 Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage <small>Source: MICS, 2006</small>	96%
Estimated number of births per year <small>Source: UN Population Division, 2006</small>	20,000
Estimated number of HIV+ pregnant women <small>Source: UNAIDS/WHO unpublished estimates, 2007</small>	<200 [<100 - <500]*
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	- -

PROVIDE PAEDIATRIC HIV TREATMENT






Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	- -
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	- -
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	- -

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24 <small>Source: MICS, 2006</small>	M: -	F: 53.6%
Sex before age 15 among young people aged 15-19 <small>Source: MICS, 2006</small>	M: -	F: 4.7%
Condom use at last higher-risk sex among young people aged 15-24 <small>Source: MICS, 2006</small>	M: -	F: 51.2%
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio <small>Source: MICS, 2006</small>	1.01
% of orphans receiving external support	-



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Tunisia

Estimated adult HIV prevalence rate 0.1%
 Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	92%
Source: MICS, 2000	
Estimated number of births per year	172,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	<100 [<100]*
Source: UNAIDS/WHO unpublished estimates, 2007	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	0 -
Source: MOH, 2006	

PROVIDE PAEDIATRIC HIV TREATMENT






Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	0 -
Source: MOH, 2006	
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	0 -
Source: MOH, 2006	
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	3 -
Source: MOH, 2006	

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: -	F: -
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Turkey

Estimated adult HIV prevalence rate -

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	81%	
<small>Source: DHS, 2003</small>		
Estimated number of births per year	1,378,000	
<small>Source: UN Population Division, 2006</small>		
Estimated number of HIV+ pregnant women	-	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	4	-
<small>Source: MOH, 2006</small>		

PROVIDE PAEDIATRIC HIV TREATMENT






Estimated number of children living with HIV	-	
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	0	-
<small>Source: MOH, 2006</small>		
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	0	-
<small>Source: MOH, 2006</small>		
Estimated number of children in need of ART	-	
Number and % of children in need receiving ART	9	-
<small>Source: MOH, 2006</small>		

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: -	F: -
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Turkmenistan

Estimated adult HIV prevalence rate

<0.1%

Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	99%
<small>Source: MICS, 2006</small>	
Estimated number of births per year	108,000
<small>Source: UN Population Division, 2006</small>	
Estimated number of HIV+ pregnant women	-
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	0 -

PROVIDE PAEDIATRIC HIV TREATMENT



Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	- -
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	- -
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	0 -
<small>Source: MOH, 2006</small>	

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE


Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: -	F: 4.8%
<small>Source: MICS, 2006</small>		
Sex before age 15 among young people aged 15-19	M: -	F: 0.1%
<small>Source: DHS, 2000</small>		
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS


Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	1.00
<small>Source: MICS, 2006</small>	
% of orphans receiving external support	-




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.


* Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Tuvalu

Estimated adult HIV prevalence rate -


PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	-
Estimated number of births per year <small>Source: UN Population Division, 2006</small>	0
Estimated number of HIV+ pregnant women	-
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	-





PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	-




PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: -	F: -
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-



PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
* Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Uganda

Estimated adult HIV prevalence rate 7.1%
 Source: UNAIDS/WHO, 2007

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	94%
Source: DHS, 2006	
Estimated number of births per year	1,406,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	79,950 [70,513 - 92,116]*
Source: UNAIDS/WHO unpublished estimates, 2007	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	19,991 25.0% [21.7% - 28.4%]*
Source: MOH, 2006	

PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	7,800
Source: MOH, 2006	


PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: 1.1%	F: 4.3%
Source: HIV/AIDS Sero-Behavioural Survey, 2004-2005		
HIV prevalence among pregnant women aged 15-24 in capital city	5.2%	
Source: UNAIDS, 2006		
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: 38.2%	F: 31.9%
Source: HIV/AIDS Sero-Behavioural Survey, 2004-2005		
Sex before age 15 among young people aged 15-19	M: 16.3%	F: 12.2%
Source: HIV/AIDS Sero-Behavioural Survey, 2004-2005		
Condom use at last higher-risk sex among young people aged 15-24	M: 55.1%	F: 52.9%
Source: HIV/AIDS Sero-Behavioural Survey, 2004-2005		
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	
% of men who used a condom the last time they had sex with a male partner	-	


PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	0.94
Source: HIV/AIDS Sero-Behavioural Survey, 2004-2005	
% of orphans receiving external support	10.7%
Source: DHS, 2004	


Number of HIV+ pregnant women receiving ARVs for PMTCT




Number of children in need receiving ART




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Ukraine

Estimated adult HIV prevalence rate 1.4%
 Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	99%
Source: MICS, 2005	
Estimated number of births per year	417,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	3,037 [1,897 - 4,857]*
Source: UNAIDS/WHO unpublished estimates, 2007	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	2,517 82.9% [51.8% - >95.0%]*
Source: MOH, 2006	

PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	800 -
Source: MOH, 2006	
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	168 -
Source: MOH, 2006	
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	650 -
Source: MOH, 2006	


PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: - F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-
HIV prevalence among most-at-risk populations	FSW: 8.0% IDU: 48.8%
Source: UNAIDS, 2006	
Knowledge of HIV prevention among young people aged 15-24	M: - F: 28.2%
Source: MICS, 2005	
Sex before age 15 among young people aged 15-19	M: - F: -
Condom use at last higher-risk sex among young people aged 15-24	M: - F: -
% of sex workers who used a condom with their most recent client	M: - F: -
% of IDUs who used sterile equipment at last injection	18.3%
Source: UNAIDS, 2006	
% of men who used a condom the last time they had sex with a male partner	71.6%
Source: UNAIDS, 2006	


PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	0.98
Source: State Statistics Committee, 2005	
% of orphans receiving external support	-

Number of HIV+ pregnant women receiving ARVs for PMTCT




Number of children in need receiving ART




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.


* Represents the uncertainty range around the estimates

Children and AIDS fact sheet: United Arab Emirates

Estimated adult HIV prevalence rate -


PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	-
Estimated number of births per year	69,000
<small>Source: UN Population Division, 2006</small>	
Estimated number of HIV+ pregnant women	-
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	-



PROVIDE PAEDIATRIC HIV TREATMENT


Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	-




PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: -	F: -
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-

% of young people aged 15-24 who used a condom at last higher-risk sex




HIV prevalence among female sex workers (%)



PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.

* Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Uruguay

Estimated adult HIV prevalence rate

0.5%

Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	-	-
Estimated number of births per year	51,000	-
<small>Source: UN Population Division, 2006</small>		
Estimated number of HIV+ pregnant women	<200	-
<small>Source: UNAIDS/WHO unpublished estimates, 2007</small>		
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	53	-
<small>Source: MOH, 2006</small>		

PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	70	-
<small>Source: MOH, 2006</small>		
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	70	-
<small>Source: MOH, 2006</small>		
Estimated number of children in need of ART	-	-
Number and % of children in need receiving ART	160	-
<small>Source: MOH, 2006</small>		


PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M:	-	F:	-
HIV prevalence among pregnant women aged 15-24 in capital city				-
HIV prevalence among most-at-risk populations	FSW:	-	IDU:	-
Knowledge of HIV prevention among young people aged 15-24	M:	-	F:	-
Sex before age 15 among young people aged 15-19	M:	-	F:	-
Condom use at last higher-risk sex among young people aged 15-24	M:	-	F:	-
% of sex workers who used a condom with their most recent client	M:	-	F:	-
% of IDUs who used sterile equipment at last injection				-
% of men who used a condom the last time they had sex with a male partner				-


PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-


Number of HIV+ pregnant women receiving ARVs for PMTCT




Number of children in need receiving ART




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
* Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Uzbekistan

Estimated adult HIV prevalence rate 0.2%
 Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	99%
Source: MICS, 2006	
Estimated number of births per year	619,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	<500
Source: UNAIDS/WHO unpublished estimates, 2007 [$<200 - 1,041$]*	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	22 -
Source: MOH, 2006	

PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	- -
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	- -
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	52 -
Source: MOH, 2006	


PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: -	F: 31.0%
Source: MICS, 2006		
Sex before age 15 among young people aged 15-19	M: -	F: 0.0%
Source: MICS, 2006		
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: 60.5%
Source: MICS, 2006		
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-


PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	0.95
Source: MICS, 2006	
% of orphans receiving external support	-


Number of HIV+ pregnant women receiving ARVs for PMTCT




Number of children in need receiving ART




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Vanuatu

Estimated adult HIV prevalence rate -

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	-
Estimated number of births per year <small>Source: UN Population Division, 2006</small>	6,000
Estimated number of HIV+ pregnant women	-
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	-

PROVIDE PAEDIATRIC HIV TREATMENT






Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	-

PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: -	F: -
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-

PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
* Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Venezuela (Bolivarian Republic of)

Estimated adult HIV prevalence rate 0.7%
 Source: UNAIDS, 2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	94%
Source: MICS, 2000	
Estimated number of births per year	595,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	2,173 [1,242 - 4,086]*
Source: UNAIDS/WHO unpublished estimates, 2007	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	310 14.3% [7.6% - 25.0%]*
Source: MOH, 2006	

PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	611
Source: MOH, 2006	


PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: -	F: -
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-


PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-


Number of HIV+ pregnant women receiving ARVs for PMTCT




Number of children in need receiving ART




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.

* Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Viet Nam

Estimated adult HIV prevalence rate 0.5%
 Source: AIS, 2005

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	91%
Source: MICS, 2006	
Estimated number of births per year	1,654,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	3,796
Source: UNAIDS/WHO unpublished estimates, 2007	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	1,385 36.5%
Source: MOH, 2006	
	[21.5% - 59.3%]*

PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	428
Source: MOH, 2006	


PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: 0.8%	F: 0.0%
Source: AIS, 2005		
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: 10.0%	IDU: 30.6%
Source: UNAIDS, 2006		
Knowledge of HIV prevention among young people aged 15-24	M: -	F: 43.6%
Source: MICS, 2006		
Sex before age 15 among young people aged 15-19	M: 0.3%	F: 0.5%
Source: AIS, 2005		
Condom use at last higher-risk sex among young people aged 15-24	M: 67.6%	F: -
Source: AIS, 2005		
% of sex workers who used a condom with their most recent client	M: -	F: 90.4%
Source: UNAIDS, 2006		
% of IDUs who used sterile equipment at last injection	-	81.8%
Source: UNAIDS, 2006		
% of men who used a condom the last time they had sex with a male partner	-	-


PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	0.84
Source: MICS, 2006	
% of orphans receiving external support	-


Number of HIV+ pregnant women receiving ARVs for PMTCT




Number of children in need receiving ART




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates


Children and AIDS fact sheet: Yemen

Estimated adult HIV prevalence rate -

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	41%
<small>Source: FHS, 2003</small>	
Estimated number of births per year	839,000
<small>Source: UN Population Division, 2006</small>	
Estimated number of HIV+ pregnant women	-
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	-


Number of HIV+ pregnant women receiving ARVs for PMTCT



PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	-


Number of children in need receiving ART




PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: -	F: -
HIV prevalence among pregnant women aged 15-24 in capital city	-	-
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: -	F: -
Sex before age 15 among young people aged 15-19	M: -	F: -
Condom use at last higher-risk sex among young people aged 15-24	M: -	F: -
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	-
% of men who used a condom the last time they had sex with a male partner	-	-

% of young people aged 15-24 who used a condom at last higher-risk sex




HIV prevalence among female sex workers (%)



PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	-
% of orphans receiving external support	-

Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Zambia

Estimated adult HIV prevalence rate 15.6%
 Source: DHS, 2001-2002

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	93%
Source: DHS, 2001-2002	
Estimated number of births per year	470,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	73,668 [65,911 - 82,984]*
Source: UNAIDS/WHO unpublished estimates, 2007	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	25,578 34.7% [30.8% - 38.8%]*
Source: MOH, 2006	

PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	-
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	-
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	7,200
Source: MOH, 2006	


PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: 3.0%	F: 11.1%
Source: DHS, 2001-2002		
HIV prevalence among pregnant women aged 15-24 in capital city	20.7%	
Source: UNAIDS, 2006		
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: 46.1%	F: 40.5%
Source: SBS, 2005		
Sex before age 15 among young people aged 15-19	M: 27.0%	F: 18.0%
Source: SBS, 2005		
Condom use at last higher-risk sex among young people aged 15-24	M: 38.0%	F: 26.0%
Source: SBS, 2005		
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	
% of men who used a condom the last time they had sex with a male partner	-	


PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	1.03
Source: SBS, 2005	
% of orphans receiving external support	13.4%
Source: SBS, 2005	


Number of HIV+ pregnant women receiving ARVs for PMTCT




Number of children in need receiving ART




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

Children and AIDS fact sheet: Zimbabwe

Estimated adult HIV prevalence rate 18.1%
 Source: DHS, 2005-2006

PREVENT MOTHER-TO-CHILD TRANSMISSION OF HIV

Antenatal care coverage	95%
Source: DHS, 2005-2006	
Estimated number of births per year	372,000
Source: UN Population Division, 2006	
Estimated number of HIV+ pregnant women	59,327
Source: UNAIDS/WHO unpublished estimates, 2007 [54,031 - 64,481]*	
Number and % of HIV+ pregnant women receiving ARVs for PMTCT	10,127 17.1%
Source: MOH, 2006 [15.7% - 18.7%]*	

PROVIDE PAEDIATRIC HIV TREATMENT

Estimated number of children living with HIV	-
Number and % of infants born to HIV+ women receiving cotrimoxazole prophylaxis within 2 months of birth	6,780 -
Source: MOH, 2006	
Number and % of infants born to HIV+ women receiving a virological test for HIV diagnosis within 2 months of birth	0 -
Source: MOH, 2006	
Estimated number of children in need of ART	-
Number and % of children in need receiving ART	4,367 -
Source: MOH, 2006	


PREVENT INFECTION AMONG ADOLESCENTS AND YOUNG PEOPLE

Estimated HIV prevalence among young people aged 15-24	M: 4.2%	F: 11.0%
Source: DHS, 2005-2006		
HIV prevalence among pregnant women aged 15-24 in capital city	18.6%	
Source: UNAIDS, 2006		
HIV prevalence among most-at-risk populations	FSW: -	IDU: -
Knowledge of HIV prevention among young people aged 15-24	M: 45.6%	F: 43.7%
Source: DHS, 2005-2006		
Sex before age 15 among young people aged 15-19	M: 5.2%	F: 4.9%
Source: DHS, 2005-2006		
Condom use at last higher-risk sex among young people aged 15-24	M: 68.0%	F: 42.4%
Source: DHS, 2005-2006		
% of sex workers who used a condom with their most recent client	M: -	F: -
% of IDUs who used sterile equipment at last injection	-	
% of men who used a condom the last time they had sex with a male partner	-	


PROTECT AND SUPPORT CHILDREN AFFECTED BY HIV AND AIDS

Estimated number of children orphaned by all causes	-
Estimated number of children orphaned by AIDS	-
Estimated number of double orphans by all causes	-
Orphan school attendance ratio	0.95
Source: DHS, 2005-2006	
% of orphans receiving external support	31.2%
Source: DHS, 2005-2006	


Number of HIV+ pregnant women receiving ARVs for PMTCT




Number of children in need receiving ART




% of young people aged 15-24 who used a condom at last higher-risk sex



HIV prevalence among female sex workers (%)



Orphan school attendance ratio



M/F = male/female; FSW = female sex workers; IDU = injecting drug users. Dash (-) means data not available.
 * Represents the uncertainty range around the estimates

United Nations Children's Fund
3 United Nations Plaza
New York, NY 10017, USA
Tel.: (+1 212) 326-7000
pubdoc@unicef.org
www.unicef.org

UNAIDS Secretariat
20, avenue Appia
CH-1211 Geneva 27
Switzerland
Tel.: (+41 22) 791-3666
Fax: (+41 22) 791-4187
unaids@unaids.org
www.unaids.org

World Health Organization
20, avenue Appia
CH-1211 Geneva 27
Switzerland
Tel.: (+ 41 22) 791-2111
Fax: (+ 41 22) 791-3111
info@who.int; publications@who.int
www.who.int

Visit the *Unite for Children, Unite against AIDS* website:
www.uniteforchildren.org
or contact us by email: aidscampaign@unicef.org

© The United Nations Children's Fund (UNICEF)
Cover photos: © UNICEF/HQ05-0862/Shehzad Noorani;
© UNICEF/HQ06-1414/Giacomo Pirozzi

ISBN: 978-92-806-4269-8

April 2008